

UNCLASSIFIED

407959

R

UNIVERSITY OF WASHINGTON
LABORATORY OF RADIATION BIOLOGY
SEATTLE, WASHINGTON 98105

ES DOE ARCHIVES
RG _____
Collection <u>1320</u>
Box <u>9</u>
Folder <u>34</u>



Job <u>1320</u>
Box <u>9</u>
Folder <u>34</u>
HISTORIAN'S OFFICE INVENTORY CONTROL

REPORT NO. UWFL-59

UNCLASSIFIED

UWFL-59

Biology and Medicine

STRONTIUM-90 AND GROSS BETA ACTIVITY IN THE FAT AND
NON-FAT FRACTIONS OF COCONUT CRAB (Birgus latro) LIVER
COLLECTED AT RONGELAP ATOLL DURING MARCH 1958

by

Diptiman Chakravarti and Ronald Eisler

Laboratory of Radiation Biology
University of Washington
Seattle, Washington

Lauren R. Donaldson
Director

March 1959

DOE ARCHIVE

Operated by the University of Washington under Contract No.
AT(45-1)540 with the United States Atomic Energy Commission

ABSTRACT

The values for strontium-90 and gross beta activity in the fat and non-fat fractions from the livers of twelve coconut crabs (Birgus latro) collected at Rongelap Atoll during March 1958 are presented.

Although fat constituted an average of 47 percent by weight on a wet weight basis (74 percent on a dry weight basis), gross beta activity of the fat fraction amounted to less than 0.5 percent of the total activity on a wet weight basis. Fat content on a wet weight basis had a range of 31 percent to 65 percent. There is a linear relationship between strontium-90 activity and gross beta activity. Since the fat content of coconut crab liver is variable and the fat fraction contains practically no radioactivity, it is suggested that the radioactivity (and mineral content) of liver samples be compared on the basis of the non-fat solids.

ACKNOWLEDGEMENT

The authors wish to thank Dr. E. E. Held of the Laboratory of Radiation Biology, University of Washington, for his willing assistance and most helpful suggestions in the course of this investigation.

DOE ARCHIVES

STRONTIUM-90 AND GROSS BETA ACTIVITY IN THE FAT AND
NON-FAT FRACTIONS OF COCONUT CRAB (Birgus latro) LIVER
COLLECTED AT RONGELAP ATOLL DURING MARCH 1958

INTRODUCTION

There has been a large individual variability in the levels of radioactivity per unit weight of biological samples collected in the vicinity of the Eniwetok Test Site (1, 2, 3, 4, 5). This variability may be great enough to mask or obscure differences which might exist between species or with time or locality of collection. Practical considerations do not permit, in most cases, increasing the number of samples in an attempt to elucidate possible differences. The work reported in this paper points out a source of variability that exists in comparing the radioactivity of various samples of coconut crab liver.

Birgus latro, the coconut crab, is of particular interest since it is edible and known to concentrate strontium-90. In the course of preparing samples of Birgus latro liver for radioassay, it appeared that the fat content varied considerably from specimen to specimen. A crude determination indicated that the fat contained little or no radioactivity, which was expected because of the low mineral content of fat.

It was therefore decided to determine accurately the fat content and the proportions of strontium-90 and gross beta activity in the fat

DOI-31 1958

and non-fat fractions. The objective was to determine whether more uniform results could be obtained when radioisotopic content was expressed on a basis of non-fat solids rather than total solids as had been done in the past.

MATERIALS AND METHODS

Liver samples of coconut crab (Birgus latro) were collected from Kabelle, Rongelap and Eniaetok Islands at Rongelap Atoll in March 1958. The samples were oven dried at $98 \pm 2^\circ \text{C}$ and partially pulverized. The fat was extracted from the dried samples by a modification of the Johnson method (6). Petroleum ether was used as the extracting solvent. The fat-free solids were wet ashed with concentrated HNO_3 and H_2O_2 . The ash obtained from non-fat solids was dissolved in a known volume of 1 N. HNO_3 and the strontium-90 levels of the samples were determined by the method of Kawabata and Held (7). The gross beta activity was also measured from an aliquot of the solution. The fat content was determined on a dry weight basis by weighing and on a wet weight basis by using the wet weight to dry weight ratios in Table 1. The fat samples were dry-ashed in a muffle furnace at 550°C overnight. The ash obtained from the fat fraction was dissolved in a small volume of 1 N. HNO_3 and was transferred to a plate for determining the gross beta activity. The gross beta activity of the fat fraction was less than one percent of that in

the non-fat fraction, making strontium-90 determinations impractical with the facilities available.

All counting was done with an Anton end-window geiger tube, number 1001-T, which was standardized against a National Bureau of Standards strontium-90 standard.

RESULTS AND DISCUSSION

The ratio of wet to dry weight and the fat content of the liver samples are presented in Table 1. The fat constituted an average of 47 percent by weight on a wet basis with a standard deviation of 9.71, and 74 percent by weight on a dry weight basis with a standard deviation of 9.29. The average ratio of wet to dry weight was 1.603, with a standard deviation of 0.156, indicating that the moisture level of the samples was relatively constant.

Strontium-90 levels expressed as disintegrations per minute per gram of non-fat solids and of total solids given on a dry weight and a wet weight basis are presented in Table 2.

Table 3 presents the gross beta activity in the fat and non-fat fractions of the liver. Although fat constituted an average of 47 percent of the wet weight and 74 percent of the dry weight (Table 1) of the total solids, gross beta activity of the fat fraction amounted to less than 0.5 percent of the total sample on a wet weight basis, and less than 1.0 percent on a dry

weight basis.

The gross beta activity of the samples on a wet and dry weight basis is given in Table 4.

There is a linear relationship between strontium-90 activity and gross beta activity (Tables 2 and 3). The percentage of gross beta activity due to strontium-90 at Kabelle, Eniaetok and Rongelap Islands on a non-fat solid dry weight basis and based upon the average values at each island are 32, 35, and 31 percent respectively.

The strontium units for the liver of coconut crab of earlier collections from Rongelap have been reported previously (4, 5, 8). To report the strontium units for the March 1958 collection, Table 5 has been included.

As a result of these studies, radiochemical analyses of the liver of coconut crab should be made on the basis of non-fat solids rather than the entire liver as has been done in the past.

Table 1. Percentage of fat content and the ratio of wet weight to dry weight of coconut crab (Birgus latro) liver* collected at Rongelap Atoll in March 1958.

Sample number	Percentage of fat in liver		Wet weight Dry weight	Ratio
	Wet	Dry		
34	44.70	78.81	1.763	
35	48.24	74.38	1.542	
36	31.50	55.37	1.758	
37	38.01	67.88	1.786	
38	42.34	67.99	1.606	
61	40.38	71.83	1.779	
62	42.76	73.04	1.708	
84	56.10	83.20	1.483	
85	64.62	87.69	1.357	
86	50.76	71.02	1.399	
87	61.49	88.55	1.440	
88	43.89	70.97	1.617	
Mean	47.07	74.23	1.603	
Standard deviation	9.71	9.29	0.156	

*Sample weights ranged from 9 to 18 grams

DATE RECEIVED

Table 2. Strontium-90 in non-fat solid and total solids of coconut crab (Birgus latro) liver.

Sample number	Location of collection	Non-fat solids d/m/g		Total solids d/m/g	
		Wet	Dry	Wet	Dry
34	Rongelap Atoll	260±10*	458±17*	55±2*	97± 4
35		353±13	544±20	90±3	140± 5
36	Kabelle	276±12	484±20	130±5	288±10
37	Island	605±35	1080±44	194±8	347±14
38		420±20	674±33	134±6	216±11
Mean		383	648	121	218
Standard deviation		140	255	52	103
84		236±10	350±15	40±2	59± 3
85		245±11	332±13	30±1	41± 2
86	Rongelap	159± 8	222±10	46±2	64± 3
87	Island	409±23	589±33	47±3	67± 4
88		224±13	362±21	65±4	105± 6
Mean		255	371	46	67
Standard deviation		93	134	13	23
61	Eniaetok	248± 9	442±17	70±3	124± 5
62	Island	321±20	548±34	87±5	148± 9
Mean		285	495	79	136
Standard deviation		52	75	12	17

*Counting error is less than 5 percent

Table 3. Gross beta activity of fat and non-fat fractions and percentage of gross beta activity in fat of coconut crab (Birgus latro) liver.

Sample number	Location of collection	Fat d/m/g*	Non-fat d/m/g*		Percentage of gross beta activity due to fat	
			Wet weight basis	Dry weight basis	Wet weight basis	Dry weight basis
34	Rongelap Atoll	4	1116	1967	0.29	0.77
35		8	1086	1674	0.67	1.36
36	Kabelle Island	6	618	1086	0.45	0.68
37		14	1929	3446	0.44	0.84
38		2	1219	1958	0.11	0.22
Mean		7	1194	2026	0.39	0.77
84	Rongelap Island	1	753	1117	0.18	0.42
85		0	661	897	0.00	0.00
86		-2	513	718	0.00	0.00
87		1	1169	1684	0.13	0.45
88		1	545	882	0.13	0.26
Mean		0.2	728	1060	0.09	0.23
61	Eniaetok Island	6	824	1466	0.49	0.99
62		6	984	1681	0.45	0.97
Mean		6	904	1573	0.47	0.98

*Counting error is less than 8 percent

NOV 1963

Table 4. Gross beta activity of coconut crab (Birgus latro) liver.

Sample number	Location of collection	Total sample d/m/g*	
		Wet weight basis	Dry weight basis
34	Rongelap Atoll	617	431
35		563	428
36	Kabelle Island	423	485
37		1196	1110
38		702	627
Mean		700	616
Standard deviation		295	288
84		330	188
85	Rongelap Island	234	110
86		253	208
87		450	194
88		306	257
Mean		315	191
Standard deviation		85	52
61		492	413
62	Eniaetok Island	564	454
Mean		528	434
Standard deviation		51	29

* Counting error is less than 8 percent

Table 5. Strontium units and calcium in coconut crab (Birgus latro) liver.

Sample number	Location of collection	Sr ⁹⁰ d/m/g Wet weight basis	mg Ca/gm Wet weight basis	Strontium units*
34	Rongelap Atoll	55	7.25	3448
35		90	10.03	4079
36	Kabelle Island	130	23.93	2469
37		194	15.72	5609
38		134	17.18	3545
Mean		121	14.82	3830
Standard deviation		52	6.51	1152
84		40	8.39	2167
85		30	5.40	2525
86	Rongelap Island	46	12.67	1650
87		47	6.18	3457
88		65	10.99	2688
Mean		46	8.73	2497
Standard deviation		13	3.09	668
61		70	8.00	3977
62	Eniaetok Island	87	7.22	5477
Mean		79	7.61	4727
Standard deviation		12	0.55	1055

*Strontium unit = micro-micro curie of Sr⁹⁰ per gram of calcium.

LITERATURE CITED

1. Held, Edward E. 1957. Land crabs and radioactive fallout at Eniwetok Atoll. U. S. AEC Report UWFL 50. Off. Tech. Services, U. S. Dept. of Commerce.
2. Applied Fisheries Laboratory*, University of Washington. 1949. Bikini radiobiological resurvey of 1948. U. S. AEC Report UWFL-16.
3. Applied Fisheries Laboratory*, University of Washington. 1953. Radiobiological studies at Eniwetok Atoll before and following the Mike shot of the November 1952 testing program. U. S. AEC Report WT-616 (UWFL-33). (Confidential)
4. Applied Fisheries Laboratory*, University of Washington. 1955. A radiological study of Rongelap Atoll, Marshall Islands, during 1954-1955. U. S. AEC Report UWFL-42. Off. Tech. Services, U. S. Dept. of Commerce.
5. Applied Fisheries Laboratory*, University of Washington. 1955. Radiobiological resurvey of Rongelap and Ailinginae Atolls, Marshall Islands, October-November, 1955. U. S. AEC Report UWFL-43. Off. Tech. Services, U. S. Dept. of Commerce.
6. Winton, A. L., and K. B. Winton. 1945. The Analysis of Foods. John Wiley and Sons, New York. 999 pp.
7. Kawabata, T., and E. E. Held. A method for the determination of strontium-90 in biological materials. Applied Fisheries Laboratory*, University of Washington. (unpublished)
8. Dunning, G. M. (Ed.). 1957. Radioactive contamination of certain areas in the Pacific Ocean from nuclear tests. U. S. AEC Report. Supt. of Documents, U. S. Govt. Printing Off.

*Now Laboratory of Radiation Biology