

Sr⁹⁰ in Land Plants
Collected at Kabelle[- March 1958
(Exclusive of pandanus)

Sample	Sample No.	Sr ⁹⁰		mg Ca gm dry sample	μuc Sr ⁹⁰ gm Ca	Limit of detectability
		d/m/g/dry	d/m/g/wet			
Guettarda Leaves	R-78	185.6 ± 6.3	36.7 ± 1.2			
" "	R-48	203.8 ± 8.1	56.0 ± 2.2			
" "	R-64	93.7 ± 4.0	26.0 ± 1.1			
Morinda Leaves	R-45	468.0 ± 18.6	88.6 ± 3.5			
" "	R-76	318.0 ± 12.4	55.3 ± 2.2			
" Fruit	R-46	54.0 ± 2.4	8.7 ± 0.4			
Lepturus Leaves	R-47	167.0 ± 6.4	----	17.016	4500	
" "	R-92	319.3 ± 13.1	154.1 ± 6.3	11.660	12000	
Fimbristylis Leaves	R-58	49.4 ± 2.3				
Tacca Tuber, Central Part	R-93C	54.7 ± 2.6	22.6 ± 1.1	4.212	5900	
Letturus, leaves	R-55	336 ± 1.9	-	15.62	9700	-
Tacca, corn unpeeled	R-93c	2.6 ± 0.4	1.1 ± 0.2	4.212	280	-