



IN REPLYING, ADDRESS THE

DEPARTMENT OF
HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE

Robertson,
Any comments
an answer is needed
2 pages.
by

Division of Special Health Services
Occupational Health Field Headquarters
1014 Broadway, Cincinnati 2, Ohio

409486

January 23, 1957

R

Dr. E. P. Cronkite
Medical Department
Brookhaven National Laboratory
Upton, New York

Dear Gene:

I recently read with great interest, the publication "Some Effects of Ionizing Radiation on Human Beings," TID 5358, July 1956.

In the conclusion of the article it is stated that internal contamination certainly could not have contributed to the acute medical picture. It is stated on page 76 of the article that "the total amount of radioactive material in the G.I. tract at one day post detonation was estimated to be 3 mc in people from Rongelap."

It occurred to me that this may have been a sufficient quantity to have contributed to the gastrointestinal syndrome. I estimate that based on an equation of K. Z. Morgan (page 37 of June 1954 Nucleonics) parts of the G.I. tract would have received over 200 rad. I would appreciate your comments on this matter.

Incidentally, in the event you do not recognize my name, you and I commuted together between Edgewood and Silver Spring, Maryland for a short course some six or seven years ago.

Sincerely yours,

Arthur Wolff, Sr. Veterinarian
Clinic Services

BEST COPY AVAILABLE

It is assumed the average density is 2,200 cm³ of water and 1.29 cm³ of air. Values of T , E , and $\Sigma(RBE)E$ are obtained from Table IV of the Internal Dose Handbook (5). The dose, D , is given by

$$D = \frac{93}{100} \int_{h_0}^{h_1} \frac{W}{7} dt$$

$$= \frac{93/100 I_0 \Sigma(RBE)E}{2.6 \times 10^{-3} \times 7m'} \int_{h_0}^{h_1} e^{-0.693t/T_r} dt \quad (7)$$

$$D = \frac{51.1 I_0 \Sigma(RBE)E (h_1 - h_0)}{m'G} \quad (8)$$

where $G = 1$ when the radioactive half-life $T_r \gg 1$. G is given by

$$G = \frac{0.693(h_1 - h_0)}{(e^{-0.693h_0/T_r} - e^{-0.693h_1/T_r})T_r} \quad (9)$$

If it is assumed the mass of the material, m' , in the critical portion of the GI tract irradiates the intestinal wall from half the solid angle for fraction $h_1 - h_0$ of a day, the maximum permissible concentration in water is given by

$$(MPC)_{GI} = \frac{2I_0}{f_w 2,200} = \frac{1.8 \times 10^{-6} m' DG}{(h_1 - h_0) f_w \Sigma(RBE)E} \quad (10)$$

In this case f_w is the fraction reaching the critical portion of the GI tract and is given by $f_w = (1 - f_1)$ where f_1 is the fraction going from the GI tract to the blood.

Usually, when a radioisotope is inhaled, some of it is swallowed. If a considerable portion enters the blood stream from the GI tract and is deposited in some organ, Eq. 6 must be used to find the (MPC)_a. If a portion of the GI tract receives the largest dose so that it is the critical tissue, Eq. 10 must be modified as follows

$$(MPC)_{GI} = \frac{2,200 f_w}{2 \times 10^7 f_a} (MPC)_{a,GI} = \frac{2 \times 10^{-6} m' DG}{(h_1 - h_0) f_w \Sigma(RBE)E} \quad (11)$$

where $f_a = 0.62(1 - f_1)$ is the fraction retained in the critical GI tissue, and the other terms are as defined previously. In most cases, i.e., where $T_r > 1/2$ day, the lower large intestine

Maximum permissible intake of a radioisotope in a single exposure

Whereas values of maximum permissible body burden and maximum permissible concentration in air and water for the equilibrium condition have been given in the Internal Dose Handbook (5) for some 70 radioisotopes, no maximum permissible single-exposure values have been published by the National Committee on Radiation Protection. Therefore, the single-exposure values published here—in the table—do not have official status. However, many persons are working with radioactive materials under conditions in which an accident could lead to a single exposure; it would be convenient to have available tables of values that apply in such a case. It was with this in mind that an effort was made to determine and list in the table the maximum permissible single-exposure values for the radioisotopes considered in the Internal Dose Handbook (5).

In the case of a single exposure to a radioisotope, the maximum permissible concentration in water (MPC)_w is given by

$$D = \frac{93}{100} \int_{h_0}^{h_1} \frac{W}{7} dt$$

where D = total dose delivered to the critical tissue in units of $\text{mrem} \times \text{cm}^2$ delivered in units of $\text{mrem} \times \text{cm}^2$
 W = total amount of radioisotope taken into body by inhalation or ingestion in a single event (for purposes of calculation assumed to be constant and not more than 1 day)
 maximum permissible concentration in water (MPC)_w and air (MPC)_a are calculated during a single exposure

$$(MPC)_{GI} = \frac{2,200 f_w}{2 \times 10^7 f_a} (MPC)_{a,GI}$$

$$(MPC)_{GI} = \frac{2 \times 10^{-6} m' DG}{(h_1 - h_0) f_w \Sigma(RBE)E}$$