

Project 2.62

RADIOACTIVITY BACKGROUND AND OCEANOGRAPHIC CONDITIONS
IN THE PACIFIC PROVING GROUNDS
AT THE START OF OPERATION REDWING

(A field report based on data collected 5 April - 5
May 1956 on the M/V HORIZON and LCU 1135)

RG 326 US ATOMIC ENERGY
COMMISSION

BEST COPY AVAILABLE

Location LANL
Records Center
Collection TR-4271 H-8 B-192
Folder Rad Background in
Pacific Proving Grounds of Op.
Redwing Project 2.62

Report prepared
10 May 1956

Approved for limited distribution:

by John D. Isaacs
John D. Isaacs

CLASSIFICATION CANCELLED
BY AUTHORITY OF DOE/OC

Carl Johnson 2/10/87
REVIEWED BY DATE
J. Diaz 2/18/87

Assistant to the Director
Scripps Institution of Oceanography

COPIED/DOE
LANL RG

PERSONNEL

Project 2.62 Officer:

Feenan D. Jennings

Participants in collection of data on the M. V. HORIZON:

Paul L. Horrner	Chief Scientist
De Courcey Martin	Radiochemist
Robert L. Wisner	Biologist
Alan C. Smith	Principal Marine Technician
Maxwell Silverman	Senior Geological Technician
Daniel M. Brown	Senior Marine Technician
James Costello	Marine Technician
Jack Lucas	Marine Technician

Participants in collection of data on the LCU 1136:

Donald W. Pritchard	Chief Scientist
Richard A. Schwartzlose	Principal Marine Technician
Robert Brennen	Marine Technician

Participants in data processing included Robert G. Paquette, physical oceanographer, in addition to the two groups listed above.

Credit is due the crews of the M. V. HORIZON and LCU 1136 for the excellent cooperation and assistance given both scientific parties.

ABSTRACT

Background radioactivity and oceanographic conditions in the Pacific Proving Ground are of significance; in evaluating the contaminating effects of REDWING events, in the understanding of the fallout problem at sea and extrapolation to land fallout patterns. Thus, as a part of Project 2.62, a month-long study has been made of radioactivity in water, organisms, and bottom sediments, and of currents and physical character of the water over a 140,000 square mile area around Bikini Atoll.

The waters are slightly radioactive, with values of 150 to 1500 gamma counts per minute per liter (cpm/l), whereas the natural radioactive background, due to potassium-40, is about 94 cpm/l.¹ A field of maximum activity (800 to 1500 cpm/l) exists at 800 to 1200 meters depth at locations to the west of Bikini Atoll, within 150 miles of it, and between $10\frac{1}{2}^{\circ}$ and $13\frac{1}{2}^{\circ}$ N. Radioactivity is associated with particulate matter (possibly organic) at the surface only; at all other depths it is mostly in solution. Organisms collected from the upper layers and deep sea fauna captured in a trawl as deep as 2500 meters depth are about equally radioactive; the level of activity in these marine creatures is about 30 to 50 times as much per unit weight as that in equivalent weight of water. Lagoon waters and surface waters in the open sea around the lagoons are slightly more radioactive than other areas (1100 to 2100 cpm/l).

1. All values of radioactivity given in this report are gamma rays emitted per minute.

COPIED/DOE
LANL RG

The highest ^{level} amount of radioactivity in the area studied is on the ocean ^{& lagoon} floor in the region of CASTLE fallout. The top of one deep sea core taken 10 miles northwest of Bikini Lagoon tested 29,500 cpm/gm; sediment from the northwest end of the lagoon emitted 45,000 cpm/g. Presumably both were from BRAVO event at CASTLE. Elsewhere in the lagoon, the bottom sediment generally emitted from 1500 to 4000 cpm/g compared with a usual 1000 to 3000 cpm/g in the fallout area outside the lagoon.

Sampling of bottom living organisms outside the lagoon is, so far, unsuccessful. Bottom dwelling organisms in Ailinginae and Bikini Lagoons (particularly the latter) were quite radioactive. Other lagoons were not studied. The molluscs displayed the highest activity and this was outstandingly concentrated in the livers and kidneys (52,000 to 84,000 gamma cpm/g).

In the open sea area, measurements of currents verified the general circulation obtained at CROSSROADS and by Japanese cruises, but more explicit information was obtained. Meanderings and eddies dominate the flow in the latitude of Bikini; the main flow westward is located at the north side of the PPG. A large, counterclockwise-revolving eddy at Bikini is found down to depths of at least 500 meters. The current at the surface attains 0.6 to 0.7 knot speed and averages 0.3 to 0.4 knot; thence it decreases to a maximum of 0.3 knot and average of 0.15 knot at 500 meters depth.

The thickness of the wind stirred layer above the thermocline, where temperature decreases rapidly, varies from 40 meters near Bikini to 170 meters in the northwestern part of the area. The

waters are most stratified (i.e., density gradients are greatest) at depths of 50 to 175 meters, being shoalest in the southeastern part of the area, deepest in the northwestern portion, and of intermediate depth at Bikini. Water masses in the area are primarily Equatorial Pacific, North Pacific Intermediate, and West North Pacific Central.

Underwater sound propagation conditions at depths above the thermocline are good, but will become less favorable as summer progresses. Propagation in the deep water is bottom limited where depths are less than 1000 fathoms. The axis of the deep sound channel is found at 800 to 1200 meters (1.3 to 1.5 times as deep as off the U. S. West Coast).

The physical hydrography of Bikini Lagoon is similar to that obtained at CROSSROADS. Surface waters move downwind, and deep waters flow in the opposite direction to complete a cellular circulation. At the upwind end the deep waters are diverted both north and south into two secondary horizontal cells. During moderate to strong trades, water flows into the lagoon along the windward reefs and flows-out through Enyu channel and ^{flows out} at the leeward reefs and channels. With southerly and southeasterly winds most of the inflow occurs through Enyu channel.

Speed of the surface current was measured to be about 1.6% of the wind speed instead of 3% as previously found. The deep flow was apparently more variable in direction and of considerably less speed than that obtained during previous investigations. As the trade winds materially decrease, the surface

current essentially stops immediately, but the deep flow continues for some time.

It is calculated that when the east northeast trades are prevailing, water in the lagoon is completely exchanged on the average every 40 days. At times of light southerly or southeasterly winds an estimated 60 to 100 days are required. No conditions were observed during the field work in which very rapid flushing of the lagoon occurred.

6

CONTENTS

	Page
TITLE PAGE	1
PERSONNEL	2
ABSTRACT	3
CONTENTS	7
LIST OF FIGURES	9
LIST OF TABLES	10
1. INTRODUCTION	11
2. RADIOACTIVITY BACKGROUND IN THE OPEN SEA AND LAGOONS OF THE PPG AREA by DeCoursey Martin, Jr. and Robert L. Wisner	13
2.1 Assay Equipment	13
2.2 Sample preparation and counting	14
2.3 Radioactivity in ocean water	15
2.4 Radiobiological assay of the near surface and deep sea fauna	18
2.5 Radioactivity background of the ocean floor	22
2.6 Radiobiological assay of certain fauna of Bikini and Ailinginae Lagoons	24
2.7 Lagoon water samples	26
2.8 Lagoon bottom samples	28
2.9 Lagoon Phytoplankton samples	29
2.10 Miscellaneous surveys	29
2.11 Air sampling	30
3. OCEANOGRAPHIC CONDITIONS IN THE PPG OPEN SEA AREA, by Paul L. Horrer	31
3.1 Currents, general information	31
3.2 Currents at various levels	32

	Page
3.3 Temperature and density	35
3.4 Mixing processes and stability	37
3.5 Water masses in the area	40
3.6 Underwater sound propagation	44
4. CIRCULATION IN BIKINI LAGOON, by Donald W. Pritchard	46
4.1 Discussion of water movements in Bikini Lagoon	46
4.2 Results of present field studies	48
4.3 Conditions during periods of light winds	50
4.4 The renewal of water in Bikini Lagoon	51
4.5 Movement and flushing of radioactive contaminated waters in the Lagoon	54
REFERENCES	61
DISTRIBUTION	63
FIGURES	

LIST OF FIGURES

- 1A. Positions of observations.
- 1B. Positions of observations.
2. Bottom sediment samples, gamma radioactivity.
3. Levels of radioactivity in a sediment core of the PPG.
4. Bikini Lagoon bottom samples, gamma radioactivity.
5. Surface currents from geomagnetic-electro-kinetograph.
- 6A. Direction and speed of surface currents.
- 6B. Direction and speed of currents at 75 meters depth.
- 6C. Direction and speed of currents at 150 meters depth.
- 6D. Direction and speed of currents at 250 meters depth.
- 6E. Direction and speed of currents at 500 meters depth.
7. Surface temperatures.
8. Average temperature in the upper 250 meters of depth.
9. Example of vertical distributions of physical data.
10. Depth of thermocline.
11. Stability of the most stable 25 meter layer.
12. Depth of the most stable 25 meter layer.
13. Average stability in the upper 250 meters of depth.
14. Depth (in meters) of various water masses.
15. Observed currents in the upper layer of Bikini Lagoon.
16. Observed currents in the lower layer of Bikini Lagoon.
17. Current velocities - a five hour set of observations near the center of the lagoon.
18. Current velocity, 30 meter drift drogues.

DRAFT

~~CONFIDENTIAL~~
LIST OF TABLES

	Page
1A. Radioactivity of PPG open sea area waters.	16
1B. Radioactivity of lagoon waters.	16
2. Radioactivity assay of plankton samples from the PPG area.	19
3. Radioactivity analysis of some deep-sea fauna near Bikini Atoll.	21
4. Radioactivity levels of certain fauna from Bikini Lagoon.	25
5. Radioactivity levels of certain fauna from Ailinginae Lagoon.	27
6. Estimates of flow into and out of Bikini Lagoon.	53