

age of the population shown in Table V-4 of BEIR-III.

2. For estimates of cancer risk both the relative risk coefficient and the absolute risk coefficient were used to give a range of estimated risk. The absolute risk coefficient gives a lower value, is less variable with the population and is not dependent upon the spontaneous cancer incidence, which is not known for the Bikini population. The relative risk coefficient gives a high value, but since it is based on the spontaneous cancer incidences, which is unknown for the Bikini population, it is probably less reliable than the estimates calculated from the absolute risk coefficients.

3. For estimating increased cancer incidences, the bone marrow dose was used because it was slightly higher than the whole body dose. This probably introduced a small element of conservatism.

4. For estimating birth defects neither BEIR-I or BEIR-III is very clear about what is meant by parental dose, thus it is not clear whether birth defects should be based on the dose to one parent or both parents. In the latter case, the 30-year whole body dose would be doubled. We assumed the BEIR-I risk of 0.2% rem was based on both parents being irradiated. Also because we believed the risk coefficient from BEIR-I

males and 49 females. These values are tabulated in the appendix.

7. The spontaneous incidence of birth defects was taken to be 10.7% of all live births from BEIR-III.

8. The normal incidence of cancer deaths was assumed to be 15%. A value less than the approximately 20% given for the U.S. population was used because the Bikini people have been and will probably be exposed to much lower limits of environmental carcinogens than people living in the U.S. and because of limited medical services and prevalence of other risks such as drowning, poisoning, etc. Other causes of death are probably higher in the Bikini population than in the U.S. population. We also suspected the average life span was less than in the U.S. population, which might tend to reduce the number of cancers that would occur in the elderly.

9. The largest dose a person might receive in a year was estimated to be three times the average dose. Data in the appendix for individuals show that the highest individual dose is more than twice the average but less than three times.

II. POPULATION ESTIMATES

To estimate the number of births, deaths and the magnitude of the Bikini population after 30 years, information was used from the final draft of the Marshall Islands Five Year Health Plan prepared by the Trust Territories' Department of Health Services' Office of Health Planning and the Resources Department. The document is undated, but the presence of data from 1976 indicates that it must have been prepared in the period of 1977 to 1979 when we received it. It was noted that there are apparent inconsistencies among several of the different tables. For example, Table III-1 gives data for the Marshall Islands for the period 1955-1975 and Table III-5 gives data for the infant mortality rate for 1976. In Table III-1, the infant death rate per 1000 births for 1970 through 1975 is given as 28.3, 33.6, 25.4, 46.4, 21.1 and 37.0. However, Table III-5 indicates the infant mortality rate to be only 17.04. We used the data of Table III-1 in the following estimates; because it is more complete and it provides a self-consistent set of data. However, in view of the discrepancies, the results can only be considered as approximations. This probably makes little real difference in view of the uncertainties in the risk coefficients that were used. There is also a bias built into the data because of the inclusion of Ebye and Majuro in the overall Marshall Island rates. This arises from the different death rates (particularly infants) at these two locations. In many respects the population of Ebye and Majuro are quite dissimilar from the Bikini population because they have the advantages and disadvantages of a more technical environment.

For the estimates the last 5 or 6 year average of the data were used because they are probably the most representative of current conditions. From this, the following were obtained:

1. Rate of increase of the population has been about 3.8%/year.
2. Infant death rate is about 3.2% per birth.
3. Overall death rate is 0.54% per year.
4. Birth rate is 4.2% per year.

A population of 550 was assumed for the one that might move back permanently to Bikini Atoll. Values for other initial populations were obtained by ratios of the results.

The total population at the end of 30 years is given by the compounding equation:

$$P_{30} = 550 (1 + 0.038)^{30} = 1684$$

The number of births in 30 years are given by:

$$B = 0.042 \times 550 \int_0^{30} (1.038)^x dx$$

where x is the time between 0 and 30. This gives

$$B = \frac{0.042 \times 550}{\ln 1.038} [1.038^{30} - 1] = 1277$$

Similarly, the number of deaths in the 30 year period would be:

$$\text{Deaths} = 0.0054 \times 550 \int_0^{30} (1.038)^x dx$$

$$\text{Deaths} = \frac{0.0054 \times 550}{\ln 1.038} [1.038^{30} - 1] = 164$$

One other datum needed is the reduction in 30 year dose to those born after the return because of the decrease in radiation levels and the smaller amount of time in the 30 year period that is spent on the island. For this, the total population dose for those born after returning assuming an initial dose rate of 1 rad/year is given by:

$$P = 550 D_1 \int_0^{30} e^{-\lambda x} (1.038^x) dx$$

λ is the half-life of decrease of the radiation dose, taken here as 30 years.

Because this integral cannot be solved analytical, an approximate solution was obtained by calculating this function for each of 30 years and

The assumption of no deaths in the original 550 returning was made for simplicity and the lack of good death rate data.

We also compared the age characteristics of the Marshallese from Table IV-3 and the U.S. population in 1970. This comparison is given in the attached curve. The slopes are similar above age 35 but the magnitudes are distorted by the high birth rate in the Marshall Islands. However, in terms of the relative risk the similar slopes suggest that if the natural cancer rates in the two populations are similar, the relative risk for people above 35 in both populations would be similar because most of the cancer occurs at ages from about 40 and above. However, the magnitude of the relative risk in the U.S. used for the Marshallese will be high by a factor of somewhere around 2-3 because of the distortion caused by the very high proportion of young people who have a relatively low natural cancer incidence.


Using the preceding calculations for a population of 550, calculations were made for other population sizes. For a population of 550 (from preceding):

Deaths in 30 years = 164 \approx 160

Births in 30 years = 1277 \approx 1300

For a population of 140 (the number that returned to Bikini):

COMPARISON OF AGE CHARACTERISTICS
OF MARSHALLESE AND 1970 US POPULATIONS



Deaths in 30 years $\frac{164}{550} = \frac{x}{140}$, $x = 41.7 \approx 40$

Births in 30 years $\frac{1277}{550} = \frac{x}{140}$, $x = 325. \approx 300$

For a population of 235:

Deaths in 30 years $\frac{164}{550} = \frac{x}{235}$, $x = 70.07 \approx 70$

Births in 30 years $\frac{1277}{550} = \frac{x}{235}$, $x = 545.62 \approx 550$

For a population of 350:

Deaths in 30 years $\frac{164}{550} = \frac{x}{350}$, $x = 104.36 \approx 100$

Births in 30 years $\frac{1277}{550} = \frac{x}{350}$, $x = 812.63 \approx 800$

III. RISK COEFFICIENTS

At the time the Bikini book was prepared no agency in the U.S. government had accepted the risk coefficients in BEIR-III. Thus we were constrained to use risk coefficients from BEIR-I. While not included in the printed book, risk estimates based on BEIR-III were calculated for comparison purposes. The following gives the origin of the risk coefficients used.

A. BEIR-I

1. Cancer (Tables 3-3 and 3-4)

Cancer deaths/year in U.S. from 0.1 rem/year (pop = 197,863,000)	<u>Derived</u> Cancer deaths/ 10^6 person rem
--	---

	<u>Absolute</u>	<u>Relative</u>	<u>Absolute</u>	<u>Relative</u>
Leukemia	516	738	26	37
Other Cancers				
30 year	1210	2436	61	123
elevated risk				
lifetime	1485	8340	75	421
elevated risk				

20X10
 13-20
 1/10/10/10-100

1/10/10/10-100

Range 1726-2001 3174-9078 87-101 160-458

From the above the minimum estimate of cancer risk would be given by a risk coefficient of $87/10^6$ person rem and the maximum by $458/10^6$ person rem. Thus, these two risk coefficients were used to define a range of estimated cancer deaths.

2. Genetic Effects (from Page 1 & 2 BEIR-I)

a. Based on specific defects 5 rem/30 year reproductive generation would cause in the first generation 100-1800 cases of dominant diseases and defects per year (3.6 million births/year) or 5 times this amount at equilibrium. The 1800 cases represent an increase of 0.05% incidences per year first generation and 0.25% at equilibrium. In addition there would be a few chromosomal defects and recessive diseases and a few congenial defects due to a single gene defect and chromosome aberrations.

The total incidence at equilibrium is 1100 to 27,000/year. These at equilibrium, the maximum would be 0.75% or 0.15% in the first generation.

These are equivalent to 0.15% per rem at equilibrium and 0.03%/rem in the first generation.

b. Based on overall ill health. Overall ill health: 5% - 50% of ill health is proportional to the mutation rate using 20% and doubling dose of 20 rem, 5 rem per generation would eventually lead to a 5% increase in ill health.

Thus the rate of overall ill health is 1%/rem at equilibrium or 0.2%/rem in first generation.

For estimating the potential genetic derived health defects in the Bikini population it was decided to use a risk coefficient of 0.2% per rem in the first generation recognizing that it was probably very conservative.

B. BEIR-III

1. Cancer (Table V-4 of Typescript Edition)

Lifetime Risk of Cancer Death
(deaths/10⁶/rad)

<u>Model</u>	<u>Single exposure to 10 rad</u>		<u>Continuous exposure to 1 rad/yr</u>	
	<u>Absolute</u>	<u>Relative</u>	<u>Absolute</u>	<u>Relative</u>
L-Q, $\overline{LQ-L}$	77	226	67	182*
L-L, $\overline{L-L}$	167	501	158	430*
Q-L, $\overline{Q-L}$	10	28	---	---

* In printed version these were 169 and 403, respectively. We used the risk coefficients that were derived for continuous exposure.

2. Birth Defects--pages 166-169 (mean parental age = 30 years)
1 rem per generation (1 rem parental exposure) per 10⁶ live offspring 5 to 75 birth defects, this is 0.0005--0.0075%--First generation.

Since the spontaneous rate is given as 10.7%, in the U.S. population, 1 rem will increase the rate from 10.7% to 10.7005--10.7075%.

In terms of the spontaneous rate 1 rem per generation gives $\frac{0.0005}{10.7} = 0.000047 = 0.0047\%$ increase and $\frac{0.0075}{10.7} = 0.0007 = 0.07\%$ increase.

IV. CALCULATIONS OF RISK

Table 1 gives the radiation dose values provided by Dr. Robison for use in developing estimates of increased health risks in the Bikini population.

A. Risks for 14 Different Living Conditions

1. Cancer Risks

Table 3 shows the calculations for estimates of increased cancer risk for 14 different living conditions.

2. Birth Defects Risks

Table 3 gives the calculations for the estimates of birth defects.

B. Risk Estimates Based on BEIR-III

Table 4 gives risk estimates based on BEIR-III risk coefficients. These were calculated for comparison purposes only and were not used in the Bikini book. The highest estimates for cancer risk result from using the linear relative risk model and are about the same as those given in Table 2 for the relative risk model. The lowest estimates result from the linear-quadratic absolute risk model and are slightly less than those for the absolute model in Table 2. Thus, as far as estimates of cancer risk are concerned, those obtained using risk coefficients from BEIR-I are in the same general range as those obtained using risk coefficients from BEIR-III.

Risk estimates for birth defects obtained using the risk factor from BEIR-I gives values about three times those obtained using the upper value of the range of risk factors given in BEIR-III. If BEIR-III risk factors for birth defects represent a more enlightened assessment of this potential consequence of radiation exposure than the factor taken from BEIR-I for overall health defects, then the estimates in the Bikini book may be conservative by a factor of three.

Females

Identification Number	Age	Total Whole Body Dose (mrem)
6111	32	250
6097	19	950
6115	43	1600
6109	15	760
6091	13	1300
6046	43	600
6061	32	1400
6122	70	1600
6030	10	1600
6129	13	850
6027	6	1200
6010	8	2000
6105	5	1500
6059	19	400
6124	54	390
6058	18	1200
6036	27	340
6110	32	1400
6051	19	1200
6092	8	2400 (highest value)
6080	7	310
6038	6	1400
6103	9	1600
6028	7	1800
6044	6	2200
6062	21	1100
6034	46	1800
865	45	1300
6050	22	710
6094	10	2100
6112	35	420
6035	20	1400
6045	28	270
6108	24	730
6063	24	1100
525	37	470
934	43	2100
6106	6	1100
6025	5	1300
6113	25	880
6060	22	790
6032	32	1400
6123	50	1000
6098	16	720
6065	19	910
6114	32	290
6064	30	1300
6081	9	610
6048	13	660

44,320 (Total for 41 under age 40)

Average = 1080.98 mrem

Total for all 49 females = 54,710

Average = 1116.55 mrem

APPENDIX

Estimates of Radiation Doses Received By Person Who Visited at Bikini for About 10 Years Until August 1978

A. Bone Marrow Doses - Calculation of Average Dose (Values in mrem)

<u>Male</u>		<u>Female</u>	
1600	2600	260	430
1600	1600	1000	1500
300	710	1700	280
1300	510	810	770
1200	2100	1400	1100
1300	1800	700	480
1600	680	1500	2200
890	500	1700	1200
2400	1100	1600	1300
1300	350	900	900
1500	2700	1200	820
1900	1600	2100	1400
900	210	1500	1100
2100	2100	410	760
310	1400	400	1000
1500	1900	1300	300
370	1600	340	1400
1300	1900	1500	620
2300	1600	1200	<u>670</u>
1900	3000 (highest value)	2400	56,200 mrem
1600	<u>72,360 mrem</u>	320	
480	n = 50	1400	n = 49
1800		1600	
2000		1900	
2500		2300	Average dose to all people
2300		1100	
1900		1900	72.36 rem
590		1400	<u>56.20 rem</u>
1500		740	128.56
2600		2200	<u>128.56</u>
			<u>99</u> = 1.2986 = 1.3 rem
			per person

B. Whole Body Dose

Males

Identification Number	Age	Total Whole Body Dose (mrem)
6001	66	1400
6127	13	1500
6130	29	300
6076	39	1300
813	23	1200
6019	48	1100
6132	12	1500
6066	32	830
6070	28	2200
6118	22	1200
6117	22	1400
6128	31	1800
6015	11	870
6033	27	2000
6007	35	300
6008	32	1400
6071	32	350
863	27	1200
6086	46	2100
6067	32	1700
6073	24	1400
6072	20	460
6119	17	1700
864	51	1900
966	56	3200 (highest value)
6009	6	2200
6049	8	1900
6042	7	580
6014	5	1500
6012	7	2400
6016	10	2400
6013	5	1600
6005	38	700
6135	35	500
6125	35	2100
6067	56	1700
6002	65	670
6006	37	490
6096	48	1100
80	69	330
6017	49	2300
6058	56	1500
6004	28	200
6018	34	1900
6126	35	1400
6003	22	1700
6023	8	1500
6131	14	1800
6011	11	1400
6133	11	2800

Total for all 50 males = 70,530

Average = 1410.6 mrem

53,230 (Total for 39 under age 40)

Average = 1364.87 mrem

ESTIMATED RADIATION DOSES TO RESIDENTS OF ENEU AND/OR BIKINI ISLANDS

RESIDENTS OF BIKINI ISLANDS

Residence Island	Years on/ Years off	Time on Eneu (%)	Bi
Eneu	Permanent	100	1
Eneu	Permanent	100	1
Bikini	Permanent	0	1
Bikini	Permanent	0	1
Eneu	Permanent	90	1
Eneu	Permanent	90	1
Eneu	1/1	100	1
Eneu	1/1	100	1
Eneu	1/1	90	1
Eneu	1/1	90	1
Eneu	1/2	100	1
Eneu	1/2	100	1
Eneu	1/2	90	1
Eneu	1/2	90	1
Eneu	1/3	100	1
Eneu	1/3	100	1
Eneu	1/3	90	1
Eneu	1/3	90	1

Imported Food
(50% of Diet)

Yes
No

Yes
No

Yes
No

Yes
No

Yes
No

Yes
No

Yes
No

Yes
No

Yes
No

* Doses are rounded off.

** Numerical value given is three times the a

Table 2
CANCER RISKS

1	2	3	4	5	6	7	8	9	10	Natural Rates	
Living Conditions	Initial Population	30-Yr Bone Marrow Dose (rem)	30-Yr Person (rem)	# of Births Expected in 30 Yr	30-Yr Dose (0.36 x Col. 3)	30-Yr Additional Person (rem)	Total Person (rem)	Absolute Risk *	Relative Risk +	Deaths	Cancer 15A
EU-100%	550	3.0	1650	1300	1.08	1404	3054	.266	1.399	160	24
. Imported food	550	6.0	3300	1300	2.16	2808	6108	.531	2.797	160	24
. No imported food											
KINI-100%	550	25.	13750	1300	9.0	11700	25450	2.214	11.66	160	24
. Imported food	550	47.	25850	1300	16.92	21996	47846	4.16	21.92	160	24
. No imported food											
EU-330 days	550	3.4	1870	1300	1.224	1591	3461	.301	1.585	160	24
KINI-35 days	550	6.5	3575	1300	2.340	3042	6617	.576	3.03	160	24
. Imported food											
. No imported food											
NEU-1 year on and 1 year off	350	1.5	525	800	.54	432	957	.083	.438	100	15
. Imported food	350	3.1	1085	800	1.116	892.8	1978	.172	.906	100	15
. No imported food											
NEU-330 days	350	1.7	595	800	.612	489.6	1085	.094	.497	100	15
KINI-35 days year on and 1 year off	350	3.3	1155	800	1.188	950.4	2105	.183	.964	100	15
. Imported food											
. No imported food											
NEU-1 year on and 2 years off	235	1.03	242	550	.371	203.9	446	.0388	0.204	70	11
. Imported food	235	2.1	494	550	.756	415.8	909.8	.0792	.417	70	11
. No imported food											
NEU-330 days	235	1.2	282	550	.432	237.6	520	.045	.238	70	11
KINI-35 days year on and 2 years off	235	2.2	517	550	.792	435.6	952.6	.083	.436	70	11
. Imported food											
. No imported food											

* 87×10^{-6} per person rem

+ 458×10^{-6} per person rem

Table 3

BIRTH DEFECTS

Sept. 10, 1980

1	2	3	4	5	6	7
Living Conditions	Initial Population	# of Births in 30 Yr	Spontaneous Birth Defects (10.7%)	30-Yr Whole Body Dose (rem)	% Increase (0.2%/rem)	No. of Increase Birth Defects*
1	550	1300	139.1+140	2.8	.56	.78
2	550	1300	139.1+140	5.4	1.08	1.51
3	550	1300	140	24	4.8	6.72
4	550	1300	140	44	8.8	12.32
5	550	1300	140	3.2	.64	.896
6	550	1300	140	5.9	1.18	1.65
7	350	800	85.6+90	1.4	.28	.252
8	350	800	85.6+90	2.8	.56	.50
9	350	800	90	1.6	.32	.288
10	350	800	90	3.0	.6	.54
11	235	550	58.85+60	.96	.192	.1152
12	235	550	58.85+60	1.9	.38	.228
13	235	550	58.85+60	1.1	.22	.132
14	235	550	58.85+60	2.0	.4	.24

SAME AS IN Table 1

* Values were rounded for use in the Bikini book.

Table 4

Risk Estimates Based on BEIR-III

	Cancer					Birth Defects					% Increase***
	Total Person rem	Relative Risk		Absolute		Number of Births in 30 Yrs	30-Yr Whole Body Dose (rem)	Birth Defects (5-75/10 /rem)	Spontaneous Number		
		L-Q 182*	L-L 430	L-Q 67	L-L 158						
1	3054	.556	1.31	.205	.483	1300	2.8	.0182-.27**	139	0.19	
2	6108	1.11	2.63	.409	.965	1300	5.4	.035-.527	139	0.38	
3	25450	4.63	10.94	1.71	4.02	1300	24.0	.156-2.34	139	1.68	
4	47846	8.71	20.57	3.21	7.56	1300	44.0	.286-4.29	139	3.1	
5	3461	.63	1.49	.23	.547	1300	3.2	.021-.312	139	0.22	
6	6617	1.20	2.85	.44	1.05	1300	5.9	.038-.575	139	0.414	
7	957	.174	.41	.064	.15	800	1.4	.0056-.084	85.6	0.098	
8	1978	.36	.851	.133	.313	800	2.8	.011-.168	85.6	0.196	
9	1085	.197	.467	.073	.17	800	1.6	.0064-.096	85.6	0.112	
10	2105	.383	.905	.141	.33	800	3.0	.012-.18	85.6	0.21	
11	446	.081	.192	.0298	.0705	550	.96	.0026-.04	58.85	0.068	
12	910	.166	.39	.061	.144	550	1.9	.0052-.078	58.85	0.13	
13	520	.095	.224	.035	.082	550	1.1	.0030-.045	58.85	0.076	
14	953	.173	.41	.064	.151	550	2.0	.0055-.0825	58.85	0.14	
									9	10	

* Risk Coefficient

182 x 10⁻⁶ man rem** eg. $\frac{2.8 \text{ rem} \times 5 \times 1300 \text{ births}}{10^6}$

*** Based on highest value in Column 8.

September 10, 1982

MI

CALCULATION OF POTENTIAL RADIATION CAUSED HEALTH EFFECTS
FOR PERSONS LIVING IN THE NORTHERN MARSHALL ISLANDS

Potential health effects for persons living in the northern Marshall Islands are calculated using the same assumptions and same methods used for the Bikini population (copy attached). Risk coefficients from both BEIR I and BEIR III were used providing not only a range of estimates but also a comparison of the most conservative (linear, relative risk model) with what would be described by many radiation biologists as the most probable (linear-quadratic, absolute model).

POPULATION ESTIMATES

The following population estimates are derived by simple ratios from the Bikini calculation (copy attached) for a population of 550. These calculations predicted 1277 births, 164 deaths over a period of 30 years and a final population of 1684 after 30 years for an initial population of 550.

$$\text{Deaths in 30 years: } \frac{164}{550} = \frac{\text{deaths in population of interest}}{\text{initial population of interest}}$$

$$\text{Births in 30 years: } \frac{1277}{550} = \frac{\text{births in population of interest}}{\text{initial population of interest}}$$

$$\text{Population after 30 years: } \frac{1684}{550} = \frac{\text{population after 30 years}}{\text{initial population of interest}}$$

Also from the Bikini population, the estimate of the full 30 year dose received by children born during the 30 year period is 0.36 of the dose persons living the entire 30 year period would receive.

September 10, 1982

RISK COEFFICIENTS

Both BEIR I and BEIR III risk coefficients are used. These are as follows:

BEIR I

Cancer--Minimum: Absolute risk of leukemia ($26 \times 10^{-6} \text{ rem}^{-1}$) + 30 year elevated risk for other cancers ($61 \times 10^{-6} \text{ rem}^{-1}$) = $87 \times 10^{-6} \text{ rem}^{-1}$.

Maximum: Relative risk of leukemia ($37 \times 10^{-6} \text{ rem}^{-1}$) + lifetime elevated risk ($421 \times 10^{-6} \text{ rem}^{-1}$) = $458 \times 10^{-6} \text{ rem}^{-1}$.

Genetic Effects: 0.2% per rem in first generation.

BEIR III

Cancer--Minimum: Absolute lifetime risk of cancer for continuous exposure, $67 \times 10^{-6} \text{ rad}^{-1}$ (low LET) based on linear quadratic model.

Maximum: Relative lifetime risk of cancer for continuous exposure, $430 \times 10^{-6} \text{ rad}^{-1}$, based on linear model.

Genetic Effects--Minimum: 75×10^{-6} increase per rem in first generation.

Maximum: 5.0×10^{-6} increase per rem in first generation.

August 5, 1980

100% IMPORTED FOOD	ENEU X	BIKINI IMPORTED FOOD	ENEU 330 IMPORTED FOOD	BIKINI 35 X	ENEU IMPORTED FOOD	1 YEAR ON 1 YEAR OFF X
--------------------	--------	----------------------	------------------------	-------------	--------------------	------------------------

Max. Annual

WB	124	242	1065	1974	88	168
BM	130	261	1094	2063	92	181

3X Max. Annual

WB	372	726	3195	5922	264	504
BM	390	783	3282	6189	276	543

30 Year

WB	2800	5400	24,000	44,000	1400	2800
BM	3000	6000	25,000	47,000	1500	3100

1 YR ENEU 330 DAYS BIKINI 35 DAYS 1 YEAR OFF IMPORTED FOOD	X	1 YEAR ENEU 2 YEARS OFF IMPORTED FOOD	X	1 YR ENEU 330 DAYS BIKINI 35 DAYS 2 YEARS OFF IMPORTED FOOD	X	1 YEAR ENEU 3 YEARS OFF IMPORTED FOOD	X	1 YR ENEU 330 DAYS BIKINI 35 DAYS 3 YEARS OFF IMPORTED FOOD	X
--	---	---------------------------------------	---	---	---	---------------------------------------	---	---	---

Max. Annual

WB	106	186	88	168	107	186	107	186
BM	110	198	92	179	110	197	92	179

3X Max.

WB	318	558	264	504	321	558	264	504
BM	330	594	276	537	330	594	276	537

30 Year

WB	1600	3000	960	1900	1100	2000	760	1500	860	1600

B

100% IMPORTED FOOD	ENEU X	BIKINI IMPORTED FOOD	ENEU 330 IMPORTED FOOD	BIKINI 35 X	ENEU IMPORTED FOOD	1 YEAR ON 1 YEAR OFF X
--------------------	--------	----------------------	------------------------	-------------	--------------------	------------------------

Max. Annual

WB	124	242	1065	1974	88	168
BM	130	261	1094	2063	92	181

3X Max. Annual

WB	372	726	3195	5922	264	504
BM	390	783	3282	6189	276	543

30 Year

WB	2800	5400	24,000	44,000	1400	2800
BM	3000	6000	25,000	47,000	1500	3100

1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD
---	---	---	---	---	---	---

Max. Annual

WB	106	186	88	168	107	186	107	168	107	186
BM	110	198	92	179	110	197	92	179	110	197

3X Max.

WB	318	558	264	504	321	558	264	504	321	558
BM	330	594	276	537	330	594	276	537	330	594

30 Year

WB	1600	3000	960	1900	1100	2000	760	1500	860	1600
BM	1700	3300	1030	2100	1200	2200	810	1700	920	1800

100% IMPORTED FOOD	ENEU X	BIKINI IMPORTED FOOD	ENEU 330 IMPORTED FOOD	BIKINI 35 X	ENEU IMPORTED FOOD	1 YEAR ON 1 YEAR OFF X
--------------------	--------	----------------------	------------------------	-------------	--------------------	------------------------

Max. Annual

WB	124	242	1065	1974	88	168
BM	130	261	1094	2063	92	181

3X Max. Annual

WB	372	726	3195	5922	264	504
BM	390	783	3282	6189	276	543

30 Year

WB	2800	5400	24,000	44,000	1400	2800
BM	3000	6000	25,000	47,000	1500	3100

1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	X	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	X	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD	X	1 YR ENEU 330 DAYS BIKINI 35 DAYS IMPORTED FOOD
---	---	---	---	---	---	---

Max. Annual

WB	106	186	88	163	107	186
BM	110	198	92	179	110	197

3X Max.

WB	318	558	264	504	321	558
BM	330	594	276	537	330	594

30 Year

WB	1600	3000	960	1900	1100	1600
BM	1700	3300	1030	2100	1200	1800

picocuries per gram

These numbers are 0-5 centimeters soil increment

(dictated by W. L. Robison, Lawrence Livermore, August 6, 1980)

B

ENEWETAK ISLANDS		CESIUM	STRONTIUM	PLUTONIUM	AMERICIUM
ALICE	Bokoluo	52	96	46	22
BELLE	Bokombako	84	161	62	24
CLARA	(no native name)	32	50	25	9
DASEY	Louj	7.9	49	28	11
IRENE	Boken	6.7	34	24	4.5
JANET	Enjebi	24	40	15	6.2
KATE	Mijikadrek	7.3	16	15	8.8
LUCY	Kidrinen	13	23	20	13
MARY	Bokenelab	9.2	21	12	6.4
NANCY	Elle	8.7	23	19	12
OLIVE	Aej	14	14	11	6.4
PEARL	Lujor	9.3	15	26	6.8
RUBY	Eleleron	1.5	5.9	6.1	1.2
SALLY	Aomon	3.9	5.5	7.6	2.9
TILDA	Bikile	4.5	7.8	4.1	2.4
URSULA	Lojwa	1.5	2.9	1.2	0.7
VERA	Alembel	5.7	0.83	4.3	2.6
WILMA	Billae	1.7	0.72	2.0	1.2

picocuries per gram

These numbers are 0-5 centimeters soil increment

(dictated by W. L. Robison, Lawrence Livermore, August 6, 1980)

ENEWETAK ISLANDS		CESIUM	STRONTIUM	PLUTONIUM	AMERICIUM
ALICE	Bokoluo	52	96	46	22
BELLE	Bokombako	84	161	62	24
CLARA	(no native name)	32	50	25	9
DASEY	Louj	7.9	49	28	11
IRENE	Boken	6.7	34	24	4.5
JANET	Enjebi	24	40	15	6.2
KATE	Mijikadrek	7.3	16	15	8.8
LUCY	Kidrinen	13	23	20	13
MARY	Bokenelab	9.2	21	12	6.4
NANCY	Elle	8.7	23	19	12
OLIVE	Aej	14	14	11	6.4
PEARL	Lujor	9.3	15	26	6.8
RUBY	Eleleron	1.5	5.9	6.1	1.2
SALLY	Aomon	3.9	5.5	7.6	2.9
TILDA	Bikile	4.5	7.8	4.1	2.4
URSULA	Lojwa	1.5	2.9	1.2	0.7
VERA	Alembel	5.7	0.83	4.3	2.6
WILMA	Billae	1.7	0.72	2.0	1.2

	1 yr Encew Belen 350 d	350 d	1 yr Encew Belen 350 d	350 d	1 yr Encew Belen 350 d	350 d	1 yr Encew Belen 350 d	350 d	1 yr Encew Belen 350 d	350 d
	1 yr off		2 yrs off		2 yrs off		3 yrs off		3 yrs off	
	Impr. feed	X	Impr. feed	X	Impr. feed	Y	Impr. feed	X	Impr. feed	X
May an										
WB	106	186	88	168	107	186	88	168	107	186
Bm	110	198	92	179	110	197	92	179	110	197
3X May										
WB	318	558	264	504	321	558	264	504	321	558
Bm	330	594	276	537	330	594	276	537	330	594
30yr										
WB	1600	3000	960	1900	1100	2000	760	1500	860	1600
Bm	1700	3300	1030	2100	1200	2200	810	1700	920	1800

	100% IMPORTED FOOD	ENEU X	BIKINI IMPORTED FOOD	X	ENEU 330 IMPORTED FOOD	BIKINI 35 X	ENEU IMPORTED FOOD	1 YEAR ON 1 YEAR OFF X
--	--------------------------	-----------	----------------------------	---	------------------------------	----------------	--------------------------	------------------------------

Max. Annual

WB	124	242	1065	1974	88		168
BM	130	261	1094	2063	92		181

3X Max.
Annual

WB	372	726	3195	5922	264		504
BM	390	783	3282	6189	276		543

30 Year

WB	2800	5400	24,000	44,000	1400		2800
BM	3000	6000	25,000	47,000	1500		3100

	1 YR IMPORTED FOOD	ENEU 330 DAYS BIKINI 35 DAYS X	1 YEAR ENEU 2 YEARS OFF IMPORTED FOOD	X	1 YR ENEU 330 DAYS BIKINI 35 DAYS 2 YEARS OFF IMPORTED FOOD	X	1 YEAR ENEU 3 YEARS OFF IMPORTED FOOD	X	1 YR ENEU 330 DAYS BIKINI 35 DAYS 3 YEARS OFF IMPORTED FOOD	X
--	--------------------------	--------------------------------------	--	---	---	---	--	---	---	---

Max. Annual

WB	106	186	88	168	107	186	88	168	107	186
BM	110	198	92	179	110	197	92	179	110	197

3X Max.

WB	318	558	264	504	321	558	264	504	321	558
BM	330	594	276	537	330	594	276	537	330	594

30 Year

WB	1600	3000	960	1900	1100	2000	760	1500	860	1600
BM	1700	3300	1030	2100	1200	2200	810	1700	920	1800

REPOSITORY PNNL
COLLECTION Marshall Islands
BOX No. 5684
FOLDER Untitled

DOCUMENT DOES NOT CONTAIN ECI

Reviewed by J. Schwartz Date 4/30/97