RADIOTOXICITY RESULTING FROM EXPOSURE TO FALLOUT SIMULANT

II. THE METABOLISM OF AN INHALED AND INGESTED SIMULANT OF FALLOUT PRODUCED BY A LAND-BASED NUCLEAR DETONATION

Research and Development Technical Report USNRDL-TR-118 NM 006-015.04

U.S. Army

11 January 1957

by

S.H. Cohn W.B. Lane J.K. Gong R.K. Fuller W.L. Milne

Health and Biology

Technical Objective AW-6

Biological and Medical Sciences Division Captain A.R. Behnke, Jr., (MC) USN, Acting Head

> Chemical Technology Division E.R. Tompkins, Head

Scientific Director P.C. Tompkins Commanding Officer and Director Captain Richard S. Mandelkorn, USN

U.S. NAVAL RADIOLOGICAL DEFENSE LABORATORY San Francisco 24, California

ABSTRACT

The present study was designed to reproduce in the laboratory an acute exposure of mice to early fallout (2 days old) such as might result from a land-based nuclear detonation. Biological data were obtained on the uptake, distribution, and retention of the inhaled fallout simulant. Further, as a basis for comparing the effects of different types of simulants, the metabolic behavior of this dry-particle fallout simulant of limited solubility was compared to that of two previously studied fallout simulants; an ionic liquid aerosol and a mud-slurry aerosol. The simulant was also administered by gavage to provide data on uptake of fallout by ingestion as compared with inhalation. From these data, an evaluation was made of the radiation dose to individual tissues from inhaled fallout as compared to the concomitant external radiation dose that the animals would receive if exposed to the same airborne simulant.

ii

<u>UNCLASSIFIED</u>

SUMMARY

The Problem

The present study was designed to reproduce in the laboratory an acute exposure of mice to early fallout (2 days old) such as might result from a land-based nuclear detonation. Biological data are required on the uptake, distribution and retention of the inhaled fallout simulant. Further, as a basis for comparing the effects of different types of simulants, the metabolic behavior of this dry-particle fallout simulant of limited solubility must be compared with that of two previously studied fallout simulants. The simulant administered by gavage provides data on the uptake of fallout by ingestion as compared with inhalation.

These data are required for an evaluation of the radiation dose to individual tissues and organs from inhaled fallout as compared with the concomitant external radiation dose that the animals would receive if exposed to the same airborne simulant.

Findings

Following a 3-hr exposure to a dry-particle simulant of a land-based nuclear detonation, the activity was quickly cleared from the lungs and appeared primarily in the gastrointestinal (Gl) tract. Lesser concentrations of activity were also found in the head, liver, skeleton and thyroid. The radioactivity was removed very rapidly from the GI tract, as compared to the rate of loss of the simulant from the respiratory system.

The ratio of activity in the GI tract to that in the respiratory system following exposure to the ionic, mud-slurry and dry-particle simulants was 3.2, 12, and 80, respectively. The rate of biological elimination of the dry-particle simulant material from the skeleton and liver was considerably greater than was noted with the previously studied ionic simulant. In terms of total respiratory and GI tract activity at 1 hr, the skeletal activity was twice as high for the ionic simulant as for the dry-particle simulant, while the reverse was true for the liver activity.

Following the administration of the simulant by gavage, the initial distribution of activity in the GI tract and in the tissues was very similar to that following inhalation exposure. It was found that the absorption across the GI tract provided an important portal of entry for the dry-particle simulant into the systemic circulation following an inhalation exposure. The composition of the simulant material in the various tissues during the first 4 days following exposure appears to be dominated by one or a group of fission products, as seen from the similarity of the radioactive decay rates for most of the tissues. The results were similar for both inhalation exposure and administration by gavage. Exceptions to these findings were the thyroid and the skeletal tissues, the former having an affinity for the short-lived iodine isotopes and the latter for the longer-lived fission products.

The thyroid received the highest dose to any tissue from the internally deposited fission products. The GI tract received the next highest dose which was, however, less than 10 per cent of the dose to the thyroid. The dose to the skeleton, while lowest in the 15-day period studied, will probably be greater than that to other tissues over a longer period of time, since the skeletal activity falls off more slowly than that in other tissues. The internal radiation dose to individual tissues was, with the exception of the dose to the thyroid, lower than the concomitant external dose received by the animals.

iv

<u>UNCLASSIFIED</u>

ADMINISTRATIVE INFORMATION

This work was done as a part of Bureau of Medicine and Surgery Project Number NM 006-015.04, Phase 1, Technical Objective AW-6, as described in the U.S. Naval Radiological Defense Laboratory Research Progress Report to the Bureau of Medicine and Surgery, NAVMED 1343, of 31 December 1955.

The work also is part of the technical program for the Department of the Army established between Department of the Army, Office, Chief of Research and Development, and Bureau of Ships (Joint Agreement, 23 November 1955).

Acknowledgment

The authors wish to thank R.R. Odom, HN, USN, and Mr. L.L. Wiltshire for their technical assistance.

seif fat fe freig - excass		1 ilin .	6 1	entrog en Ar de	in mice au .
ter to reall	i	lynas i e	111	lanst	il blo enbys
tripping of the	١	11 Hole of	11, 1	ig-miles	CHURCH PERSON
THERE		10.11		11111 281	on en liter;
THAT POOP OF LIGHT		(sid)	1 () 1 ()	little 6 i	and a state of the area
ad'i s		LARIE	ι.	up 4	and a strate added
		* \$ 11 :	۰.	14411	e to Aprilia
1), 1 (1), 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,		- 4 3 2		at the	side spoklat

Exposure to fallout from a nuclear detonation, or to an aerosol formed in controlled nuclear fission, results in a biological hazard both from the external radiation of the organism and from radiation emanating from internally deposited material. Previous studies have indicated that the short-term effects which appear following a combined exposure (i.e., external and internal exposure) result primarily from the external radiation. The amount of material sufficiently great to penetrate the natural filtering defenses of the living organism and produce immediate effects is necessarily associated with a very large amount of external radioactive material. There are, however, situations in which the internal radiation is of primary concern. Such situations include the long-term effects produced by internally deposited isotopes with long radioactive and biological half-lives; long-term effects of accumulated small doses of short-lived as well as long-lived isotopes; and damage to individual tissues resulting from selective localization of isotopes.

In order to assess these situations, it is necessary to understand the metabolism of the fission products, i.e., the uptake, distribution, and retention in the body of the various isotopes which are inhaled and ingested. While a number of the long-lived fission products have been studied,¹⁻³ information is lacking in two areas: first, on the metabolism of a very early fission-product mixture (which includes many short-lived fission products) and secondly, on the behavior of longer-lived fission products competing in the body metabolism with the numerous other fission products found in the spectrum formed on bombardment of uranium. The behavior of an isotope administered singly may differ from the isotope taken in a mixture of fission products.

While physical instrumentation can be used to determine the external dose to an organism, it is not sufficient for calculating the internal exposure since complementary metabolic data are lacking on the quantity of internally deposited material in particular tissues and organs and on their turnover rates as a function of the amount present in the external environment.

In the present experiment, therefore, a direct evaluation was made of the uptake and distribution of fission products deposited internally

-1.

in mice as a result of exposure to a dry-particle simulant incorporating 2-day old fission material. The simulant employed was composed of tracer amounts of fission-product chlorides adsorbed completely on dirt particles in a wide size range. The simulant was generated as airborne particulates and represents, qualitatively, the fallout which might be encountered as a consequence of a land-based nuclear detonation. The simulant is, however, not necessarily of the same chemical form as fallout which would tend to maximize the hazard as approximated here.

Earlier studies have traced the fate of two other types of simulants an ionic simulant and a mud slurry simulant. The former represents the fallout that might occur following a nuclear detonation in sea water, and the latter a detonation in a shallow harbor. The biological response to an exposure to fallout is dependent on the properties of the carrier material. The results obtained with the dry-particle simulant in the present experiment were compared with results obtained with other types of simulants produced in previous studies.

In order to assess the relative importance of inhalation and ingestion as modes of entry for the firstion products into the body, the dry-particle simulant was administered to mice by gavage and the response to this mode of administration was compared with that following inhalation

exposure,		eigent as they	1541	4 }
	10.4	gent grave	日本語で	
· · · · · · · · · · · · · · · · · · ·	ner	eger, ett	ou the second	;
	i i i i i i i i i i i i i i i i i i i	减效 (1999年1月11日)	in gan	· 1
EXPERIMENTAL	ive p	A State of the	1+ 111	: *)
	144	the transmission of transmission of the transmission of transmiss	(d.)	
	20.9	ag į	2.1	ì
Preparation of the S	imulant	4.	F	
	* * * * * * * * * * * * * * * * * * *	,		

The simulant was prepared from neutron-irradiated uranium obtained from the Materials Testing Reactor. The mixture of fission-products was dissolved in HCl and added to a solution of previously prepared dirt particles of Ambrose clay loam of less than 40 microns in diameter. The particle-fission-product mixture was evaporated to dryness in a ball mill to insure mixing. The resultant simulant consisted of dirt particles with an average diameter of from 1 to 5 microns with the fission products adsorbed on the surfaces. The solubility of the activity adsorbed on the clay particles was 8 per cent in water and 18 per cent in HCl at pH = 1. The animals were exposed to this simulant within 2 days after the removal of the irradiated uranium from the reactor.

* Materials Testing Reactor Irradiation Service, Phillips Petroleum Co., Idaho Palis, Idaho.

-2-

Aerosol Generation

The experimental appartus for generating the aerosol and exposing the animals (see Fig. 1) was previously described.^{4,4} The equipment consisted of: (1) an aerosal generator, (2) the animal exposure chamber, (3) sample collector for collecting particles (for determination of size and aerosol concentration) and (4) supporting equipment (pump, etc.).

A generator was designed to provide a stream of airborne dry particles at a constant concentration level $(3.2 \times 10^{-2} \,\mu\text{c/ml})$.⁶ The rotation of the outer tube provided is constant advance of a brush against the surface formed by the dry particles. The particles thus agitated were picked up by the airstream and delivered to the exposure chamber at a flow rate of 20 cu ft/hr.

In the present study, **A** mice were exposed to the dry-particle-fallout simulant in a manner idea cal to that used with the ionic simulant.⁴ In addition, a solution containing a measured amount of the dry simulant was administered by stonich tube to 36 mice.

The animals were sacrificed in groups of 6 each, starting at 1 hr after exposure and at various intervals up to 15 days. The experimental procedure for measuring the activity of the fission products in the tissues has been described in a previous report.⁴

The distribution of gamma activity was determined in the following organs and tissues: rear story tract (starting at the upper end of the larynx) and lungs, liver, bia, thyroid, head, and GI tract and contents (from top of esophagus to mal region). The autopsies were carried out with care to avoid cross untamination.

The tissue assays were performed by determining the gamma activity with a sodium iodide gamma scintillation counter. The tissues were mounted in Coors porcellin crucibles and positioned under the crystal. Beta activity was not measured due to the inherent difficulties with preparation of samples for counting. However, previous experiments have shown that the gamma activity measurements satisfactorily reflect the total fission-product activity.

RESULTS AND DISCUSSION

Initial Uptake of the Inhaled Aerosol by Animal Tissues.

The distribution of the radioactive fallout simulant in the tissues and organs of mice exposed to the airborne simulant for 3 hr, and serially

. 3.

434



Pig. 1 Mouse inhalation Chamber. Method of exposing mice to radioactive aerosol is illustrated.

۰,

!

· · ·

UNCLASSIFIED

网络

sacrificed at periods ranging from 1 hr to 15 days following exposure, is shown in Fig. 2. At 1 hr after exposure, approximately 90 per cent of the internal activity (as determined by gamma activity measurements) was found in the GI tract, although all the activity originally entered the body by inhalation. It appears that most of the radioactive maferial was associated with large particles which presumably were rapidly moved back up the respiratory tract by ciliary action and swallowed, while the smaller particles penetrated to the alveolar tissue. A small but significant fraction of the material in the GI tract was absorbed into the body following either inhalation or gavage; the remaining material was excreted, By the third day only 1 pericent of the maximum activity in the GI tract remained.

The major portion of the radioactive substance which entered the systemic circulation did or in a matter of hours. Loss of activity from the blood was very rapid. At 1 hr the activity in the liver was approximately 2 per cent of the activity in the GI tract, while both the respiratory tract (including lung) and skeleton contained approximately 1.2 per cent of the activity. The thyroid contained 0.2 per cent of the GI tract activity and the head 9 per cent. The high level of activity in the head probably reflects the presence of large particles which were inhaled and trapped in the nasal passages.

The initial distribution of activity differs markedly from that observed following exposure of mice to the ionic liquid acrosol⁴ and the slurry type acrosol⁵ (see Table 1). On the basis of the activity found in the respiratory tract being 1, the activity of the GI tract was 80 ± 13 for the dry particle simulant (Type III), as compared to 3.2 ± 0.3 for the liquid acrosol (Type I), and 12 ± 1.1 for the ionic-mud slurry acrosol (Type II),

If the total activity found in both the respiratory tract and the G1 tract is used as the basis for comparison, the amount of activity retained at 1 hr in the skeleton is 2.3 ± 0.1 per cent with the Type I simulant and only 1.3 ± 0.1 per cent with Type III, suggesting that the dry particles may not be as available to bone as the ionic form. On the other hand, the activity retained by the liver when the animal is exposed to Type III simulant (again in terms of total GI and respiratory tract activity) was 1.9 ± 0.1 per cent as compared to 1.1 ± 0.1 per cent for Type I. The higher retention of the particulate simulant by the liver is probably associated with the natural function of particle entrapment by the reticuloendothelial system. These findings suggest that the particulate aerosol retains its adsorbed fission products in the body.

Initial Uptake of Ingested Simulant

In the experiment in which the dry-particle simulant was fed the mice by stomach tube, the initial distribution of activity in the GI tract and in the soft tissues and skaleton is very similar to that following inhalation

- 5 -



A State Sec.

TABLE 1

Distribution of Activity in Mice | Hr After Exposure to Fallout Simulants

11

s a construction of the second s		<u>'1'</u> 23	La Carte de la companya de la company	* * • • • •
is in the line of			e of Fallou Simulant	•
Tissue.	- - -	Inic I	Ionic Mad Shurry	III - Harting
Respiratory: Tract and Lun	g(4)			1. 5.55 1. 5.55
Gastrointestinal Tract and	Contents	8.2.1 + .3 1.	12 () () () () () () () () () () () () ()	80 <u>+</u> 18
Skeleton e († 1915) († 1934)	:	0.096 + .011	0.22 # [604v los () -	0,99 +
ter Head (ar each an teach and teach an teach	· ·.	0.28 + .064	1.1 1.26.100 (0.1)	6.7 ± ,76
Energ Liver 2021 - 1993 Anno 2014 (1994) - 19		0.043 ± .0078	0.31 (11) (16) (15) (16) (17) (16) (16) (17) (16) (17) (16) (16) (16) (16) (16) (16) (16) (16	1.5 ± .71
Thyroid (0.0083 ± .003	0.028	0.15 ± .07
(a) Respiratory tract acti respiratory tract activ	vity is exp vity.	ressel as 1. All ot	her values are expressed in ter-	ins of ratio to
1		∔ ≯. ∦	11 1 1 C	ŧ

TABLE 2

đ.

Distribution of Activity in Mice at 1 Hr Following Administration of the Dry-Particle Simulant

Tissue	Method of Adr	ninistration of Simulant
Gastrointestinal Tract and Contents(a)	31	
Respiratory Tract and Lung	0.0125 ± .002	0,0064 ± .001
Skelcton	0,01 24 <u>j</u> ± .001	0.0028 ± .001
Liver	0.0194 (± .004	0.010 ± .003
(a) Gastrointestinal tract activity is expl ratio to gastrointestinal tract activity	ressed as 1. All other	values are expressed in terms of

UNCLASSIFIED

DAT

1-14

1

exposure (Fig. 2B). The comparison of the amount of activity in the various tissues following analation and gavage of the Type III simulant is presented in Table 2. Following gavage, approximately 1.3 ± 0.2 per cent of the activity in the GI tractiat I hr gained entry into the blood stream and was retained by the liver and skeleton, the primary sites of internal deposition of the simulant. In comparison, about 3.2 + 0.3 per cent of the GI tract activity was found in the liver and skeleton following the inhalation exposure to the same simulant." Thus, absorption from the GI tract accounts for an average of 23 per cent of the activity deposited in the skeleton following an inhalation exposure, and 50 per cent of both the liver and respiratory tract activity. Only half the activity in the respiratory system, therefore, derives from material directly inhaled and absorbed through the alveolar dissue; the other half is obtained from material which enters the circulation upon absorption through the GI tract. These data emphasize the fact that the GI tract is a significant portal of entry of a dry-particle aerosol into the systemic circulation following an inhalation exposure of Thus, the physical characteristics (such as particle size) which have to be considered in evaluating the absorption of particles must be considered with respect to the intestinal membranes as well as the alveolar tissue.

111

The radioactivity in all the tissues except the skeleton, and thyroid following both inhalation and gavage decayed rapidly, decreasing to a fraction of 1 per cent of their 1 hr activity by the 15th day. The skeleton and thyroid, however, retained from 4 to 9 per cent of their 1 hr activity at the 15th day. The drop in activity in each tissue was a function of two simultaneous processes: the radioactive decay of the fission products and the biological loss of the simulant material itself from the tissues,

Radioactive Decay of Tissue Activity

The composition of the radioactive material deposited in a tissue is reflected in the radioactive decay ourve of that tissue. In order to gain information on the selective uptake, the gamma activity of the tissues of animals sacrificed at 1 hr post-exposure was measured at intervals over a 30-day period. These decay curves for various tissues (following administration by both inhalation and gavage) are shown in Fig. 3. The similarity of the curves (with the exception of the skeleton) indicates that the isotopic distribution of the internally deposited fission products in most of the tissues was very similar. The lower rate of decay of the activity in the skeletal tissue reflects a fractionation in the uptake of the fission-product mixture due to the affinity of bone for certain of the elements which are characterized by relatively long half-lives. The activity of material deposited in the liver decayed with a somewhat higher rate than other soft tissues. The apparent increase in activity at 1 day proba ably reflects the growth of a gamma-emitting daughter from a beta-emit ting parent radionuclide.

-8-



The composition of the fission-product mixture in the various tissues (with the exception of the thyroid) appears to be dominated by one or a group of fission products, as seen from the similarity in the radioactive decay rates for all the tissues during the first 4 days. The thyroid, of course, has an affinity for iodine¹³¹ and the shorter-lived iodine radioisotopes.

A gamma spectrum analysis will be used in future experiments to identify the major fission-product components of the simulant material,

Following the administration of the simulant material by stomach tube, the radioactive decay of the material deposited in the various tissues (Fig. 3B) was very similar to that observed following the inhalation exposure. The basis for the similarity probably lies in the fact that in either mode of administration a large fraction of the radioactive material enters into the systemic circulation by the GI portal of entry.

Biological Loss of Simulant

The gamma activity of each tissue taken from mice sacrificed at various time intervals following exposure was corrected for radioactive decay back to 1 hr after exposure by the use of the above-mentioned radioactive decay curves. The resultant curve approximately describes the biological loss of the fallout simulant in that tissue (Fig. 4).

The activity of the GI tract and its contents decreased the most rapidly, due to the excretory nature of this organ. As previously noted,⁴ the biological decay of the radioactive material in the GI tract was very rapid and could be separated into two components. The initial rapid loss of material with a few hour's half-life probably corresponds to the rapid loss of material from the GI tract via excretion. The second component with a half-life of several days was probably related to the slower pulmonary elimination of the lung "fixed" activity and to the normal excretion of internally deposited material. The rate constant for the loss of activity from the blood is also quite high initially, due to the rapid exchange of material from blood to other tissues. After this initial rapid loss, the activity in the blood approaches a constant level.

Following administration of the simulant by gavage, the material rapidly passed through the GI tract at approximately the same rate that was observed following inhalation. The rate of loss of the simulant through the GI tract was such that the stomach and small intestine contained 1 per cent of the maximum activity at 2 and 5 days, respectively, post-radiation. The activity in the large intestine fell to 1 per cent of the original activity by the 18th day.



j

ł

The very rapid clearance of radioactivity from the GI tract contrasts with the relatively slow biological loss observed in the respiratory system. At 5 days following an inhalation exposure, 42 per cent of the maximum activity of the respiratory tract is still present. The loss of activity from the lung for the first 5 days was approximately the same following the inhalation of simulants I and III. The curve of biological decay for the respiratory system can be described by 2 rate constants. The initial rapid loss of material (half-life about 10 hr) is probably associated with the 'second process (with a half-life of approximately 7 days) is probably associated with the slower loss of the smaller particles "fixed" in the alveolar tissue. With simulant III the biological decay of respiratory activity following administration of the simulant by gavage is much more rapid than that following inhalation (Fig. 4B) reflecting the different mode of fixed of the material in the lung. Following inhalation, a large fraction of the activity is on the air side of the alveolar membrane, while following gavage the material is deposited on the blood side of this tigue.

Following gavage, the pativity in the liver built up to a maximum at around 3 days, in contrast to the more rapid build-up following the inhalation exposure. After 3 days the biological turnover of activity by this organ is the same for both modes of administration. For both types of administration the activity in the thyroid builds up to a maximum in 2-1/2 days, and then drops off.

With an inhalation exposure, the concentration of activity in the skeleton tapered off to a relatively constant value after an initial rapid loss. It stayed fairly level in the period 5 to 15 days. With gavage, nowever the concentration appeared to increase in the 10-day period studied.

Evaluation of the Internal Radiation Hazard

While calculation of radiation dose from fallout with any degree of precision is difficult, an approximation based on experimental data is feasible and was performed here. In order to evaluate the dose to individual tissues following an inhalation exposure, the activity per gram tissue as a function of time was determined (Fig. 5). The greatest activity per gram tissue was observed in the thyroid at 1 hr following exposure. At this time the activity in the GI tract (including contents) was next highest, followed by that in the respiratory tract,

The total dose received by each organ for comparable energies is, of course, proportional to the area under its curve (see Table 3). The thyroid, for example, received 170 rad. This was by far the highest dose received by any of the tissues measured during the 15 day experimental period studied. The ratio of the dose received by the thyroid to

-12-

Calculation was performed by graphic integration.

2

· ••••• '#••• •							
<i></i>				/			41
	10	-			2 - 26 - 26 - 26 - 26 - 26 - 26 - 26 -		
.i !	Preduce	- 103 c -	De Date (C.S	(1 yl anti	an surva a	11.14	#6Q
	- Antoine - California	•		an en ander an a		an in the second se	A State and a state of the stat
		151					
	e. E	ľ	The second	1 14 (14 (14 (14)			
	(1/81) 107	1				The second se	
	011	F		au		LINKET	
	01	7		80.03		ast 15	
10 1 K	1.8 建				Burd hug	v fastickult	
	4.0 ¥				1	1. 4. 1. 1	N.
	a.1				3 14	610010	
1	8 10*	<u> </u>	9				
· · ·	Š	lt .				See Stanly 199	
x	5						
ug X B	riwistleit		an much	15 19	THYPRIPEQUA	Yathaa	Sum Ball.
sin	nt Co	-	Stewe Li		studia of	1-	1 01 4404
-		AR.	a second		с÷,		in Alganta
nt with	3 10*0	891	, Dice to sole in	· 11 3 6 1 1	DE CONTE LE	he to to	भाषतिकोः भ
- Hed	ad party		10 011 4111		rais no	ans goin	14-11-12-13
6.51811	id grant		CONTRACTOR OF	in the second	() Saund	19 39 X	TUN BOUL
VIII (Seveli	13月) 19月1日 - 19月1日 -						and the first
11111		[N]				日本 - 日本 1、日本日 - 日本 - 日本 - 日本 - 日本 - 日本 - 日本 - 日本 - 日本	
1()	VILL	1			. i. y. luci	1 65 11 11	il all
	101		The state			nel ant.	WHOT ME
	4						
		0 BLOOD	N Poliven		SKELETON Respiratory T	RACT	dji i
		G	ASTRONISTINA	TRACT	HEAD		injust v
	10 3		1		<u>_</u>		
	int.	0 1	00 🐙 200	300 TIME (MB)	4 <u>90</u> : (\$0)		t si
r.	ių.		nife has		3	. Port	
ts -	្រះ		- u 🙀 🖓 🖓	All a star	114 5	1. 9151	43
	B PI	g, 5 Gar	mma Activity P	er Unit Weig	ht of Tissue Fo	llowing	1 20 1 1 1 1 2 1
	1 đợi		Experieur	e to Simula n	CILLAR CONT	l l l sat	
	;						
			ាណ៍) សម័រមាន	apa -a	The entry	to the first state of the	vila all'i >
× .	. 1.		(Paperic) Jelan	Solida a t₩ solida avia	at wast the		THE LAURE
	₩.		AND STREET	9012 - E 1911 - E	+_111 + +€' →	er erste Detreter er	19777
• •			ŧ				···· • • • • • • • • • • • • • • • • •

-13-

UNCLASSIFIED

8 A . C O

TABLE 3

CT # 1 4 1

,		和 注意。	1		
i ja 19 ja 19 ja			is and	Tota	Dote
Ti	15 UQ X	(rad,	Ne Rate /hr at 1 hr)	Interval (hr)	(rad)
Thyroid			16	0 360	170
GI Ttact			0,96	0 - 168 /	16
Respiratory T	ract and	LIREN	0.94	0 - 860 /	8.0
Head			0.15	0 - 360	6.4
Liver			0.050	0 - 168/	8.2
Skeleton			0.029	0 - 360	1.5

Dose to Tissues From Internally Deposited Fission Products

that received by the respiratory tract was much greater following exposure to the dry-particle simulant than following inhalation of the ionic simulant.

It is to be noted that, over an extended period, bone would receive the greatest integrated dose since the isotopes deposited in bone have halflives ranging up to 27 years (Sr^{90}) and, in addition, have long biological half-lives. In contrast, the longest-lived iodine isotope has an effective half-life of 7.7 days and thus delivers its dose in a short period of time.

The dose rate to the individual tissues can be determined by use of the following formula:

 $I_{ij} = K \frac{Q}{W} E$

where

I = dose rate in rad/hr

= beta activity of each tissue in d/min (gamma activity was measured experimentally in this study and was converted to beta activity on the basis that the ratio of gamma photons to beta particles at this time was approximately 1.2).

The following document is referenced for the convenience of those who have access to it, Sondhaus, C. The Ratio of Lung Bets Does to Whole-body Dose During Given Time Intervals After an Atomic Bomb Detonation, U.S. Naval Radiological Defense Laboratory Report, USNRDL-394, 12 December, 1952 (CONSIDENTIAL).

-14-

月日人工的工作

ti ii

stollowing alculation applies only impunisance the local set of the rest of the second is Sil a kit a constant factor converting Mevito erginierg/gmito radi

follows:

0.2 Mev (the average beta energy for 1¹³¹, assuming the entire activity in the thyroidydeniver from this isotope

$$hr = (1.15 \times 10^{10}) (6.8 \times 10^{7})(2 \times 10^{-1})_{(18)}$$

This calculation does not consider the tellurium precureors or the intake of tellurium isotopes. However, as pointed out by Dunning,⁷ the error in ignoring these isotopes is not any greater than the other inherent uncer tainties in this thyroid dose approximation. The short-lived iodine iso-topes are not considered separately from 1¹³¹ since the mean of the ratio of energies of I short-lived/1¹³¹ equals 1 at this time.

The dose rate to the dither tissues at lahr wash so calculated (see Table 3). The value for $E\beta$ for all tissues other than thyroid is approxmately 0.39 Mev from I to 4 days post-detonation and 0.29 Mev from to 15 days.

· HOY

and all faites ab Bolan; and SB (The assumptions implicit in this calculation are direct, the simulant is homogeneously distributed in the tissue and secondly, the energy is a completely absorbed in the tissue. The energy is sorption per tissue, is taken, therefore, as equal to the energy emission, the the

Evaluation of the External Radiation Hazard

14

Since a simultaneous exposure of the animakitothe airborne simulanty occurs ordinarily with internal radiation, it is of interest to relate the internal dose with the concomitant external dose. In this particular experimental situation, however, the animals were not placed in the agrosol and thus the external dose could not be measured directly. The

See footnote on page 14.

UNCLASSIFIED

() TREAT IN COMMON 258 4

following calculation applicationly to the general case, it is to be noted that, in this calculation, show ogeneous distribution in the aerosol is assumed, and an infinite column of aerosol is also implied. It is to be noted further, that for calculation of the external dose the gamma radiation is off primary implemented () (while the internal dose is based on the beta radiation. An approximation of the external dose rate at 1 hr that would have been received by these mice had ther been exposed to it his aerosol in a 2. T field casine calculated as follow stated to the interna-

 $I_t = 51 \frac{Q}{W} E_{(1)} h_{(1)} 01 \times 8.0$

UNCLASS

 $I_{ij} = 0.2$ Mew [the average bota oneo, for $I^{(1)}$, resuming the cutive activity in the the roldwheet star scole configuration of the second star scole of the secon

 $\frac{Q}{W} = \text{activity, } \mu c \left(\frac{1}{2} + \frac{1$

This calculation does the wey if Wey are interested by the error of the error of

Threat 1060 rad/damor 44 rad/hr in a 4 Tifield of the sub all a sound of the state of the sub all a sound of the state of the sub all a sound of the state of the sub all and the sub all and

The integrated dose that the mice would receive externally when exposed for 3 hr to the second rosol in a 2 Tifield is approximately (60 rad. The ratio of the second rose to the dower received by the hing from the internal mainters was 8, which corresponds closely to theoretical calculations of the second second rose in the second rose in the second rose in the second rose in the second rose is the second rose in the second rose is the second rose in the second rose is the second ro

I the estad and the mail of the

The supported to the solution

a due drive ville school affinan

SUMMARY AND CONCLUSIONS

The present study was designed to reproduce in the laboratory an stor apute exposure of mice to they fallout (2 days old) such as might result from a land-based nuclear to phation. Biological data were obtained on

See footnote on page 14.

-16-

UNCLASSIFIED

Redformmer on a series

the uptake distribution and retention of the inhaled fallour simulanf. Further, as a basis to comparing the effects of different types of simulants, the metabolic behavior of this dry-particle fallour simulant of limited solubility was compared with that of two previously studied tallout simulants; an ionic liquid aerosol and a mud-slurry agrosol. The simulant was also administered by gavage to provide data on the uptake of fallout by ingestion as compared with inhalation.

Following a 3-br posure to the dry-panticle simulant, activity was quickly eladed (rom the lunguand appeared primarily in the GI tract. Lesser concentration of activity was also found in the head, liver, skeleton and thyroid. The radioactivity was removed very rapidly from the GI tract, as compared to the rate of loss of the simulant from the respiratory system.

The ratio of activity in the GI tract to that in the respiratory system following exposure to the ionic, mud-slurry and dry-particle simulants was 3.2, 12 and 80, respectively. The rate of biological elimination of the dry-particle simulant material from the skeleton and liver was considerably greater than was noted with the previously studied ionic simulant. In terms of total respiratory and GI tract activity at 1 hr, the skeletal activity was twice as high for the ionic simulant as for the dryparticle simulant, while the reverse was true for the liver activity.

Following the administration of the simulant by gavage, the initial distribution of activity in the GI tract and in the tissues was very similar to that following inhalation exposure. It was found that the absorption across the GI tract provided an important portal of entry for the dryparticle simulant into the systemic circulation following an inhalation exposure. The composition of the simulant material in the various tissues during the first 4 days following exposure appears to be dominated by one or a group of fission products, as seen from the similarity of the radioactive decay rates for most of the tissues. The results were similar for both inhalation exposure and administration by gavage. Exceptions to these findings were the thyroid and skeletal tissues, the forme having an affinity for the short-lived iodine isotopes, and the latter for the longer-lived fission products.

The thyroid received the highest dose to any tissue from the internally deposited fission products. The GI tract received the next highest dose, which was, however, less than 10 per cent of the dose to the thyroid. The dose to the skeleton, while lowest in the 15-day period studied, will probably be greater than that to other tissues over a longer period of

-17-

I N CI A SSIT I I II

di Welta

rapidly from

built mort 1

molay where

or a lutil and

unfigation of

· Witto-

VILAY

"iped-

小小小田

- føder

ះស្ថិរភេ

had

Section.

11/8 11

Ignv:

1 Ann

1 this

'ağ 1

-tily-

¥111

abin

前小

te ciùi ret

(111) · 6W 19

umie sindi l hreit y de diroi.

and a state of the second

ちんち まいちょうちんしいた いちち

Rh

1+

f

17

.

11

3

1.

j.

n ilie e

マロロ戦

ilt nl

- 64V

illa s

11 191

-1

19 × Y7

dyol

11 A 1		2.8			
	ne clare the	a kolota lina	ëvity falls off	more alowly the	n that in other
		1757 997 7369 53	11 * 1717) - 115 16 1 7 F	9 3*97 * 1 50 * * * * * * * * *	
1 . A . A . A . A . A . A . A . A . A .	aue The	internal	lation dose to	individual tissu	es was with th
- 1 111	THITCHERGE	THE REPORT OF THE REPORT OF THE		1001001401	
∴ đe X	ception of th	ie dose for	h thyroid, lov	ver than the cond	comitant extern
	TOP JIER LITTLE		A TOPPING	រច <i>ាលប្រ</i> ោ កព័ត្ ថា ចារថា	181970 8847 83755
	SE TECEIVER	DV TOCKED D		3 H SI	STREET STREET

within nathraid.	- ALADIG	atter strategister	「「「「「「「「「」」」」「「」」」「「」」」	- / LUBIHOA DATE:	1)
Man The The	23 F H B	S & SHANGO	as blepth after	ออน สายเสมชาวิต เก	
information in the	H AND	gave of	ban remini	- Is esw freine :	?
		and differ	TRUEIN RULER	· sailout by inc.	۰,
			pproved by:		
BAW VIIVE	6 Hatt	Here yest	Jan Barris	" Eline Astorn How	
3. 10641 C. 91	11100	10 States 1	and the second	A State of the second se	11

 ϕ^{\dagger}

11

f • •

3

3

	+ •		1.	, CU				A Par	141	$(a) \cup (b)$
1.	j.			IVIA	1.5		101.1912	onre	วิรัยย	n + 1
1	. (()	Prof.	11.27		ب و فع	t KvY	bas	eton	Le dit.
i	1	6 B	Capt	ain (MC)	USN		Acta	2 10	
		A.	Actii	ng H	ead,	Biol	ogića	l an	A Sector	
			Medi	cal :	Scien	Ce s	Divt	í i ón 🖯		1: I
		<2 V		22		*			1.00	

ŧ ť

ų

												·					್	1		1		41.4	1.1			
	1	1	11	3	1 10.	in L		12	.91	11	1	i_j	U)	Ì.	11		1	lo	Ĵ	Ĵ.	(p	Ν.	e e e	11		
18				.	ASH'	r o	T.	<u>un</u>		50	10	Ð		4	ر د ارد	Ri I	1		C U I X	Q1	Y.	n		\mathbf{o}^{\dagger}		
1	, t				an		1	10.12	11		ci'i) 171		0	Č.			5	: r	, n	Ċ	E.	•••		
·				,		;	, ,	: 7.¥ - 64.≹			ereg u Alva	ية جا معا					1		يت : سنة	₹, † 	1	÷.	1 % 	; ,		, 1
1			·	1	erana e Good a co			11	1 '	1.4	147	41	11	+ 1 + +		1		9. F	13	. 13	4 *	Ŷ	. 1	,		ţ
	14	. •	. 1	1	្រភ្លូរ			÷ιQ	11		ήΨ	1	14		1.3	9	1	;	1]		χļ	4	នា		٠.	
1	7				an		1	157	H	• •	74	ľ	6]	1	1	Ċ.	4	f f	ry,	鮒	Ĵ	ų		• *	••	ì
	ţe,	٠.	'	11	- Ser 1		, '	d g	1	٠,	1	W	1	13	~* †	Ý	(Ŷ	ŗν	ţ\$	Ø į	۱.	ĻE	đa	1	ł	ł
	LI.	•			w.		-	11			1.5	Лį	H		, .	1a	4	41	(1	4	\$	9Ì	ÚÌ.	1		i
100					- 5												- E.			P 1						1

1 201	ngetseteiner	liss off galwollis
ers 🏚 👘	ay in the G	dist in all the start of a lar
	WY WIRL	a grievollog tale a
÷۴,۰	10 10 10 10 10 10 10 10 10 10 10 10 10 1	an use the GIT term
· 1.:	· inge alter	indumie upsis of
·/i	11 11 1 11 11 11 11 11 11 11 11 11 11 1	is suffre to an in the state
r v	1111180	di gnerubigene if
- 4	al company	in leader by one on
' *O	$(\theta_{1}, \theta_{2}) = (\theta_{1}, \theta_{2})$	antipochier and the
ň	A CONTRACT OF A CONTRACT. CONTRACT OF A CONTRACT. CONTRACT OF A CONTRACT OF A CONTRACT OF A CONTRACT OF A CONTRACT. CONTRACT OF A CONTRACT OF A CONTRACT OF A CONTRACT OF A CONTRACT. CONTRACT OF A CONTRACT OF A CONTRACT OF A CONTRACT. CONTRACT OF A CONTRACT. CONTRACT OF A CONTRACT OF A CONTRACT. CONTRACT OF A CONTRACT OF A CONTRACT. CONTRACTACT OF A CONTRACT. CONTRACTACT OF A	a dipd rol galie of
5 / 1	ALC SHILL	a sight of games a
1,1	$\{e_{i}, e_{i}, \dots, e_{i}\}$	indite as guile a
	1 (f)	- 9711 - 1, Blite)

.11 on damy rate re-1 1900 · Peall Lighters 11 1 April n Bal: , [†] word , wain it's 110 $r \to r + 1$ I add it uaab bahly be greet · 1 the solar . . .

. +18-

REFERENCES

- 1. Abrams, R., et al., Metabolism of Inhaled Fission Product Aerosola U.S. Atomic Energy Commission Report, MDDC-248, 24 May 1946.
- 2. Hamilton, J.G. The Metabolic Properties of Fission Products and Actinide Elements, Rev. Mod. Physics, 20:718 (1948).
- 3. Scott, K.G., et al., Deposition and Fate of Plutonium, Uranium and their Fission Products Inhaled as Aerosols. <u>Arch. Path.</u> 48:31 (1949)
- 4. Cohn, S. H., Lane, W. B., Gong, J. K., Sherwin, J. C., Fuller, R. K. Wiltshire, L. L., and Milne, W. L. Uptake, Distribution, and Retention of Fission Products in Tissues of Mice Exposed to a Simulant of Fallout from a Nuclear Detonation. I. Simulant of Fallout from a Detonation Under Seawater. U.S. Naval Radiological Defense Laboratory Technical Report USNRDL-TR-77, 5 Dec 1955. Also A. M. A. Arch Ind. Health 14:333 (1956).
 - . Cohn, S.H., Lane, W.B., Gong, J.K., and Milne, W.L. The Preparation and Biological Application of Airborne Simulants of Fallout From Nuclear Detonation: An Experimental Approach for the Estimation of Inhalation Hazard Following Exposure to Radioactive Fallout. Presented at American Cancer Society Air Pollution Symposium, Atlantic City, Sept 1946.
 - Sherwin, J.C., Fuller, R.K., Lane, W.B., and Wiltshire, L.L. An Apparatus for Exposing Mice to Radioactive Aerosols. U.S. Naval Radiological Defense Laboratory Technical Report USNRD1 -TR-78, 2 Feb 1956

Dunning, G.M. Two Ways to Estimate Thyroid Dose from Radioiodine in Fallout, Nucleonics, 14:38 (1956).

(not with pimota) houngo (syst) bas do trassi loganta Continued Prantantion (rotto not al Committee) (ATTARO) mapping (ORITE) ere, the might meet to and here to proceed Comment hasman initiatal an and taking any apiralerndel leathathean Stinter and the San brend and Ole derald , builden it 1 N endagne Denartment, Aberdeen Proving Graum -045 ((), Ghomical Corps Toming Command (Lillarove) 1 11 CO, Ohe micht Weld Requirements Agency COPIES CO, Chunigal Warfare Langratories 1 (miles a chief Bignal Older (SIGRD-8B) NAVY Wighter Medical Been arch Laboratory 161 (1-VEIChief, Bureau of Shipe (Code Sed) (mit A healt rain W 43 $(i\dot{i})$ (auravit Chief, Bureau or Acronautics (Code AE40) Myon h 1.1 Chief, Bureau of Supplier and Mocounts ((Code W)) () Chief, Bureau of Yards and Docks (D-440) (640) a 59 10-11 12 Office of Naval Research (Code 444) Strand In Spille <u>թ</u>.գ. Chief of Naval Research (Code 811) stago , mispand 13 ðð Chief of Naval Operations (Op-36) Work (Margin C. 14 67 15 Commander, New York Naval Shipyand (Material Lab.) 33 Director, Naval Research Laboratory (Code 2021) 16-18 $(\cdot a)$ CO, Office of Naval Research, New Sorki 19-20 0.521 Naval Medical Research Institute on Hammanian 15 CO, Naval Unit, Army Ohemical Center (19.00) 22 CO, Naval Unit, CmlC Training Command 23 11 MININE 1 24 CO, U.S. Naval Civil Engineering (Resuland Evaluate). 25 U.S. Naval'School (CEC4Officeus) (Messacon Callor) Commander, Naval Air Material Center, Phildelphia 26 27 CO, Naval Schools Command, Preasure Island (10) 28 CO, Naval Damage Control Training Center, Philadelphia 29 U.S. Naval Postgraduate School, Monterey 1978 91 30 CO, Fleet Training Center, Norfolk "CO, Fleet Training Center, San Diego 31-32 11222233311 Office of Patent Counsel, Mare Island 33 34 Commander Air Force, Atlantic Fleet (Code 16F) 35 Commandant, U.S. Marine Corps 1. 36 Commandant, Marine Corps Schools; Quantico, (Library) 37: 00 Commandant, Marine Corps Schools, Quantico (Development. (1) 'Center)' - I 1. 16.06.5 hit 10 1 3 119 18 ie i 11() 11 fors be IO. C. A DINS · 1급 00 1.11.11 ARMY ALL FRANK AND ាត់អា 1. hait magazie 50111 118 "" Chief of Engineers (ENGEB, Dhein) (191) A 38⁰ 140 68 Chief of Engineers (ENGNB) main or good 39 ().) 1.190 White -21-

の設定

40-41	Chief of Research and Development (Atomic Division)
42	Chief of Transportation (TC Technical Committeel
43	Chief of Ordnan OPDTR
44	CG, Chemical Research and Development Command
45	CO, Chemical Claups Materiel Command
46-47	Chemical and Regiological Laboratories ACmic Maryland
48	CO. Chemical Coops Medical Laboratories
40	President, Cherry Board
50 51	Ordenne Dans Aberdeen Provine Crowder
	Ordnance Department, Aberdeen Froving Ground T
52	CO, Chemical Command (Library)
§ 53 -	CO, Chemical Requirements Agency
54-55	CO. Chemical Wallare Laboratories
56	Office of Chief Biggel Officer (SIGRD-88)
67	CO. A rmy Medical Besearch Laboratory
E9 E0	Wolton Dood/A Windigals Contain
20+27	Walter Accuss and and A weather to a soft a
OV	CG, Continentation which on mand, which which the velocity of the second
61	Army Medical Arms of Graduate School (Dept of Hiophysics)
62-63	Brooks-Army- Menored Cepter
64	CG. Quartermanne Bes. and Devy Command
65	Office of Quartemeter General Rando D.Div
66	Director, Operatings Besearch Office (Librarian)
67	CO Dugway Brown Ground
40	Surrow Concerts Alerian Manager Manager Aleria
00	
07	Surgeon Menereration and and the state interest and the state
新7U	Director, Evant man all Laboratory, Nucleonica Hersion)
N 71	CG, Engineer Researd Dev. Lab, [Library]
72	CO, Transpontation Res. and Deva Command, Fort Eustis
73	Director, Officeren Special Weapons Development
74	Director, Supsizin Research Unit Fort Sam Houston
75	CO Ordnance Materials Research Office. Wateriown
76	CO Energiand A monol
/0	SUU, F FARKIOFORDERAL
77	TOKYO Armyallogpere, Studie lavel
	atter a state of the second state of the secon
	AIR FORCE IM Jun Autout Player Star
78	Commander, Waint Air Development Center (WERDO)
79	Commander, Working Air Development Center (WODTV)
\$ <u>9</u> 0	Commander Am Des and Des Command (DDMDA)
N 00	
01	Commander, Alt Mes, and Der, Command, MALART
82	Commandant, School of Aviation Medicine his states
83	USAF, SAM, Randoph Field (Brooks), tailing a first
§ 84	CG, Strategic Air Command, Offutt Air Force Bage (IGABD)
§ 85	CG, Strategic Air Command (Operations Analysis Office)
86-87	Commander, Kirsland Air Force Base
88	Office of Surgeon General
89	Director. Air University Library, Maxwell Air. Porce Basen
	Commander Technical Training Wing 3415th Thick
8. 7 ∪ +7↓ 8. 02	CG Combridge Research Center (CRHTM)
4 76 (5	Or' Cumpting Carter Contes (Ottes W)

UNCLASSIFIED

2 		*	
	OTHER DOD TIVITIESTADATE TO DEST	A Land de MA	
	ALVAL SABOARA TA SACARA	A London	· V .
93	Chief, Armed Torces Special/Weapond Pr		- 08
04	A FSW D SWITH Sandia Remarks heaven		[2:
75 07			¢
72-71	AFSWF, Hq. maleid Command Meandlands	ARTIT PSVL	1. 1. 1. 1. 1.
78/狭空	Assistant Sequetary Of Defenant Heridands	Dev.)I unt	13
99-100	Assistant Sectorary of Delene Civil Dele	ose Divisio) . <u>a</u> t
101	Armed Force Medical Clarate Anothin	White Istor	101.000
102-111	Armed Service Technicat Information Ag	ACYNTER	1
ang di san	VUAUMMEINTELOTIA	Wend Flinket	6121.00
	AEC ACTIVIT ESTAND OTHERS INTAL	AH . LINCT	
			11/12-11
112	Alco Product Inc.	an our and	} ∖i
113 4116	Argonne Canding Research Hospital		
116 121	A reasonad Window and a bear on the parameter (1)	A DEDECT	402464
1772172	Atomic Book and a destroy of a former and and	THO DOIDED	· : : : : : : : : : : : : : : : : : : :
174 174	Atomic Thomas and a date Marchineton	p antapp	- 70 5 -96
1274120	Atomic Energy Sommission Control of States	CA DELETT.	11\$5-80
120 120	Restalle Merel Manual Transferrer	M ALDU	
1674130		Via svinU	$= \frac{1}{2} (c_{11})$
1,31,41,36*	Dettis UTARE AND DEIDEN AND OLARO IO	dianoviau.	012 (1)
133	Boeing Arrph Bucompany Heintoinad to v	University	615.810
134-137	Brookhaven - Brookhall Laboratory - (1) have	Sen svin D	e i t
138 🚯	Brush Beryland Company Tale dood love	tinn ovin U	ΞúΕ,
139	Chicago Pate Group (1 moter dati) lo y	de martale.	(·
140	Columbia Un Saty (Dailla) Junila W In V	TRANSTON .	
141	Colubmia University (Hassialis) on to all	MIL LINW	
142	Committee of Atomic Crevalting Washin	ton	\$28.23
143	Committee of Recta of Atomic Radiation	a she was a straight the	
144-145	Consolidated Jultee Airgraft Corporation		i da mili
146	Convair-General Dynamics Corporation	TET GLADEN	
147_148	Division of Pow Materials Denven	and eller freiter	
140 5 5	Dow Chemine Sampanya Dittely and		1
150 ····································	Dow Chemier Company, Producting	ha i sa kreduži	
150	Dow Chemican Sumpany Rocky Flats	, , , , , , , , , , , , , , , , , , ,	
171~174 当 1863	duPont Company, Aiken		
1.22% 1.22%	duPont Company, wiimington		
150-157	General Eler Company (ANFP)		
158-103	General Electric Company, Richland		
164-165	Goodyear Atomic Corporation		
166	Hawaii Maria Laboratory	·	
167	Iowa State College		
168-170	Knolls Atomic Power Laboratory		
171	Lockheed Ail maft Corporation (Cleveland	1.5	
172	Lockheed Ainstatt Corporation, Marietta	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	
173-174	Los Alamos mientific Laboratory	5.6 8.8	
175	Mallinckrodt shemical Works	新い	
76	Massachuset Institute of Technology (Ha	rdv) 🦈	
77	Mound Labor tory	t,	
178	National Advisory Committee for Aeronau	tics	
		i i	

-23-

179	National Bureau of Standards ((Library) and a starter	
180	National Bureau of Standards (Taylor)	
181	National Leads Campany of Ohio and the set	
182	New Brunswich Laboratory and an and	
183	New York Openations Officere hlor	
184	Nuclear Development Conporation of America	
185 (0	Cat Ridges Institute of Mucles & Studies	
186-191	Cak Ridge National Maboratory	
192	Patent Branc scentis hington index	
103-198	Phillips Petroleum Company	St 1 SAUL
100-200	Rublic Health Service Washington and the	
201	Radioisotopes (Eporatory (Thoma)	
202	RAND Cornoration	
203-204	Sandia Corporation (1986) (1978)	
205	Union Carbide Muclear Company (C-31 Plant)	
206-207	Union Carbide Disclear Company (K-25 Plant)	
208-211	United Aircraft Comparation	
212	UCLA Medical Research Laboratory	
213	University of California Medical Center	
214-216	University of California Radiation Laboratory, Berke	elev
217-218	University of California Radiation Laboratory, Liver	more
219	University of Chicago Radiation Laboratory.	
220	University of Rochester (Technical Reportiunit)	
221	University of Utah (Stoner)	
222	University of Washington (Applied Fisheries, Lab.)	
223	Weil, Dr. George L. In Million Hill	
224-227	Western Reserve University	が (1) (1) (1) (1) (1) (1) (1) (1) (1) (1)
228-252	Technical Information Extension, Oak Ridge	
	Turpar Del sele the anthritesanth.	
	USNRDL MAT A HIT HAT AND	
	the second s	as i s

ŧ

253-290 USNRDL, Technical Information Division t a t

.

DATE ISSUED: 21 January 957

		/4	THE HEAD STRATES	
,		۰ i ·	We with the state	
	۰.	- 1		
		11 1	and the second sec	
	1	. 1	B. I. D. KIN	
•	4 1	1411	the design of the	
	, i	1 6,11	A DE AND AND A STATE AND A STATE	
		141	added to make the first of	
	1.	10.10	A DAMAN MARKET	
	t	11 - C	and show in the state of the	
		1111.	Lat man Market	
		11	A DE ANTINA A DE A	

-24-

11

141 161

401 -

· · •2

1. Fallout - Metabolism Naval Radiological Defense Laboratory. . Fallout - Metabolism Naval Radiological Defense Laboratory. 2. Intestine - Effects of 2. Intestine - Effects of USNRDL-TR-118 USNEDL-TE-118 radiation radiation RADIOTOXICITY RESULTING FROM EXPOSURE TO RADIOTOXICITY RESULTING FROM EXPOSURE TO Lo Coin, S.H. FALLOUT SIMULANT. II. THE METABOLEM OF AN Cohn. S.H. FALLOUT SIMULANT. II. THE META BOLISMA OF AN II. Lane. W.J. INHALED AND INGESTED SIMULANT OF FALLOUT INHALED AND INGESTED SIMULANT OF FALLOUT Lane, W.B. 11 III. Title PRODUCED BY A LAND-BASED NUCLEAR DETONA-III. Title PRODUCED BY A LAND-BASED NUCLEAR DETONA-IV. Title: Metabolism. IV. Title: Metabolism ... TION, by S.H. Cohn, W.B. Lane, and others. TION, by S.H. Cohn, W.B. Lane, and others, V. NM 006-015.04 V. NM 006-015.04 11 Jan. 1957, 24 p. tables, diagrs, UNCLASSIFIED 11 Jan, 1957, 24 p. tables, diagrs. UNCLASSIFIED The present study was designed to reproduce in the The present study was designed to reproduce in the laboratory an acute exposure of mice to early fallout laboratory an acute exposure of mice to early fallout (2 days old) such as might result from a (2 days old) such as might result from a land-besed nuclear detonation. Biological land-based nuclear detonation. Hological dist were obtained on the uptake, distribudata were obtained on the uptake, distribu-UNCLASSIFIED tion. and retention of the inhaled fallout **UNCLASSIFIED** tion, and retention of the inhaled fallout (over) (over) 1. Fallout - Metabolism Naval Radiological Defense Laboratory Naval Radiological Defense Laboratory 1. Falloutin Metabolism 2. Intestine - Effects of URREDE-TE-118 Intestine - Effects of USNRDL-TR-118 TO: BRITANE PROMITORING TO: PADEDTOXIC TO RESIDETONS PLANE DEPOSITION THE METADOLISM OF AN Cohn. S. H. FALLOUT SIMULANT. II. THE METABOLISM OF AN Cohn. S.H. T. THEALED AND INGESTED SIMULANT OF FALLOUT Lane, W.B. III. INHALED AND INGESTED SIMULANT OF FALLOUT IL Lane, W.B. ECOLICED BY A LAND-BASED NUCLEAR DETONA-Title III. PRODUCED BY A LAND-BASED NUCLEAR DETONA-III. Title TICH, by S. H. Cohn, W. B. Lane, and others. IV. Title: Metabolism IV. Title: Metabolism TION, by S.H. Cohn, W.B. Lane, and others, 11 Jan. 1957. 24 p. tables, diagrs. UNCLASSIFIED V. NM 006-015.04 11 Jan. 1957. 24 p. tables, diagrs. UNCLASSIFIED V. NM- 006-015.04 The present study was designed to reproduce in the The present study was designed to reproduce in the laboratory an acute exposure of mice to early fallout laboratory an acute exposure of mice to early fallout (2 days old) such as might result from a (2 days old) such as might result from a land-based nuclear detonation. Biological land-based nuclear detonation. Biological data were obtained on the uptake, distribudata were obtained on the uptake, distribu-**UNCLASSIFIED** UNCLASSIFIED tion, and retention of the inhaled fallour tion, and retention of the inhaled fallout (over) (over)

A Martin Association

simulant. Further, as a basis for comparing the effects of different types of simulants, the metabolic behavior of this dry-particle fallout simulant of limited solubility was compared to that of two previously studied fallout simulants, an ionic liquid aerosol and a mud-shurry aerosol. The simulant was also administered by gavage to provide data on uptake of fallout by ingestion as compared with inhalaiton. From these data, an evaluation was made of the radiation dose to individual these from inhaled fallout as compared to the concomitant external radiation done that the animals would receive if exposed to the same airborne simulant.

UNCLASSIFIED

simulant. Further, as a basis for comparing the effects of different types of simulants, the metabolic behavior of this dry-particle fallout simulant of limited solubility was compared to that of two previously studied fallout simulants; an ionic liquid aerosol and a mud-slurry aerosol. The simulant was also administered by gavage to provide data on uptake of fallout by ingestion as compared with inhalztion. From these data, an evaluation was made of the radiation does to individual tissues from inhaled fallout as compared to the concomitant events, radiation done that the animals would receive if exposed to the same also

and a second of the second second

UNCLASSIFIED

timelants. Further, as a basis for comparing the effects of different types of simulants, the metabolic behavior of this dry-particle fallout simulant of limited relability was compared to that of two previously studied fallout simulants; an one liquid aerosol and a mud-slurry aerosol. The simulant was also administered by gavage to provide data on uptake of fallout by ingestion as compared with inhala tion. From these data, an evaluation was made of the radiation dore to individual times from inhaled fallout as compared to the concomitant external radiation done that the animals would receive if exposed to the same airborne simulant.

UNCLASSIFIED

UNCLASSIFIED

simulant. Further, as a basis for comparing the effects of different types of simulants, the metabolic behavior of this dry-particle fallout simulant of limited solubility was compared to that of two previously studied fallout simulants; an ionic liquid aerosol and a mud-slurry aerosol. The simulant was also administered by gavage to provide data on uptake of fallout by ingestion as compared with inhalation. From these data, an evaluation was made of the radiation dose to individual tissues from inhaled fallout as compared to the concomitant external radiation dose that the animals would receive if exposed to the same airborne simulant,

			· ·	
				Ç.
Naval Radiological Defense Laboratory. USNRDL-TR-118 RADIOTOXICITY RESULTING FROM EXPOSURE TO FALLOUT SIMULANT. II. THE METAROLISM OF AN INHALED AND INGESTED SIMULANT OF FALLOUT PRODUCED BY A LAND-BASED NUCLEAR DETONA- TION. by S.H. Cohn, W.B. Lane, and others. 11 Jan, 1957. 24 p. tables, diagrs. UNCLASSIFIED The present study was designed to reproduce in the laboratory an acute exposure of mice to early fallout 2 days old) such as might result from a table-based miclear detonation, Biological	 Fallout - Metabolism Intestine - Effects of radiation Cohn, S.H. Lane, W.B. Title Title: Metabolism NM 006-015.04 	Naval Radiological Defense Laboratory. USNRDL-TR-118 RADIOTOXICITY RESULTING FROM EXPOSURE TO FALLOUT SIMULANT. II. THE METABOLISM OF AN INHALED AND INGESTED SIMULANT OF FALLOUT PRODUCED BY A LAND BASED NUCLEAR DETONA- TION. by S.H. Cohn. W.B. Lane, and others. 11 Jan. 1957. 24 p. tables, diagrs. UNCLASSIFIED The present study was designed to reproduce in the laboratory an acute exposure of mice to early fallout (2 days old) such as might result from a land-based nuclear detonation. Biological data ware obvided on the untake distribu-	 Fallout - Metabolism Intestine - Effects of radiation Cohn, S.H. Lane, W.B. Title Title: Metabolism NM 006-015.04 	
and recention of the inhaled fallout	UNCLASSIFIED	tion, and retention of the inhaled fallout	UNCLASSIFIED	
Naval Padiological Defense Laboratory.	1. Fallout - Metabolism	Naval Radiological Defense Laboratory.	1. Fallout - Metabolism	
ALLOUT SIMULANT. II. THE METABOLISM OF AN INHALED AND INGESTED SIMULANT OF FALLOUT PRODUCED BY A LAND-BASED NUCLEAR DETONA- TION, by S.H. Cohn, W.B. Lane, and others. 11 Jan, 1957. 24 p. tables, diagrs. UNCLASSIFIED The present study was designed to reproduce in the Jaboratory an acute exposure of mice to early fallout (2 days old) such as might result from a land-based nuclear detonation. Biological data were obtained on the uptake, distribu-	L. Cohn, S. H. II. Lane, W. B. III. Title IV. Title: Metabolism V. NM 006-015.04	EADEORUMICATY RESULTING FICH EXPOSURE TO FALLOUT SIMULANT. II. THE METABOLISM OF AN INHALED AND IN-JESTED SIMULANT OF FALLOUT PRODUCED BY A LAND-BASED NUCLEAR DETONA- TION. by S.H. Cohn, W.B. Laue, and others. 11 Jan. 1957. 24 p. tables, diagrs. UNCLASSIFIED The present study was designed to reproduce in the laboratory an acute exposure of mice to early fallout (2 days old) such as might result from a land-based nuclear detonation. Biological data were obtained on the uptake, distribu-	Tadiation I. Cohn, S. H. II. Lane, W. B. III. Title IV. Title: Metabolism V. NM- 006-015.04	
tion, and retention of the inhaled fallout (over)	UNCLASSIFIED	tion, and retention of the inhaled fallout	UNCLASSIFIED	

•

A LANDAL STATISTICS

simulant. Further, as a basis for comparing the effects of different types of simulants, the metabolic behavior of this dry-particle fallout simulant of limited solubility was compared to that of two previously studied fallout simulants, an ionic liquid aerosol and a mud-slurty aerosol. The simulant was also administered by gavage to provide data on uptake of fallout by ingestion as compared with inhalation. From these data, an evaluation was made of the radiation dose to individual tissues from inhaled fallout as compared to the concomitant external radiation dose that the animals would receive if exposed to the same airborne simulant.

UNCLASSIFIED

simulant. Further, as a basis for comparing the effects of different types of simulants, the metabolic behavior of his dry-particle fallous simulants of Hantedryses we solubility was compared to that of two previously studied fallout simulants, an ionic liquid aerosol and a mud-simily aerosol. The simulant was also administered by gavage to provide data on uptake of fallout by ingestion as compared with inhalation. From these data, an evaluation was made of the radiation doe to individual tissues from inhaled fallout as compared to the concomitant ethories does that the animals would receive if exposed to the same alcorate timulant.

UNCLASSIFIED

UNCLASSIFIE

UNCLASSIFIED

imilant: Further, as a basis for comparing the effects of different types of simuthe metabolic behavior of this dry-particle fallows simulant of limited as a second state of the second

UNCLASSIFIED

simulant. Further, as a basis for comparing the effects of different types of simu-

ionic liquid aerosol and a mud-slurry aerosol. The simulant was also administered by gavage to provide data on uptake of fallout by ingestion as compared with inhalation. From these data, an evaluation was made of the radiation dose to individual tissues from inhaled fallout as compared to the concomitant external radiation dose that the animals would receive if exposed to the same airborne simulant.

UNCLASSIFIED

UNCLASS