

the Commission at the information meeting scheduled for  
Wednesday, May 11, 1966.

W. B. McCool

Secretary


DISTRIBUTION

COPY NO.

Secretary	1,25-30
Commissioners	2-6,31-34
General Manager	7 - 8
Deputy Gen. Mgr.	9
Asst. Gen. Mgr.	10
Dir. of Regulation	11
Exec. Asst. to GM	12 - 13
Asst. GM for Admin.	14
Asst. GM for Reactors	15
Asst. GM for R&D	16
General Counsel	17
Biology & Medicine	18
Congr. Relations	19
Inspection	20
Military Application	21
Operational Safety	22
Public Information	23
Reactor Dev. & Tech.	24

SPECIAL REVIEW	Revisions	Class.	Date
FINAL			
13 DETERMINATION			
Class: UNCL			

By: J. Hahn  
5/2/85



UNITED STATES  
ATOMIC ENERGY COMMISSION  
WASHINGTON, D.C. 20545

May 4, 1966

MEMORANDUM FOR CHAIRMAN SEABORG  
COMMISSIONER PALFREY  
COMMISSIONER RAMEY  
COMMISSIONER TAPE

*fr* THROUGH GENERAL MANAGER *John V. Kneig*

SUBJECT: CURRENT STATUS OF THYROID NODULES AMONG THE MARSHALLESE  
EXPOSED TO FALLOUT FROM THE BIKINI TEST OF MARCH 1, 1954

In our last memorandum on this subject, October 11, 1965, it was noted that a cumulative total of 11 instances of nodular thyroid glands had been found in the 69 surviving Marshallese who had been exposed to fallout from the subject test while on Rongelap and on Ailingnae. All but one of these cases were in young people who were less than 10 years old at the time of exposure. The single exception was in a 41-year-old woman (approximately 30 at time of exposure) who was found to have a carcinoma of the thyroid at operation.

Dr. Robert A. Conard of the Medical Department of Brookhaven National Laboratory, and leader of the annual (or more often) medical surveys of these people, has just returned from the latest survey conducted during February and March of 1966. He reports that an additional five cases of nodular thyroids have been diagnosed in this group. The present cumulative incidence is shown in the following table.

Age at Exposure	Number Exposed	Total Incidence of Nodules Through March 1966	
		Number	Percent
1-5	16	9	56
6-10	9	4	44
11-15	7	0	-
16-20	5	0	-
> 20	<u>32</u>	<u>3</u>	<u>9</u>
	69	16	~ 25

CONFIRMED TO BE UNCLASSIFIED  
BY AUTHORITY OF DOE/OC  
*Jose Diaz* 5/21/81  
DATE  
BY: TFEHMER 3/25/86

the remaining 10 cases to Brookhaven for further diagnostic study and possible surgery during May 1966. One is a 46-year-old adult woman and the others are girls presently 13, 13, 13 and 20 years old. They will be accompanied by the Rongelap schoolteacher who will act as interpreter. The travel and treatment of these people has been developed with the cooperation of the Honorable Wilfred Goding, High Commissioner, Trust Territory of the Pacific Islands.

All these Marshallese presumably have been taking tetraiodothyronine, a synthetic thyroid hormone, for the last seven months. The intent of the therapy is to prevent and/or cause regression of the nodules. Some success has been seen, but it is too soon to draw conclusions.

Forty children who were on Utirik at the time and received estimated doses of 14 R plus proportionately smaller internal thyroid doses were examined for nodular thyroids; none were found.

Among 194 unexposed controls who are now living on these islands, three cases of thyroid nodules were found, all in adults more than 47 years. Other forms of thyroid disease were not seen.

You will be kept informed of the progress of these studies; the next noteworthy item will be the clinical and possibly surgical findings in the five Marshallese coming to Brookhaven.

*C. L. Dunham, M.D.*  
C. L. Dunham, M.D.

Director

Division of Biology and Medicine

UNITED STATES GOVERNMENT

# Memorandum

TO : R. E. Hollingsworth, General Manager  
THRU : Howard G. Brown, Jr., AGMA  
FROM : John A. Harris, Jr., Director  
Division of Public Information

DATE: MAY 5 1966

SUBJECT: PROPOSED ANNOUNCEMENT AND RESPONSES TO INQUIRIES ON FIVE MARSHALLESE PEOPLE WHO HAVE THYROID NODULES

DPI:FT

Attached for your approval are a proposed public announcement and responses to inquiries on circumstances surrounding five new cases of thyroid nodules detected in the Marshallese people during the March, 1966, medical survey.

Two upcoming events are likely to receive news media attention:

1. the arrival, this month, of five Rongelap people in this country for clinical examination and possible surgery; and
2. the appearance of an article, in June, in the New England Journal of Medicine on the latest findings by Dr. R. A. Conard of BNL.

We recommend that the announcement be approved for issuance by Brookhaven National Laboratory just before the arrival of the five Marshallese. The questions and answers are for use to the extent necessary in response to telephone inquiry only, or for guidance before interview with media representatives.

Attachments

APPROVED:


\_\_\_\_\_  
General Manager

\_\_\_\_\_  
Date



Upton, New York, May , 1966. Five people from Rongelap Atoll in the Marshall Islands of the Pacific will be brought to the Medical Research Center of AEC's Brookhaven National Laboratory at Upton, L.I., N.Y., this month for clinical examination and possible surgery. They have nodules -- a thickening or hardness -- in their thyroid glands which appear to be the result of exposure to fallout radiation from an atmospheric nuclear detonation at Bikini 12 years ago. Of the five, one is a mature woman, three are girls in their early teens, and one is a young woman under 21 years of age.

The five additional cases, noted early this year during the continuing annual medical surveys of the islanders, bring the total number of thyroid cases to 16 out of the original 82 Rongelap people who were exposed. The thyroid nodules were first detected in March, 1964, during the tenth annual medical survey of the people of Rongelap and Utirik Atolls. Thirteen of 19 children in the more heavily exposed group, all of whom had been exposed at less than ten years of age, had nodules. All nodules of those children who were examined or treated surgically were found to be benign. Only one person, a woman of 40 years, had cancer of the thyroid, for which she received surgery a year ago, and is now doing well. In addition, there were two boys



with hypothyroidism in the exposed group who had previously shown growth retardation. These boys have improved as a result of the thyroid hormone therapy instituted six months ago.

~~Additional amounts of radiation resulted from absorption of radioiodine in the~~  
thyroid glands. Since that time the U.S. Atomic Energy Commission and the Trust Territory of the Pacific Islands have sponsored annual medical surveys of the exposed people. The surveys are carried out by a Brookhaven National Laboratory team, headed by Dr. Robert A. Conard from the Laboratory's Medical Research Center.

Immediately after the 1954 exposure the people were evacuated to Kwajalein Atoll for treatment and were subsequently on Majuro Atoll until 1957 when they were returned to Rongelap. On their return to their home atoll the United States provided them with a new village and other facilities and services to assist their re-establishment. The current population of Rongelap is about 228 persons. Last year Congress appropriated \$950,000 as "compassionate relief" payment to the exposed population.

#

2. Q. What is the total number of thyroid nodules cases detected in Marshallese to date?

A. There have been a total of 16 cases in the 69 people now living of the original 82 people exposed. The deaths that occurred during the past twelve years were among elderly people and were not considered to be radiation-connected. Thirteen of the 19 young people, or 68 per cent, in the more heavily exposed group on Rongelap had nodules. All had been exposed at less than ten years of age.

3. Q. Are thyroid nodules malignant or benign?

A. In all of the children examined surgically the thyroid nodules have been benign. Those not operated on yet are assumed to be benign. The possibility of an occasional malignancy cannot be ruled out, however. One of the people in the exposed group, an adult woman, has cancer of the thyroid which has been treated surgically.

4. Q. Are the nodules in the thyroid radiation-connected?

A. We believe so. Since they have been found in the exposed population much more frequently than in the unexposed group, we must assume that the increased frequency of the nodules is radiation-connected.

one adult woman had a thyroid malignancy which was removed last year.  
She is doing well.

6. Q. What is being done for the additional five Marshall Islanders in whom

A. Arrangements have been made to bring the five to the Medical Department at Brookhaven National Laboratory for further examination and treatment.

7. Q. When were these people exposed to radiation and how many were in the group?

A. A total of 82 Rongelap people (64 on Rongelap Atoll, and 18 visiting on nearby Ailinginae Atoll) and 157 on Utirik who were accidentally exposed to fallout radiation from a nuclear device tested on March 1, 1954.

8. Q. Were there people on the same, or an adjacent, atoll who were less exposed? What were their radiation dose estimates?

A. Yes. The 18 Rongelapese who were visiting on Ailinginae Atoll were lightly exposed. They received an estimated whole body dose of 69 R, and a proportionately smaller thyroid dose. About 40 children on Utirik who were mildly exposed to fallout radiation (14 R) have shown no thyroid gland abnormalities. These children on Utirik are in the same age range as those in the heavily exposed Rongelap population.



9. Q. What is the "unexposed group;" how many people?

A. Some 200 unexposed people now living on Rongelap, including 75 children, are examined as part of the annual medical surveys for comparison purposes.

10. Q. How many children were in the heavily exposed group?

A. There were twenty-five between 1 and 10 years of age at the time of the incident.

11. Q. What was the estimated dose to the thyroid glands of the children?

A. Their thyroids received doses possibly between 200 to 1400 rem largely from radioiodine.

12. Q. What kind of shape are these people in now; and what other delayed effects, besides the nodules, have been observed?

A. The general health and death rate of the exposed population on Rongelap have been about the same as for the comparison group. Other possible residual radiation effects noted in the 82 exposed Rongelapese were: slight retardation of statural growth and bone maturation in two boys exposed at less than 5 years of age; possibly a higher incidence of miscarriages in exposed women during the first four years; slower recovery of some of the peripheral blood elements, and non-malignant changes in some of the areas of previous beta radiation burns of the skin. No definite radiation effects on birth rate, aging, or leukemia have been noted. The numbers of people here are far too small for statistical analysis except in the thyroid nodule case.

14. Q. What was the estimated whole body dose received?

A. The average whole body dose was approximately 175 roentgens.

15. Q. What medical assistance has AEC given?

A. Since the time of the incident the U.S. Atomic Energy Commission and the Trust Territory of the Pacific Islands have sponsored annual medical surveys of the exposed people and have collaborated in seeing that their medical needs have been met.

16. Q. Who runs these surveys?

A. The surveys are headed by Dr. Robert A. Conard of the Medical Research Center at AEC's Brookhaven National Laboratory.

17. Q. What was the thyroid dose received by the exposed adults?

A. The dose to the thyroid glands of the adults from the radioiodines absorbed was estimated to be about 150 to 160 rem.

~~Exposed~~ ~~burns~~ ~~skin~~ ~~around~~ ~~the~~ ~~neck,~~ ~~head,~~ ~~arms,~~ ~~and~~ ~~legs~~ ~~during~~  
the first day of the fallout. These places received burns from  
beta radiation. The burns healed but the pigment, in some in-  
stances, did not regenerate.

19. Q. What is the extent of the present New England Journal of Medicine  
report?

A. The present report summarizes the medical findings over a twelve-  
year period relating to the people of Rongelap Island who were  
exposed. The initial findings are only briefly reviewed; more  
attention is given to the more recent examinations.

20. Q. Are there public reports on the medical surveys in addition to the  
current article in the Journal of the New England Medical Society?

A. Yes. See list below:

After Exposure to Fallout," AEC report BNL-412 (T-80), March 1958.  
(OTS \$0.20).

4. R. A. Conard, et al, "March 1957 Medical Survey of Rongelap and Utirik People Three Years After Exposure to Radioactive Fallout," AEC report BNL-501 (T-119), June 1958. (OTS \$1.00).
5. R. A. Conard, et al, "Medical Survey of Rongelap People Four Years After Exposure to Fallout," AEC report BNL-534 (T-135), May 1959.  
(OTS \$1.25).
6. R. A. Conard, et al, "Medical Survey of Rongelap People Five and Six Years After Exposure to Fallout," AEC report BNL-609 (T-179), February 1961. (OTS \$2.00).
7. R. A. Conard, et al, "Medical Survey of Rongelap People Seven Years After Exposure to Fallout," AEC report BNL-727 (T-260), May 1962. (OTS \$2.00).
8. R. A. Conard, et al, "Medical Survey of Rongelap People Eight Years After Exposure to Fallout," AEC report BNL 780 (T-296), Jan.1963. (OTS \$2.00).

9. R. A. Conard and A. Hicking, "Ten Year Summary of Medical Findings in Marshallese People Exposed to Fallout Radiation," AEC report BNL-8760.
10. R. A. Conard et al, "Medical Survey of the People of Rongelap Island, Eleven Years After Exposure to Fallout Radiation (March, 1965)" BNL-9698.
11. R. A. Conard and Hicking, "Medical Findings in Marshallese People Exposed to Fallout Radiation, Results from a Ten-year Study," Journal of the American Medical Association (JAMA), May 10, 1965, Vol. 192, No. 6, pp 457-459.
12. R. A. Conard, New England Journal of Medicine, June , 1966 (in press).