snp - 270 - 546

HEADQUARTERS JOINT TASE FORCE SEVEN APO 187 (HOW), c/o Postnester San Francisco, California

12 April 1954

E

MELORANDUM FOR RECORD

ľ

SUBJECT: BRAVO Shot, Operation CASTLE

1. <u>PURPOSE</u>: To make a matter of record operational aspects that were considered prior to ERAVO event of Operation CASTLE and to analyze the resultant situation in light of available pre-shot and post-shot information.

2. <u>GENERAL INFORMATION</u>: Operation CASTLE is planned to consist of a series of seven detonations at the Pacific Froving Grounds, which encompasses Eniwe tok and Bikini Atolls. BRAVO is the code name that was given the firing of the first device, SHRIMP, at 0645 M on 1 March 1954, off Namu Island, Bikini Atoll.

Subsequent to BRAVO detonation radioactive debris fell on certain inhabited atolls of the northern Marshall Islands. Radiation intensities rose to levels sufficient to warrant evacuation of four atolls and all personnel were removed from these atolls to Kwajalein in accordance with the operational energency plan of JTF SEVEN. Areas evacuated and gamma dosages received are indicated below:

ATOLL	POPULATION	DISTANCE FROM GROUND ZERO	DOSES RECEIVED
Ailinginae	17	79 NM	80 R (computed)
Rongelap	. 82	100 NM	100-130 R (computed)
Rongerik	28 🖸	133 NM	40 - 98 R (film badge)
Utirik	154	270 NM	17 R (computed)

(#) 28 American Service personnel; 25 USAF Weather Detachment plus 3 USA Signal Corps personnel.

All evacuees are under competent medical care.

3. <u>PREVIOUS EXPERIENCE AND CHARACTERISTICS OF NUCLEAR</u> <u>DETONATIONS</u>: Redioactive debris is an interent characteristic of all nuclear detonations. It originates from fission fragments

ation (Concelled) (tomate the Serie disadicy its at Joint Arrive Deriper + avol said

which are the residue of bomb elements and surface materials, soil and water, made radioactive by accompanying radiation fields. Debris is sucked high into the atmosphere by after winds of the explosion. Where this radioactive debris will fall is a major pre-shot consideration and primarily influences the decision to detonate a nuclear explosion at a certain time.

The area over which radioactive debris is spread and the intensity of fall-out on the ground are determined by the yield of the explosion as well as by wind pattern since the larger the yield, the more surface materials are sucked up into the cloud and the more fission fragments are available. The relationship between yield and fall-out is known only qualitatively.

4. <u>PRE-SHOT INFORMATION</u>: The operational aspects of the BRAVO experience were planned and conceived in the light of experience gained from previous operations. These factors were considered:

a. The basis for forecasting where fallout will go is experience gained from overseas test operations CROSSROADS, SANDSTONE, GREENHOUSE and IVY and to a certain extent from tests at the Nevada Proving Ground. Prior to the firing of BRAVO, only one megaton yield device (IVY-MINE) had been detomated. Although conscientious efforts were made to document the fallout from MINE, only about 5% of the total debris could ever be accounted for.

The technique used for forecasting fallout patterns is to consider the cloud as a small area source (about a 15 mile radius); then add vectorially forecast winds from the surface to approximately 100,000 feet. The next step is to outline an area on the ground where fallout is expected. This area is computed by taking into consideration particle size, diffusion into the atmosphere, wind pattern, yield and source radius. Such patterns have been largely confirmed by experience in Nevada as well as by the meager data available here.

c. The surface radex was plotted, with an insurance factor added, i.e., smaller particles than previous experience indicated necessary were considered. This doubled distances from ground zero where fallout was predicted to occur.

fred incert

ъ.

122.7

d. The upwind intensity of radiation levels at various distances was considered to be of the same order of ragnitude as for IVY-LIKE. Radiation versus distance lines ware transposed to Bikini Atoll.

2

PFAR

SEUNCE

A critical problem in predicting fall-out in-6. , forecasting the stability or lack of stability of the attern after shot time. Since radioactive particle tradetermined primarily by the winds at each level, it is ud that winds must be from favorable directions or very-;hin the outer limits on favorable directions during the fallout. The critical fallout period was considered to the order of twelve to eighteen hours for significant ; to occur. The variation in time arises from consideraof wind shear, with more diffuse and less significant inies at a given time associated with large angular and thear. For this reason, it was required that actual wind sations and forecasts immediately before shot time and __out shot day be continuously considered in their relation :s forecast conditions for the first twenty-four hours ine shot.

a. Weather

Weather conditions during the five days prior to Indicated a favorable trend for BRAVO day with easterly elow 15,000 feet and winds of a southerly component The situation presented at E-6 hours for the subsequent period (18 hours after shot time) was satisfactory. The - period to begin 18 hours after shot time was predicted an unfavorable trend as northwest winds were forecast - 10,000 to 20,000 foot levels.

b. RadSafe

(1) Resultant wind diagrams including latest
winds and forecast winds for H Hour and the 72 hour
> isjectories, which gave a fellout pattern in a narrow
> the east northeast and a wide (140°) sector to the
> the very slow resultant winds. (See Figure 1).

(2) Surface radex, H to H plus 6 hours. (See

(5) Outlooks for:

(a) <u>Bikini</u>: Unfavorable; <u>Eniwetck</u>: Favorable; Favorable, and the native populated atolls in southeast from ground zero favorable, since resultant winds in tion of these areas were considered too slow to not at fallout to the atolls involved.

(b) <u>Tesk Force fleet</u>: Favorable, provided id out at least 50 miles.



e. A critical problem in predicting fall-out involves forecasting the stability or lack of stability of the wind pattern after shot time. Since radioactive particle travel is determined primarily by the winds at each level, it is required that winds must be from favorable directions or varying within the outer limits on favorable directions during the time of fallout. The critical fallout period was considered to be on the order of twelve to eighteen hours for significant fallout to occur. The variation in time arises from considerations of wind shear, with more diffuse and less significant intensities at a given time associated with large angular and speed shear. For this reason, it was required that actual wind observations and forecasts immediately before shot time and throughout shot day be continuously considered in their relation with the forecast conditions for the first twenty-four hours after the shot.

5. <u>PRE-SHOT BRIEFINGS</u>: The following were presented at the pre-shot command briefings:

a. <u>Weather</u>

Weather conditions during the five days prior to BRAVO indicated a favorable trend for BRAVO day with easterly winds below 15,000 feet and winds of a southerly component above. The situation presented at H-6 hours for the subsequent 24 hour period (18 hours after shot time) was satisfactory. The 24 hour period to begin 18 hours after shot time was predicted to give an unfavorable trend as northwest winds were forecast for the 10,000 to 20,000 foot levels.

b. RadSafe

(1) Resultant wind diagrams including latest observed winds and forecast winds for H Hour and the 72 hour cloud trajectories, which gave a fallout pattern in a narrow sector to the east northeast and a wide (140°) sector to the south with very slow resultant winds. (See Figure 1).

(2) Surface radex, H to H plus 6 hours (See

Figure 2).

1

(

((

Ċ

(3) Outlooks for:

(a) <u>Bikini</u>: Unfavorable; <u>Eniwetok</u>: Favorable; <u>Ujelang</u>: Favorable, and the native populated atolls in southeast quadrant from ground zero favorable, since resultant winds in the direction of these areas were considered too slow to move significant fallout to the atolls involved.

(b) <u>Task Force fleet</u>: Favorable, provided ships moved out at least 50 miles.





(e) Air routes through Wake and Kwajalein:

feverable. : (d) Surface routing inside 500 miles considered in its relation to all known transient shipping: favorable.

e. <u>Scientific</u>

X3

(1) High altitude sampling operations - favor-

1

(2) Light transmission for scientific experiments -

6. <u>COECIDSIOSS</u>:

able.

- a. Lack of fallout information from previous shots of comparable yield was a serious handicap.

e. The original source cannot be considered as a point or a relatively shall area but must be considered to be an area of about a hundred miles in diameter. This diameter also depends on yield.

d. The radioactivity of the debris can be considered proportional to yield. Radioactive material in the eloud was thus two to three times was was expected.

•. An appreciable fraction of the observed fallout ean only be accounted for by assuming that it originated in the stratosphere. For such particles to reach the ground at observed times, their diameter must have been in excess of 100 microns.

f. Forecast for shot time winds at shot time was essentially correct. Variation from forecast trajectories was approximately 10 degrees in significant upper levels; unfortunately, the variation was in the wrong direction (See Figure 5). The small variations observed at lower levels were also in an unfavorable direction. Mevertheless, the accuracy of the winds aloft forecast approached the limits of accuracy of the wind observations themselves and were well within the normal forecast error.

g. The fallout pettern extended from the Bikini atoll to the east northeast. Considerable widening of the pattern took place due to diffusion. The intensity of the



pattern on the ground was due primarily to superposition of methods cloud fallout on the stem cloud pattern; and the superposition can be attributed to the narrow cone within which the winds were acting. The theory that a significant fallout does not come from the stratosphere is not substantiated by the facts of BRAVO.

h. For future high yield shots, the forecest and observed winds for the first twenty-four hour post-shot period should receive as much emphasis as analyses made for shot time.

7. **EVACUATION:** Evecuation took place in accordance with operational emergency plan and without incident. Evacuation was not effected prior to detonation because no signifieent fallout was expected on inhabited areas.

•	ALVIN C. GRAVES Beientifie Directo	P. W. CLARKSON or Kajor General, U.S. Army Commender
.6 Inc	វិទ	• •
]. 2. 5.	Pigure 1 Figure 2 Figure 5	•
8. (Tab "A" - Weather Tab "B" - RadSate, Harre	tive Sequence of Events

. Tab =C= - Medical (plus addendum)



Reofinisted Data















1 March 1954

MEMORANDUM FOR RECORD:

SUBJECT: Command Briefing, 1100, 28 February 1954

Members present: Gen Clarkson, Gen McGinley, Dr. Graves, Adm Bruton, Col Cowart, Lt Col Harbour (in lieu of CTG 7.4) Capt Maynerd, Lt Col Bonnot, and Lt Col House.

1. In general, the hodograph gave a forecast surface radex in the low levels (zero to 15,000 feet) at about 50 miles in 6 hours, generally orientated to the west south west in approximately a 30 degree sector. In the high levels, the radex gave a 6 hour pattern centered on approximately 65 degrees in a marrow cone approximately 20 degrees wide.

2. The 72-hour cloud trajectories indicated all levels in a narrow band generally 65 degrees true from ground zero except the 10,000 foot level moving generally east through north to west and a 60,000 foot level moving southwest and then west.

3. RadSafe outlooks were given as follows:

- a. Bikini atoll: Favorable.
- b. Eniwetok atoll: Favorable.
- c. Ujelang atoll: Very favorable.
- d. Native atolls in southeast quadrant: Very favorable.

e. Control DDE: It was recommended that the Controll DDE be moved from a position 90 miles West of Ground Zero to a posttion 230 degrees True, 90 miles from Ground Zero.

f. ATF and the YAGS: The plot of the YAGS and ATF courses from about H minus 24 hours to H plus 2 hours was indicated. In general, from about H minus 5 to about H minus 3 hours the ATFs and YAGS courses run approximately 15 to 20 miles from the armed device and the ATF departs from a position on the edge of the low altitude radex at about H minus 12 hours to a position effort imately 35 miles south of ground zero at H plus 2. The horests of the YAGS and ATF were presented for whatever consideration was necessary in view of their close proximity to the device and the radex.





g. No closure was recommended on air and surface routes through Wake and Ewajalein.

h. No transient shipping was reported within 500 miles of Ground Zero. It was pointed out that 3 minus 2 and 3 minus 1 P2V sweep had been sent out on headings of 300 degrees and 330 degrees respectively. This was based on earlier forecasts. Since the winds shifted around to a narrow band to the ENE, it was recommended that a B day P2V sweep along a bearing line approximately 65 degrees be laid on.

'i. It was recommended that CINCPAC be advised of the following:

(1) 72 hours trajectories.

(2) Very favorable outlook for native populations.

(3) No interference of the air and surface routes.

j. It was recommended that no change be made in the position of the task force ships.

4. In summary, it was recommended that RadSafe conditions be considered very favorable on all points, but only favorable at Bikini shot atoll.

R. A. HOUSE Lt Col., USAF RadSafe Officer

2



1 March 1954

MELIORANDUM FOR RECORD:

SUBJECT: Command Briefing, 1800, 28 February 1954.

L'embers present: Gen. P.W. Clarkson, Dr. A.C. Graves, Jr. Ogle, Dr. D. Sewell, Gen. E. L'CGinley, Lr. J. Reeves, Col. Cowart, Capt Haynard, Lt Col Bonnot, Lt Col House.

1. In general, the 1100, 28 February 1954 briefing was confirmed except indications were presented that conditions were getting less favorable.

2. It was agreed to let the execute decision remain firm until the midnight briefing.

I

R. A. HOUSE Lt Col., USAF Rad Safe Officer

1 March 1954

「おいたのできた」」」

a manager and and an a set the set of the se

MELOFANDUM FOR RECORD:

ι.

ï

SUBJECT: Command Briefing, 0000, 1 March 1954

Kembers present: Gen. P.W. Clarkson, Gen. E. McGinley, Gen. H. Estes, Dr. A.C. Graves, Dr. W. Ogle, Mr. J. Reeves, Dr. D. Sewell, Capt. W.L. Knickerbocker, Col. W.S. Cowart, Capt. R. n. Maynard, Lt Col C.D. Bonnot, Lt Col R.A. House.

1. In general, the forecast presented at the midnight briefing was confirmed, except that in the levels between 5 and 15 thousand feet the forecast was light and variable. Th an attempt to delineate direction to these winds, it was determined that the best forecast that could be given was for the 10 thousand foot level. This was forecast to be westerly at 10 knots as the most pessignistic situation. Consequently the hodograph plot was made using the 10 thousand foot westerly wind in order to present the most pessimistic situation which would occur. This picture gave resultant winds in the direction of Rongelap and Rongerik, however, it was considered that the distance to Rongelap and Rongerik compared to the resultant wind speeds were such that no fell-out should reach those atolls. From the forecast hodograph the time of travel to Rongelap would have been about 12 to 15 hours. The hodograph plot did however show that certainly TARE site would be heavily contaminated, and most likely NAN. The hodograph gave two general fall-out The lower level running from 260 degrees around through areas. south to 90 degrees with a six hour fall-out line in the direction of the populated atolls in the southeast quadrant about 15 to 20 miles out from ground zero. The high level radex ran from about 45 degrees to 80 degrees with a six hour fall-out line to 70 miles. (Since the 6 hour fall-out lines were computed on about 100 micron particle size, it was recommanded that the distance be doubled for safety. This amounted to considering particle sizes down to about 70 microns).

2. No change in the 72 hour cloud trajectories.

- 3. RadSafe Outlooks were modified as follows:
 - a. Bikini atoll was changed from favorable to unfavorable.
 - b. Eniwetok stoll remained very favorable.
 - c. Ujelang atoll remained very favorable.



d. Native atolls in the southeast quadrant were discussed at this point and elsewhere in the briefing. The net result of the forecast was that these atolls should remain favorable due to the forecast long time of travel for fall-out to these places. Specifically, Wotho and Rongelap were considered by name and position, these being the closest native populated atolls in the vicinity of ground zero.

e. Control DDE: No change recommended over the change made at the 1100, 28 February briefing.

f. ATF: It was recommanded that the ATFs E_{f}^{2} hour position be changed from 35 miles outh of GZ to at least 50 miles south of GZ.

g. No closure was recommended on air and surface routes through Wake and Kwajalein.

h. No further shipping was reported within 500 miles of GZ.

i. An advisory to CINCPACFLT and CINCPAC was recommended in conformance with the above.

j. Task Force Ships: It was recommended that, due to the close proximity of Task Force ships to the outer edge of the six hour fall-out, these ships be moved further out on a radial line to at least 50 miles.

4. In summary, it was recommended that RadSafe conditions be considered favorable on all points, and unfavorable for Bikini shot atoll.

R. A. HOUSE Lt Col., USAF Radzafe Officer 2

1 Earch 1954

KENORANDUM FOR RECORD

SUBJECT: Final Weather and RadSafe Check, 0430, 1 March 1954

L'embers present: Gen P. W. Clarkson, Gen O. P. Weyland, Dr. A. C. Graves, Gen. H. Estes, Dr. W. Cgle, Dr. D. Sewell, Mr. J. Reeves, Col W. S. Cowart, Capt R. H. Maynard, Lt Col C. D. Bonnet, and Lt Col R. A. House.

1. The original forecast made at midnight (both weather and radsafe) was considered the best available condition for shot time.

2. The general recommendation for this briefing was one of minimizing the effects of the low level northerly and westerly winds. The recommendations as given at the midnight briefing were restated.

R. A. HOUSE Lt Col USAF RadSafe Officer







DISCUSSION OF OFF-SITE FALLOUT

Fallout off-site followed the pattern immediately established at and adjacent to the proving ground where the cloud in general moved east north easterly with prevailing winds. Task force ships southeast of NAN received the first fallout, being on the southern edge of the main strip of contamination. Fallout on the ships ranged from intensities of 1500 milliroentgens per hour on the BAIROKO which was closest to the center of the fallout path to a few milliroentgens per hour on vessels farther south.

Fallout began at Rongerik Atoll at 1348 hours, 1 March as shown by a self-recording radiation detection station placed there by the NIDO AEC and operated by personnel of the Air Weather Station. This instrument went off scale at 100 mr per hour at 1418 hrs 1 March. Based upon query from air weather personnel a monitor was dispatched with the supply PBK on the morning of 2 March. A ground reading of 2000 mr/hr was obtained at 1115 by the monitor who evacuated 8 of the personnel on his own initiation and recommended evacuation of the remainder as soon as possible based upon the high radiation levels. This was concurred in and the remaining 20 were evacuated by FEL at 1645, 2 March. Calculations estimating the dose received indicated that personnel evacuated at 1115 would have received 85 r and the remainder 95 r. This was in fair agreement with readings of film badges on personnel. Varimum film badge reading was 98r representing 3 men, 52r for 1, 44r for 1, 40r for 9. Average dose for all personnel, 54 romsgens.

Inasmuch as the data from Hongerik is the only data showing exact time the fallout occurred at any location east of the proving ground and adjacent to populated islands affected by substantial radiation, its importance is such that calculation of dosages received by native populations are based upon it for time of fallout in those locations. A detailed analysis of this data is therefore appended in the medical tab.

Survey of Rongelap was made by Pattern ABLE of Security Patrol Squadron (Patron 29) with NIOO Scintameters aboard on 2 March and found an estimated reading of 6750 mr/hr. (Later calibration for aerial survey equipment revised this to 1350). Based on this and its proximity to Eongerik it was decided that it would probably be necessary to evacuate the atoll. Consequently, the PHILIP, DDE 498, was dispatched to reach Rongelap on the morning of 3 March. In the meantime a PEM was sent to ground survey the atoll and at 1830 hours, 2 March, their ground survey showed a reading of 1400 milliroentgens per hour. Calculation of 2005 to 3 March indicated about 110 roentgens so the order to evacuate was given. The destroyer PHILIP evacuated the natives by 0935 to 1025, 3 March. Calculation of the total cose indicated that 130 roentgens were received. Detailed calculations are appended in the medical tab. A total of 65 natives were removed, 16 natives (the old and sickest) by FBM and 49 by DDE.



and the second second and the second second second second



(] (] Inasmuch as some natives of Rongelap were reported to be on Linginae, the atoll was surveyed, 17 natives located, and an intensity of 445 mr/nr was found. Evacuation was complete by 1800, 3 March. The cose computed for this group is 80r and details are appended.

Lerial survey of Utirik by ABLI Fattern indicated 620 mr/hr at 1651 hrs, 2 March (later re-calibrated to 240 mr/hr). On 3 March 1345 ground survey indicated 160 mr/hr. Decision to evacuate based upon fact that estimated dose at time of earliest evacuation would be 13r. Evacuation complete 1245 hrs, 4 March. Estimate of dose to actual evacuation time was 17r. 154 natives were evacuated. Calculations appended in the medical tab.

The only other populated atoll which received fallout of any consequence at all was Ailuk. ABLE Fattern indicated 95 mr per hour at 1845 hrs, 2 March. Based upon the best estimate of fallout time it was calculated that a dose to infinite time would reach approximately 20 roentgens. Balancing the effort required to move the 400 inhabitants against the fact that such a dose would not be a medical problem it was decided not to evacuate the atoll.

Indications from aerial surveys indicated substantial fallout occurred on the unpopulated islands of Eikar and Taka.

Very minor fallout occurred in a southwesterly and westerly direction on Eniwetok and Ujelang Atolls but levels did not exceed lOur per hour at Eniwetok nor 3mr per hour at Ujelang. This was apparently very fine particulate matter carried by the low trade wind component.

A detailed plan was made to make ground surveys of all islands which had fallout in excess of lOmr per hour at estimated fallout time to provide information as to decay rate and verification of estimation of asses. Water and soil samples from these surveys were shipped by air to HASL, WY Operations Office, Attention: Wr. Merril Eisenbud, for detailed analysis.

CLUDE

R. A. HOUSE Lt Colonel, USAF Ch. Tech. Br., J-3



2



CLOUD TRACKING OPERATIONS FOR BRAVO

8 Larch 1954

1. SUMMARY:

The BRAVO Air Rad Safe Operations were conducted essentially as planned. No hazardous air contaminations were encountered by aircraft other than the samplers. Several aircraft and crews were exposed but the levels encountered appear acceptable from both a health and a decontamination aspect. Communication and control difficulties made it difficult if not impossible to forecast the fallout in the Rongerik/Rongelap areas. Steps have been taken to remedy the defects which became apparent during ERAVO operations.

2. GENERAL:

Cloud tracking information for BRAVO was derived from five sources. The manner in which each of these functioned during BRAVO will be discussed individually in subsequent paragraphs. These sources were as follows:

> Sampling aircraft reports. Sweet-sour reports. Special Cloud tracking flights. Weather reconnaissance flights. AFOAT-1 flights.

3. SAMPLING AIRCRAFT REPORTS:

These reports were monitored and recorded by Rad Safe personnel aboard BOUNDARY TARE from plus two thru plus seven hours. Information derived from these reports indicated the sampling aircraft were working the South and Southeast edge of the cloud and therefore stayed in the immediate vicinity of Ground Zero. Because of the altitude of the sampling operations (30,000 - 45,000 ft) there is little relation between the operation of these aircraft and subsequent air or ground contamination. This data, nowever, does assist the Air Rad Safe Officer in obtaining an overall picture of the dispersal of radioactive material. No reports of cloud movement were received from the control E-36 aircraft.

4. STEET-SOUR REPORTS:

ſ

These reports are submitted by any aircraft encountering radioactive contamination and not reporting by other means. No such reports were received during BPAVO. This is not surprising since aircraft other than the samplers and trackers (reporting by other means) seak to avoid areas in which contamination is suspected.



5. SPECIAL CLOUD TRACKING (WILSON) FLIGETS:

The first of these flights, Wilson 2, was directed by Rad Safe to pre-planned post-shot search. This required the aircraft to remain in a holding pettern track approximately 50 miles West of Ground Zero from plus two to plus five hours. This portion of the track was designed for the purpose of derecting radioactive cloud movements toward Eriwetok Atoll. 02 BRAVO the maximum activity encountered was less than 15 mr/br. Through a misunderstanding of control procedures, Wilson 2 overstayed in the holding pattern. When Rad Safe realized this fect, CIC was requested to order him into the previously designated search sector at once. The delay, however, resulted in this air-craft being well behind and to the North of the cloud segments that must have caused fallout on Rongerik and Rongelap. At 1550: the aircraft reported its maximum reading during this flight. 7:18 was reported as being between 500 and 1000 mr/hr approximately 150 nautical miles from Ground Zero at a bearing of 60 degrees. This and the subsequent data appeared to verify the forecast cloud trajectories which indicated the upper cloud segments would leave the PPG on an approximate bearing of 70 degrees, thus avoiding the populated atolls. Wilson 2 subsequently reported in-flight difficulty with the instruments used.

b. On the basis of the results of the Wilson 2 flight the second tracker, Wilson 3, was instructed to search the same general area but to proceed further East to define the rate of cloud movement. At approximately 2000M information was received indicating the possibility of some contamination in the Rongerik/ Rongelap area. A message was immediately dispatched to Tg 7.4 requesting Wilson 3 to alter his search area in such a manner as to cover the populated area to the East. Communication delays prevented Wilson 3 from complying with the request. This plus the fact that no exact instrument readings (instead a range of readings) were reported made interpretation of cloud tracking data difficult.

c. Subsequent Wilson flights (for plus one day) were cancelled when it appeared that no air contamination problem existed at that time.

6. WEATHE RECONNAISSANCE FLIGETS:

Two Petrel Juliet Weather reconnaissence flights were flown on plus one day. These flights were flown to the South and to the Southeast and indicated essentially zero air contamination.



2

7. AFOAT-1 FLIGHTS:

<u>ن</u>ه م

ſ

AFOAT-1 sponsored flights from Hawaii indicated a meximum air contamination of less than 1 mr/hr in that area, (3 March(. Similar flights from Guam reported tenths of an mr/hr as a maximum reading. This was encountered 4 March, 100 nautical miles west of Ponape at 5000 ft.

8. INFLIGHT EXPOSURES:

As expected, several aircraft, including samplers, cloud trackers, evacuation aircraft and P2V security sweep aircraft, encountered areas of air contamination. In all cases it appears that the exposures were well under task force limitations for a health point of view. Standard decontamination procedures are expected to be effective so that all aircraft should be returned to service well prior to the next shot. The P2V security sweep sectors will be modified on future shots to reduce the possibility of contaminating these aircraft.

9. <u>CONCLUSIONS</u>:

a. The Air Rad Safe operations for BRAVO were generally successful but several changes in procedures are being made (see below) to provide more timely and accurate data.

b. No hazardous areas of air contamination were encountered although fallout in the Rongerik/Rongelap area would make it probable that such contamination did exist for a short period of those atolls.

 \boldsymbol{n}

c. Improved monitoring, data reporting and communications facilities are required.

d. Lower search altitudes may improve the ability to correlate air contamination with subsequent fallout.

e. No hazardous fallout appears likely in the Eawail, Ponape or Guam areas.

f. In flight exposures of Task Force personnel appear well within established limits.

10. RECONTENDATIONS:

a. A CW contact is required between BOUNDARY THE and WILSON aircraft to insure better data reporting and control. (This has been requested).

b. A TIB radiac instrument should be carried on all WILSON aircraft and exact radiation reading should be reported. (Eas been arranged).



c. Lower altitudes should be employed in tracking operations. Will be specified in future vector messages.

(

 $\left(\right)$

(

nall PAUL R. MIGNALL Colonel, USAF Air Rad Safe Officer

CONCIDENTIAL'

10 March 1954

MENORANDUM FOR RECORD

SUBJECT: Protection of Transient Shipping During Operation CASTLE

1. In order to provide protection for transient shipping in the region immediately outside the Eniwetok/Bikini Danger Area during Operation CASTLE planning factors were established and a plan of action placed in effect as follows:

- a. Planning factors:
 - (1) CASTLE clouds more than 24 hours old should not be hasardous.
 - (2) 24 hour travel of a CASTLE cloud should be approximately 500 nautical miles.
- b. Plan:
 - The Commander in Chief, Facific Fleet was requested to make advance diversions of shipping outside a sector area from southwest clockwise through north to east to 500 nautical miles from ground sero from H to H plus 24 hours.
 - (2) P2V aircraft were planned to sweep the significant forecast sector of cloud travel, using visual and search radar methods of sightings out to 800 nantical miles on D-2 days, out to 600 nautical miles on D-1 day and, if necessary, in front of the cloud on D day. P2V aircraft crews were directed to effect diversion on all ships sighted in the sector area on D-1 and D day.
 - (3) WB-29 aircraft on routine weather reconnaissance missions were directed to report all sightings of surface shipping encountered. All sightings were to be relayed to the Radar center (CIC USS BAIRDED) in the TG 7.3 fleet.
 - (4) P2V aircraft and destroyer security sweeps were directed for the Eniwetok/Bikini Danger Area. Information from these sweeps was channeled to the Radar center (CIC USS BAIRORD).

MEMORANDUM FOR RECORD (Contid)

SUBJECT: Protection of Transient Shipping During Operation CASTLE

- (5) Information from all the above sources was channeled into the task force headquarters for evaluation and consideration at the Weather/Radsafe Command Briefings.
- 2. The results of the above efforts for ERAVO were as follows:
 - a. All known transient shipping was diverted outside the hazardous fall-out area. The Petapsco (AOG-1) was sailed from Eniwetok to be out of the bazardous area by shot time. The Trust Territory ship M/V Roque was operating outside the designated sector (from Kwajalein to Utirik), arriving at Utirik on the morning of 2 March 1954 and departing Utirik on 3 March 1954. The Roque was subsequently located and monitored at Majuro and found to have insignificant levels of radiation. The Merapi was enroute from Honoluln to Eniwetok but well outside the designated sector area at shot time. The Merapi was monitored upon arrival at Eniwetok and found not contaminated.

ころ、ころにないで、「「「「「「」」」」

b. Based on the forecast significant cloud travel (forecast made on the night of B-3 days) the P2V sweep for B-2 days was directed along true bearing 300 degrees from ground sero. Ho ships were sighted on this sweep. Based on the B-2 day shot time forecast, the P2V sweep for B-1 day was directed along true bearing 330 degrees to a distance of 375 nautical miles. The reduction in distance was based on forecast reduction in resultant wind speeds. This sweep comtacted the USS General Patrick at 17-318, 162-C3E on course 266 degrees, speed 16 knots, at 1204M, 28 February 1954. As she would clear the designated sector by shot time, she was not diverted by the patrol aircraft. Based on a re-forecast (made on B-1 day) of the significant cloud movement for B day, it was decided to search in advance of the cloud along bearing 65 degrees true from ground sero out to 600 nautical miles. Two P2V's were used. The first of these became contaminated early in its mission and was forced to return to base. The second was directed to pick up the search in the approximate location of the previous abort and carry it out to the 600 miles. The only contact reported by these aircraft was the Patapsco (AOG-1) sighted at 12-31N, 170-483, at 1935M, 1 March 1954, course 30 degrees, speed 10 kmots. The Patansco was turned to an easterly heading at 2030E, 1 March 1954. The Commander in Chief, Pacific Fleet was advised later to have the Patapaco monitored upon arrival at Honolulu in the event a check erroute could not be accorplished.

CONFIDENTIN

MEMORANDUM FOR RECORD (Cont'd) SUBJECT: Protection of Transient Shipping During Operation CASTLE

- c. Between 26 February and 1 March 1954, WB-29 aircraft performed weather and cloud tracking missions in all four quadrants from ground zero. No surface shipping was sighted on these missions, three of which were flown to the east northeast of ground zero, one on 27 February and two on 1 March 1954.
- d. The P2V and destroyer search of the Eniwetok/Bikini Langer Area made m contacts. As a matter of interest, a destroyer security sweep on 17 Feburary 1954 encountered one Japanese fishing vessel, the Miyagikenajinoiokompiramaru, 26 mantical miles on true bearing of 40 degrees from Eniwetok Island. This ship was escorted toward the northern edge of the Danger Area and left on course 315 degrees, 9 knots with the recommendation that air patrol observe its subsequent movements. No further contacts with this ship were reported.
- e. As a summary, the CIC BAIROKO was contacted periodically pre-shot and reported no transient shipping in the area.

R. A. HOUSE Lt Col, USAF Chief, Tech Branch, J-3

PATTERN OF FALL-OUT FOLLOWING BRAVO EVENT (Combined Analyses - Immediate and One Week after BRAVO)

1. <u>General</u>. The pattern of ultimate fall-out of radioactive particles has been established utilizing in the cases of the most critical area (i.e. bearing about 050° True, clockwise to 120° True from Ground Zero) the following:

a. Aerial survey by P2V employing NYOO-ABC survey equipment, with readings in mr/hr extrapolated to ground level.

b. Known ground readings taken at some atolls (early and later) used with their time and intensity (actual observations) to get a feeling for the overall situation.

c. Resultant wind pattern to establish best wind for period from H minus 1 hour (USS CURTISS Observation - BIKINI) to H plus 8 hours (Rengerik sounding) together with the H minus 32 hours (Rongerik 0300M) to piece together the wind pattern above the tropopause.

d. Since the Rongerik (NYOO-AEC) survey meter trace established initial time of arrival of fall-out, this time was used in coordination with resultant wind at the cloud level which passed over Rongerik. This level was 25,000 feet vector. Its average speed from Ground Zero calculated from resultant wind plot was 10.4 miles/hr. At first, there was considerable difficulty in making fall-out arrive from the stem of the atomic cloud ()-55,000 feet-tropopause) at Rongerik in 8 hours. The 10.4 miles/hr above would make cloud arrive at Rongerik at about H plus 12 hours. However, by the method of plotting the entire cloud height (which is believed to be about 100,000 feet) for which there were available winds to 95,000 feet, and with the assumptions listed below in constructing shadows (fall-out) of stem and mushroom, there are obtained 2 areas - elliptical in shape, generally east of Ground Zero and superimposed on each other (Appendix I). The suggested fallout area (blue) for the stem is oriented about 070° True from point 52 of Ground Zero, distant 35 miles and with a 200-mile major axis, 100-mile minor axis with a series of extremely hot elliptical envelopes emanating from Ground Zero out to about 110 miles. Superimposed on this area (red) is the suggested mushroom fall-out pattern which is an ellipse oriented 080° True, 40 miles from Ground Zero, major axis at least 180 miles, minor axis 45-90 miles. It is assumed that the cloud diameter in the mushroom for the period in question was at least 70-100 miles. This shows therefore, that the early fall-out at Rongerik could come easily from the mushrooz - large particles by E plus 8 hours, and since the superimposed fall-outs from stem and mushroom cross the northern half of Rongelap Atoll, one would expect these islands to be exceedingly high with their radiation levels. This night be likened to scattaging of the hot stem material by large particles from the tropopause and above. However, the major hot fall-out element must come from the stem debris.

The assumptions used in the rough construction of the ellipse for the whole cloud are: (LtCol Lulejian report on Fall-out - ARDC, SECRET, RD):



1

, •



(1) If wind shear is $< 10^{\circ}$ for the levels in question, sinor axis is 1/8 to 1/4 of the major axis (which is the entire vector for levels looked at). (This is case of 0-5,000 feet winds and 5-20,000 feet winds.)

(2) If wind shear is $>10^{\circ}$ but $< 120^{\circ}$, minor axis is $\frac{1}{2}$ of asjor axis. This is case for 20,000 feet winds to tropopause, and somewhat less for 65,000 to 95,000 feet winds.

(3) If shear at levels making up vector under study is $>120^{\circ}$ draw circle with diameter = to entire resultant vector.

CONCLUSIONS:

 \int_{i}

1. From overall fall-out picture, it is concluded that fall-out may have reached Rongelap Island and Ailinginae later than the pessimistic time of H plus 5 and H plus 4 hours, respectively.

2. From initial land survey reports on Rongelap Atoll with levels at Erippu Island (NE part of Atoll) still at 2.8 to 3.5 r/hr on B plus 7 days, the picture for heaviest fall-out patterns north of this area is established. The relatively light fall-out at Utirik (ESE of the hot area), higher levels of intensity at Bikar (East and downwind of the hot area, i.e..o r/hr at H plus 33 hours almost in downwind line with the superimposed ellipses or hot areas but definitely beyond the hot shadow), confirm the belief in the assumed area of hot fall-out pattern above. Notho (SEE of the area and from Ground Zero) received practically nothing because resultant vector wind speed from the stem and, perhaps some of the mushroom fringe, was so low in velocity through the SE to South from Ground Zero. Enimetok received at about H plus 11 hours a build up to about 10 mr/hr for a period of about five hours.

3. This type of analysis gives a feeling only for pattern of fall-out because it does not tell exactly when the fall-out arrives. However, it is apparent that the 200-300 plus roantgens lifetime dosage line passed on or close to Ailinginae, Rongelap Island and Rongerik which are at 80-100 miles in cases of Ailinginae and Rongelap and 130 miles to Rongerik from Fround Zero. The 1,000 plus roantgens lifetime dosage lines are exceeded as one goes north from Rongelap Island to northern islands of that atoll. This analysis is based on: (1) logical use of wind patterns existing during shot time to fall-out, (2) multiple shot (tower or ground) fall-out pattern data from Nevada Proving Grounds over last 3 years, and (3) experience and data from IVY-WIKE (limited cross-wind and upwind) and CASTE-BRAYC itself.

4. Rongerik radiation intensity levels are known at caset and evacuation time; calculated roentgen dosage agrees with actual observations from film badges at this site

5. The heaviest fall-out pattern was expected to pass north of 12: and east northeast from Ground Zero.

2

Inclosure 6

6. The levels of radiation intensity at the distances of Rongelay and Rongerik were much higher than expected, and sooner than expected since necessary information in this range of yield for surface shots today is the result of some scaling, up from much lesser yields, interpretation of upper wind field patterns, coupled with forecast changes and experience of individuals with such limited data as IVI-WIKE, GREENHOUSE-DOG, EASY, GREENH and ITEM.

7. After seeing BRAVO cloud project pictures (taken from an airplane) with large quantities of visible particulate matter falling through the cirrus deck above the camera plane from tropopause and above, wind data to great heights (i.e. up to at least 100,000 feet) is a must for shot time since the fall-out problem for surface or near shots of large yields can be a definite function of the mushroom as well as the very hot stem of the cloud.

3

Appendirs

- I Plot of General Fall-cut Pattern.
- II Forecast and Observed Hodographs, B-2 to B/7 days.
- III Tabulation of Time of Arrival Data
- IV Hodograph Trend Data

S. S. L. S. S. S. S. S. S.

CAPT. JSN

JT........

6. The levels of radiation intensity at the distances of Rongelar and Rongerik were much higher than expected, and sconer than expected since necessary information in this range of yield for surface shots today is the result of some scaling, up from much lesser yields, interpretation of upper wind field patterns, coupled with forecast changes and experience of individuals with such limited data as IVY-KIKE, GREENHOUSE-DOG, EASY, GEORDE and ITEM.

7. After seeing BRAVO cloud project pictures (taken from an airplane) with large quantities of visible particulate matter falling through the cirrus deck above the camera plane from tropopause and above, wind data to great heights (i.e. up to at least 100,000 feet) is a must for shot time since the fall-out problem for surface or near shots of large yields can be a definite function of the mushroom as well as the very hot stem of the cloud.

3

Appandixt

- I Plot of General Fall-cut Pattern.
- II Forecast and Observed Hodographs, B-2 to B/7 days.
- III Tabulation of Time of Arrival Data
- IV Hodograph Trend Data

R. H. HATYIRD CAPT, JSN

Inclosure 6




































.



.





Dearmuled Drive

AIONIC ENENGT ACT - 1948

BRAVO
11000



I LACE	TIJE BEALLIG	DIGTANCE NAUT.MI.	TILE FALLOUT ARRIVED	INTENSITY LR/IR	DCSE	DATEASITY NR/HR	TIME HRS	Intensi Mr/HR	TY TINE HKS/DAS	AVERACE RESULTANT WIND SPEED TO TROPOPAUSE
Lae	16U	174	/39hrs			a 08		. •		3 2
Ujac	170	168	737			0.1				3 2
Viotho	154	100	<i>4</i> 13.2	2.3	.17	1.0				3.4
Ailinginae	115	79	3.6	12,000	220	445	458	700 .	129.7	9.0 8.1
Kongolap	109	100	5.6	9,800	350	*1400	+36	1350	30.9	8.9
Enlastok	105	98.5	5.4	46,800	4,200	*3000	453.2		2-07	8.9
Rongerik	098	133	* 8	9,000	350	2000	128.5			10.4
Taongi	U52	280		·	•	1.4	32.6	Fringe	effect on	ly
Bikar	084	285	16.3	1,400	120	600	33.0			14.4
Utirik	096	270	21.6	450	48	160	<i>4</i> 55	53/10	46.3das	10.2
Taka	097	261	20.3	290		160	<i>f</i> 34	-	-	10.4
Mejit	104	340	30.2				• • •			9.6
Ailuk	107	291	27.1	110	13	76	34.8			8.9
Jono	110	268	24.8			18	34.7			8.8
Likiop	115	2,2	26.2			6	34.9	1		8.1
Namu	142	265				•016	48.5			4.2
Ailinglapap	μ,2	338				•08	48.9	1		4.8
Arno	127	1,61,				0.6	51.6			5.3
Lia juro	128	1,1, L				2.0	51.4	•		5.5
Aur	122	100			•	•36	50.9			5.9
In Toerab	117	387	42.3			3.6	50.6)		7.1
Erikub	119	321	40.0			4.0	50.2			6.8
liotje	114	321	39.0			20.0	50.0)		6.0
Kucaio	200	225				•5	54.4	•		
Fonape	235	503				•				
Ujelang	245	286				2.0	49.5			
Pingolap	220	426								

* Reading on ground.

•

"DIATCO

NETHOLE

1:

SCORE



AFFEIDIX IV

WINDS ALOFT													
120-	1		1 -	<u>† –</u>	+ -	t	1 -	<u>†</u> —	† •	+ +	†		-12
115-		- 22	1	† − .	↓ ≥		4 -	-	┥ ── ╷	- ·	-	-	-11
110-5			-	-			4	- 1	↓ - ·			↓ → ⋅	
105-2	14- <u>3</u>	7	- 1			18-		ł —	ł — ·	┨ ── ┤	/ /		- 10
100 -		-	-	4 -	-	- 1	-						-10
ຼົ 3 5 -	1 - 1		<u> </u> -	1 -	-0	8.5-	+	4 -	i —	-	} − ·		- 95
90 -	$\mathbf{H} - \mathbf{H}$		<u>+</u> –	- 1	-0	739-	+	↓	—	- i		-	- 90
85 -	+ - +	· — ·	1 -	4 -		3 ≉7—	+ -	- ·		-	-		- 85
80 -	$\mathbf{H} = \mathbf{H}$			-	-08	3 <u>2</u> 2—	-	} −		-	-	-	- 80
75 -	$\mathbf{I} - \mathbf{I}$		-	₩ -			₩ - 1	4 —	-				- 75
70 -	1 - 1	-		 -	-09		1 -	f		-			- 75
65 ~	1 - 1		-	† .−	05	5 7 3 —	+			-		-	- 65
60 -	1 - 1	-	-	- 1		ip. —	ł — ·	{ {	-			-	-60
55 35	11-28	15-13	POPADS	<u>s</u> –	30	13-	ł — ·		-	-		-	- 55
50-27	80-2	66		- 1	-27	36-	·	—	-			-	-50
45 2 5	- 27	57-			-25	40-	 − ·		-			-	- 45
40 25	1-23	<u>1</u> —	-	- 1	-25	41-	† †	—	—	-		-	40
_ 35 26	k7-2\$	4-	-	-	25	33-			-	_	-	-	-35
30-25	29-275	4-	-	-	24	29-		—	-				-30
25 2 5	21-252	7_	-	-	-23	27-	{ — }		-	-			-25
20 - 29	<u>i</u> 7−3¢	4-51	RIFICA	T-	-30	20-	{ ─ }	·	-		-	_	-20
18 -	$\mathbf{f} - \mathbf{f}$		নহাল	ř —	} − ·	+ -			—			-	- 18
16 -		,— ł	-				4 4		-			-	- 16
74 -	┥ <u>─</u> ⁻′┲	-	-			1 ¹¹¹	ł — ł	—				-	-14
12 -	1 - 1		-			· · · ·		—	-			_	- 12
10 3 0	10-28D	9	-	-	-91	06		—	-	-		-	- 10
9 -	$\mathbf{H} = \mathbf{H}$	- +			-		<u> </u>					-	- 9
8 -	$\mathbf{h} = \mathbf{h}$	- +	-		-		—		-	-		· —	- 8
7 -		- +	-				—		-			-	- 7
6 -		- +	-	-				-	-			-	- 6
5 30	5-071	∟ †		-	-09	12-		-	-			-	- 5
4 -		- +	- +	-		<u> </u>			-			-	- 4
3 -		- +		-	-	-			-	-		-	- 3
2 - {		- {	- {	-	<u> </u>				-	-		-	- 2
15-		- +			-			· -		- +	- +	-	– 15
ı – 🛉		- +	- +	-	-	—		- +	- +		· - }	-	- 1
SFC-10	is-on	1	- 1	-]]	—	-				-]]	- SFO
hree	Z	2	Zi	21	Z	ZI	ZI	Z	ZI	Z	Z	Zi	ZI
STA					E 67	A NO		UAT	E				

APPEIDIX IV

•

MEDICAL ASPECTS OF FALL-OUT FROM BRAVO

1. Medical evaluation of personnel exposed to the radiation from fall-out in case of BRAVO depends to a great extent upon the accuracy with which dosage can be computed. Based upon extrapolation of fall-out time from Rongerik data where the fall-out time was precisely determined by automatic recording instruments, it seems plausible to conclude, after making allowance for factors giving maximum values of time and intensity, that personnel were not exposed to dosages much higher than calculated. This is particularly true inasmuch as Rongerik calculations were in good agreement with observed film badge data on personnel there.

2. The association of symptoms with a given dosage may lead to erromeous conclusions since such tabular relationships have been devised only for whole body penetrating radiation given over a period of a few minutes. It is now generally believed that the symptoms in those tabulations will appear with a smaller dose than indicated. These personnel may develop signs or symptoms out of proportion to what would have previously been expected but could be somewhat tempered by the relatively slow dose rate characteristic of fall-out. a definition of a state of the state of

3. With respect to natives, due to the language difficulty, it was extranely doubtful that information obtained by questioning would be reliable.

4. We may draw certain conclusions, however, which seem to be sound concerning immediate prognosis based upon the doses believed to have been received.

5. Considering the personnal involved in exposure to radiation they can be grouped according to location:

a. Rongerik -

í

Twenty-eight Americans were exposed showing film badge readings ranging from 40 to 98 roentgens during a period of 28.5 to 35 hours. They were evacuated to Kwajalein. It was not expected that any of these men would develop any subjective symptoms. One admitted to feeling badly until reassured after which he admitted that his feeling was probably psychological. First blood counts taken on D plus 1 showed a normal distribution. Generalized loss of hair which usually occurs after 10 days with sufficient dosage was not expected and has not occurred to date. Levels of personnel contamination were not exceedingly high and inassuch as decontamination was performed on D plus 1, beta burns are unlikely.

COLLED FUT:

b. Rongeliap -

Sixty-five natives were evacuated to Kwajalein and may have received doses as high as 130 roentgens in a period of 51 hours. In this case, the level of radiation is about the level which might cause some symptoms such as nausea, vomiting, fatiguability and loss of hair for acute doses. Allowing for the reduced effect from low dose rate it may happen that symptoms as above will occur in individuals who were already ill or in generally poor physical condition. Readings of skin and hair contamination were such that for this exposure time spotty distribution of beta burns could occur within several days. If this occurs, ulcerations might develop which may require several months to heal.

وموارغه ورخابا ومعادية ومعتدية وبالمحمد فمعاليهم وأرجادهم أرجاد والمراد والمعادية والمعادية والمعادية والمعالية

c. Allinginae -

Seventeen natives on this island were exposed to approximately 80 roentgens in 58 hours. They were evacuated to Kwajalein. It was not expected that any subjective systemic symptoms would develop. However, personal contamination of this duration could conceivably cause beta burns in a spotty distribution with ulceration as described above.

d. <u>Otirik</u> -

ł

154 natives were evacuated to Ewajalein after receiving a dose of 17 roentgans in 78 hours. No subjective systemic symptoms or changes in blood count were expected. Beta burns are unlikely but are possible statistically.

e. <u>Alluk</u> with 401 natives was not evacuated and the total dose for a life time will be less than 20 roentgens. No medical problem from radiation should occur in the population.

f. Some other islands received fall-out exposing inhabitants to insignificant quantities of radiation.

8. Task Force personnel at or in the vicinity of Bikini Atol1 -

Personnel in the concrete bunker on NAM island were evacuated to ships afloat receiving in general comparable dosage to those aboard ships all the time. Based on readings taken aboard the ships it was estimated that none of the ship's personnel would receive more than 10 roentgens whole body radiation. This doze would not cause any general symptoms of radiation sickness, however, decontamination personnel might have akin contact with concentrated radicactive deposits and possibly sustain mild beta burns.

6. All native evacuees were held at Kwajalein for observation and treatment should the need arise. The station medical complement took complete blood counts, made physical examinations and took histories. Captain H. H. Haight, (MC), USN, a radiological medical officer was sent to Ewajalein as consultant on radiation effects to the station surgeon. Daily observation was instituted in anticipation of the arrival of a medical group from the U. S. who were to investigate the patients.



7. The medical group arrived in Ewajalein on 8 March. It consisted of military and civilian modical officers and technicians from the Naval Medical Research Institute, the Armed Forces Special Weapons Project and the U. S. Kaval Radiological Defense Laboratory and was established as Project 4.1, TU 13 of Task Group 7.1 with Commander E. P. Cronkite, MC, USH, as Project Officer. Drs. G. V. Leroy and C. L. Dunham represented the Division of Biology and Medicine, AEC, and were to act as advisors to Project 4.1. A systematic organization was set up with a view toward running a sick call, performing blood studies, taking histories, making physical examinations and documenting the cases by means of records and photography. Buildings were furnished for these purposes by COMNAVSTAKKAJ and his Station Surgeon, Commander W. J. Hall, worked closely with the group. The establishment of the investigating group of Project 4.1 was essential and desirable from several standpoints. All the medical personnel were experienced in the field of stomic medicine having been participants in previcus testing using biological material as well as having had full time research projects along this line during interim periods. This allows for proper evaluation of human effects toward correlation with data on animals from which a great deal of our ideas on human effects have been extrapolated. Further, they constituted an angmentation medical group for treatment if necessary in conjunction with station medical facilities. An additional advantage was that almost all of the personnel had worked together as a unit on previous occasions.

TULIVITA

and the second second

and a star of the second second

四百日日 一日日日日

8. Some of the natives nor the Rongerik Americans had preliminary or ently systemic symptoms consistent with radiation sickness from large dosage of external whole body irradiation. A reported case of vomiting and a few cases of loss of appetite were not significant considering the sudden change in environment and dist to which they were subjected. To relieve the load on the station medical facilities, not knowing of the early arrival of the modical group, the twenty eight Americans were returned to Invetok to remain as outpetients under the supervision of the Surgeon. Task Group 7.2. Blood counts were taken at approximately three day intervals. They remained asymptomatic although there began a depression of the white blood cells of mild degree. They were returned to Kwajalein on 17 March. During the early days of March all patients remained free of systeric symptoms attibutable to irradiation but there was a definite decrease in the white bod cell count more marked in the Rongelap group. The blood pictures of the Ailinginae natives and the Rongerik Americans were quite similar which was reasonable considering they were exposed to the same order of magnitude of radiation. The Utirik group showed nothing particular from a medical standpoint and were considered as a virtually normal native population for comparison purposes pending time for obtaining base line data from non irradiated natives.

By the thirteenth and fourteenth day a tendency to epilate had become evident in the Bongelap natives involving mostly children but within a few days it had appeared in adults. The epilation was both patchy and diffuse, confined mostly to the head and particularly in children the scalp assumed a spotty appearance due to depignentation of the skin.

3

ATT THE

. .

At about the same time that epilation appeared in the Hongelap group, small skin lesions became noticeable on the folds of the neck, the forehead, shoulders, and arms. They appeared to be superficial and at first were hyperpignented. As time went on, the lesions, which became blister like, began to peel leaving a whitish depignented area in the center. The skin manifestations continued to appear throughout the month of March, all going through the same cycle and involving most of the netives. The most severe cases occurred on the fest with one exception one aan developed a deep ulcer behind one ear. By this time all of the skin lesions except the ear have virtually healed and it appears that repigmentation is taking place.

TTUEITIN

Similar findings but in a lower percentage and at a later date occurred in the Ailinginae group. One American developed what appeared to be superficial radiation lesions on the back. They were hyperpignented and behaved as the others.

Throughout, there have been no demonstrative systemic symptoms other than an epidemic of colds in the Bongelap group. A few cases of secondary infection from skin lesions and some unexplained high fever in children responded well to penicillin with no sensitization reaction.

The white blood counts reached a minimum during the latter part of March with a late depression in blood platelets becoming apparent. The level of the mean counts being well below normal mean counts. Lowest counts were about 30,000 compared to a normal mean of over 300,000 for the natives. There is a definite upswing in the entire blood picture of both the natives and Americans at the present time.

On about 20 March, several cases of radiation burns were reported aboard both the USS BAIROKO and the USS PHILIP. Examination showed that in almost all cases there were discrete areas around the belt line which corresponded well to some lesions seen on the natives. History indicated that these lesions developed sometime between 3 March and 15 March. All were in the process of healing with desquaration and mild depigmentation and were quite superficial. The whole body dose was less than lOR and there were no other symptoms.

Three M-boat operators from TG 7.3 presented film badges reading from 85 to 95E and were sent to Kwajalein to be observed by the medical team on 16 March. Since that time they have had no symptoms, no skin findings nor blood changes. It is likely some discrepancy in badging or wearing of badges must have taken place as careful examination of the badges by densitometer revealed nothing unusual in the radiation to which they were subjected.

It was decided at the outset to manage all cases in a conservative manner, treating symptoms as they arose, avoiding experimentation with treatment but being ready at any time to perform transfusions either of whole blood or platelets if indicated. Sick call was managed daily

ATUENTING-



where complaints were treated as though radiation had not been present. Skin lesions were kept clean by surgical scap with excellent results leading to a minimum of secondary infection and remarkably prompt healing. It is felt that this conservative regimen gave optimum results and that all patients are recovering satisfactorily.

Detailed reports will be rendered by Project 4.1 on all cases. Detailed statistical analyses will be required to properly evaluate the data derived. Urine samples which have been analyzed in the U. S. will be combined with this study. A detailed study of characteristics of the fallout samples, shielding properties of the measuring instruments, and weather analysis will be necessary before a more exact dose of external whole body radiation can be established. The study of all aspects should lead to a much clearer concept of dose versus effect. The picture of external gamma radiation with a broad spectral band, combined with external beta radiation, and internal hazard makes a very complicated problem in the final report.

As a corollary to immediate treatment of the personnel exposed to the radiation, evaluation of the hazard remaining upon rehabilitation must be investigated. To that end soil and water samples, animals, plants and other comestibles are being investigated with a view of determining if and when the natives may be returned to their home atolls.

All personnel who have been involved in large dose exposures and those whose dose was small but who may have to reside in an active area should be observed over a long period of time. The first year following the tests, re-examination should be at quarterly intervals. This has been discussed with the Director, Division of Biology and Medicine, AEC, who advises that it is the intention of his organization to maintain a periodic observation system.

In summary, natives from adjacent stolls and Americans from the Task Force were exposed to radiation in doses from a few roentgens to approximately 150 roentgens. Some of the more heavily irradiated may be considered to have been borderline from a standpoint of seriousness. All should recover from the effects of the exposure.

CLINTON S.

Colonel, Medical Corps Staff Surgeon

5





TAB A

SULLARY OF TEATHER SITUATION FOR BRAVO SHOT

<u>_</u>____

/ ^{...}

CEOCET

and the second states of the second

į



SULLIVEY OF MEATHER SITUATION FOR ERATO SHOT

- 1. <u>GENERAL</u>. This summery presents the weather situation as eristed prior to, at, and just after BRAVO shot time, for the general area of the Marshall Islands, with specific reference to Bikini and the SHRIMP shot site.
- 2. PRE-SHOT MEATHER.

Ĺ

a. General Synoptic Situation

(1) <u>SURFACE</u> - The surface chart was dominated by a high pressure cell located at approximately 34 degrees north 171 degrees east with a central pressure of approximately 1030 LBS. A trough, oriented northeast - southwest lar just off the coast of Japan. Minimum pressures in low latitudes lay just along the equator. Flow over the Marshall Islands area was the east northeast trade flow, with no marked regions of convergence or divergence. (Chart No. 1). During the preceding several days the high pressure cell had been moving slowly eastward, with a slight increase in the central pressure.

(2) UPPER LEVEL CHARTS. From approximately 25,000 to 55,000 feet the flow pattern was dominated by a clockwise rotation centered near 5 degrees north 175 degrees east. This system was essentially vertical from 30,000 to 50,000 feet (Chart No. 4). The flow at these levels, under the influence of this dominant system was from the west southwest. At 20,000 feet the dominant clockwise rotation lay much further to the east, with a secondary center located south





of BIKINI and west of KINJALEDS, giving westerlies at this level in the BIKINI area. To the east of BIKINI the flow was nore northwesterly (Chart No. 3). At 10,000 feet the entire flow was broken up by a group of minor eddies, both clockwise and counterclockwise, features approaching large scale turbulence in nature and showing quite erratic movements; however, there was a general west wind in the vicinity of BIKINI, with the wind speeds being very light (Chart No. 2).

b. Forecast for Shot-Time

(1) The following forecasts, based on the prognostic chart for shot time (Chart No. 5), were issued at the indicated times for EPAVO shot-time:

- (a) H-48 Hrs: <u>WEATHER</u>: Scattered cumulus, scattered cirrus, widely scattered showers. <u>WEIDS</u>: Surface easterly 10 to 15 knots Ten thousand feet easterly at 15 to 25 kmots 50 thousand feet southwesterly at 10 to 20 kmots 50 thousand feet southwesterly at 10 to 20 kmots 50 thousand feet southerly at 5 to 15 kmots.
- (b) H-38 Hrs: <u>IEATHER</u>: Scattered cumulus bases 2000, scattered cirrus based at 38,000 feet, very midely scattered showers. <u>NEIDS</u>: Surface 80 to 90 degrees at 15 to 20 knots Ten thousand feet 80 to 90 degrees at 15 to 20 knots 20 thousand feet 90 degrees at 10 to 15 knots 30 thousand feet southerly at 5 to 10 knots 40 thousand feet southerly at 5 to 10 knots 40 thousand feet southerly at 20 hnots 50 thousand feet southerly at 20 knots 50 thousand feet to 250 degrees at 20 knots 50 thousand feet to 250 degrees at 20 knots 50 thousand feet southwesterly at 220 to 250 degrees at 20 knots 50 thousand feet southwesterly at 20 to 250 degrees at 20 knots 50 knots





- (c) H-24 Hrs: <u>ITATHER</u>: 3/8 cumulus bases 2000 tops 500, 3/8 cirrus bases near 38,000 widely scattered showers.
 <u>ITEDS</u>: Surface 70 degrees at 20 knots 10 thousand feet at 14 knots 20 thousand feet 230 degrees at 16 knots 30 thousand 350 degrees at 14 knots 35 thousand feet 260 degrees at 22 knots 40 thousand feet 250 degrees at 24 knots 45 thousand feet 240 degrees at 24 knots 55 thousand feet 230 degrees at 10 knots 50 thousand feet 230 degrees at 14 knots 55 thousand feet 10 knots 70 thousand feet 70 degrees at 14 knots 50 thousand feet 90 degrees at 12 knots 90 thousand feet 100 degrees at 18 knots.
- (d) H-13 Hrs: <u>NEATHER</u>: No change in H-24 forecast. <u>NENDS</u>: No change.
- (e) H-8 Hrs: <u>MEATHER</u>: 2/8 cumulus bases 2000 tops 5000, 2/8 stratocumulus bases 6000 tops 7000, 4/8 cirrus bases near 38,000 widely scattered showers, contrail level 36,000 tropopause height 55,000 temperature of tropopause minus 78 degrees. <u>MENDS</u>: Surface 70 degrees at 20 knots - 5 thousand feet 70 degrees at 16 knots - 10 thousand feet light and variable with a westerly trend - 14 thousand feet 250 degrees at 16 knots - 20 thousand feet 270 degrees at 12 knots -25 thousand feet 230 degrees at 20 knots - 30 thousand feet 230 degrees at 25 degrees - thirty five thousand feet 230 degrees at 36 knots - 40 thousand feet 240

DEDILL



degrees at 40 knots - 45 thousand feet 250 degrees at 35 knots - 50 thousand feet 80 degrees at 10 knots - 50 thousand feet lightly and variable - 70 thousand feet 50 degrees at 25 knots - 90 thousand feet 70 degrees at 20 knots.

- (f H-4 Hrs: <u>WEATHER</u>: No change in H-8 forecast. <u>IERS</u>: Se change.
- (2) The air particle trajectory chart (Chart No. 6) was used to brief on the movement of the cloud after detonation for the several levels portrayed.
- 3. <u>SITUATION AT H-HOUR</u>
 - a. General Synoptic Situation

(1) <u>SURFACE</u> - The general synoptic patterns at this time differed from those earlier only by the continued eastward movement of the high pressure cell (Chart No. 7).

(2) <u>UPPER LEVELS</u> - The clockwise system had moved slightly to the east and south, with the winds at upper levels tending to become more westerly (Chart No. 10). At 20,000 feet the small clockwise circulation west of Kwajalein had expanded (Chart No. 9). At 10,000 feet the flow remained dominated by the minor eddies which had existed at 230300Z (Chart No. 8).

b. For the observed local conditions versus the forecast for shot time, see paragraphs 5g, h and i below. A revised air particle trajectory forecast was issued at H/14 hours, based on data available through H/9 hours (Chart No. 11). The major change, which had not been forecast, was in the 20,000 foot trajectory.





4. POST-SHOT WEATHER:

a. The weather after shot time deteriorated rapidly due to induced local cloudiness. The winds zloft for Eniwetok and Rongerik were less favorable from a fallout point of view (Incl No. 1 and 2). b. Chart No. 11 is the computed air particle trajectories for EE470 day, based on data available through H433 hours. Amplification of other levels than those presented is in order since the pattern from the surface to 25,000 feet was in a continual flux. The 25,000 foot trajectory was essentially the same for 30,000 feet. From 12,000 to 18,000 feet the trajectories lie between the 10,000 and 20,000 foot level patterns with the 18,000 foot most like the 20,000 foot trajectory and the 12,000 foot most like the 10,000 foot trajectory. In other words, between 10,000 and 20,000 feet there must have been comsiderable dispersion, with the 20,000 foot level having the most northerly component of motion. From 6,000 to 8,000 feet, the trajectories again start showing northerly components, with the 6,000 foot and below trajectories being generally from the east northeast.

5. DISCUSSION:

a. Briefings:

(1) The outlook on the morning of 27 February was very favorable from the radsafe point of view. The winds for shot time were forecast to be easterly from the surface through 20,000 feet, southeasterly at 30,000 feet, southerly at 40,000 feet, and southwesterly at 50,000 feet. The situation was changing slightly; 27 February would have been an ideal shot day (Incl No. 3).





(2). On the afternoon of 27 February, the weather outlook was still favorable from all considerations, especially radsafe. Easterly winds were forecast for shot time from the surface through 10,000 feet (080° to 090° at 15 to 20 knots), and southwesterly aloft (230° to 250° at 16 to 25 knots) up to the tropopause (55,000 feet) (Incl No. 4).

(3) The briefing at 1100¥, 28 February, indicated a forecast for shot site at shot time as 3/8 cumulus, bases 2,000, tops 5,000;
3/8 cirrus at 39,000 feet and winds to be 070° at 20 knots at the surface (Incl 5).

(4) Later that afternoon, the trend at 10,000 feet became more southerly due to the shift of the high pressure cell at 10,000 feet to the north. The remaining winds were forecast to remain essentially as given at the 1100% briefing (Incl No. 6).

(5) The briefing given at midnight prior to the shot time at 0645, 1 March 1954, was as per the inclosed forecast (Incl No. 7).
(6) The briefing at 0400K, 1 March, was essentially as briefed at midnight. Winds aloft from the CURTISS for 2400M and 0300M were discussed (Incl No. 3). The levels 7,000 feet through 11,000 feet were under close scrutiny due to their variability. A tendency for a westerly direction at that level was forecast (Incl No. 8).

b. Observed shot time winds and weather:

The observed shot time winds (0600%, 1 March 1954) from the USS CURTISS, Eniwetok and Rongerik versus the forecast winds for . shot time for the shot site are shown in Table I (next page).





OBSERTED 06001, 1 LAFOH

	FORECAST FOR BIKINI 06154, 1 Mar	USS CURTISS (AT BIAINI)	EILETOE	1252 S.
Surface	070/20	050/12	070/17	030/17
1000	•	070/17	080/20	070/15
2000		030/18	020/20	050/11
3000		090/17	CSJ/19	050/10
4000		090/14	030/17	090/18
5000	070/16	100/09	C30/13	CEC/15
6000	•	120/04	Calm	C7C/12
7000		310/04	310/09	050/05
8000		310/05	310/11	C20/03
9000		320/07	360/11	330/06
10000	Light & Variable (westerly trend)	310/10	290/11	310,′38
12000	•••	300/07	250/10	290 ′09
14000		290/14	240/07	320/13
1,5000	250/18	·	•	
15000		290/13	265,13	310/15
18000	•	280/13	280/17	290/11
20000	270/12	260/23	280/17	300/19
2 5000	230/20	250/26	242/25	250/24
30000	230/26	240/35	250/28	250/29
35000	240/28	230/35	24.0/37	242/41
10000	230/38	250/44	240/12	21,2/1,8
45000	240/40	250/45	260/23	250/42
50000	260/38	250/31	270/19	250/31
55000	050/10	200/16	300/11	340/06
60000	Light & Variable		330/04	220/03
70000	080/09		080/27	090/13
00008	080/25		080/30	
90000	070/20			

<u>Tiele</u> I

(2) The following was the forecast of clouis and weather for shot time over shot site: 2/3 cumulus, bases 2,000 feet, tops 5,000 feet; 2/8 stratocumulus, bases 6,000 feet, tops 7,000 feet; 4/8 this cirrus at 38,000 feet; widely southered light showers in the area, none over shot site at shot time; contrail formation level 36,000 feet; tropopause height, 55,000 feet with temperature of 72°C.



ſ

Ç

-

ニキッパ



(3) The following conditions were observed at shot time at Bikini (A consensus of shot time observations taken by 10 wether officers and aerographers mates) aboard the USS CURTISS: 4/2 cumulus at 2,000 feet, tops 4,000, 1/5 altostratus (barely discerpable) and 5/8 to 6/8 cirrus, thin.

Sea Level Pressure1006.1 mbs.Temperature800PDew Point72°PSurface Wind074° 15KHeight of Tropopause54,350 ft (at H/3)

A 3-36 over ground zero at approximately H minus 20 minutes reported scattered cumulus, scattered to broken cirrus with bases 38,000 to 39,000 feet and tops to 40,000 feet, and no showers. (4) The 282100M Feb wind observation from Rongerik which was received on board the ESTES was garbled in transmission and was in considerable error. It did not agree with the winds received from Eniwetok and the CUETISS for that time, in that it showed southwesterly flow from 10,000 feet; the correct observation indicated light northwesterly flow, which was in agreement with the other observations.

(5) Local weather conditions began deteriorating shortly after the detonation. Hulticloud layers developed at all levels. The winds aloft were forecast to be less desirable from a ratsafe consideration, and such was the case. Such a situation continued through 8 March 1954 (See Hodographs under TAR E).



c. Analysis of winds aloft:

(1) At 10,000 feet a belt of outdrafts across the Facific, at approximately 25 to 30 degrees north, gave easterly flow generally throughout the northern Marshalls. There was an intense floma storm in the vicinity of Hawaii. A number of weak short-duration circulations characterized by light and variable winds formed and dissipated between the equator and 20 degrees north. One clockwise circulation developed near Eniwetok on 28 February and drifted toward Rusaie, giving weak westerly flow over the shot area. This general situation persisted through EPAVO plus 3 days, with winds at all times less than 11 knots at Eniwetok. Analysis at this level was difficult and very changeable.

(2) At 20,000 feet, the major systems on HEAVO minus 5 days were a large outdraft 20 degrees north, 170 degrees east, and deep middle latitude troughs near 145 degrees east and 150 degrees west. This latter trough remained stationary while the western trough moved eastward, forcing the outdraft to 180 degrees east, thence southward and westward into the Marshalls. This trend resulted in shot site winds veering from east through south to west and continued after HEAVO day. Speeds were under 20 knots until HEAVO plus 1 day, then increased to approximately 30 knots on HEAVO plus 2 days.

(3) At 30,000 feet and 40,000 feet, a belt of clockwise circulations centered near 10 degrees north moved slowly southward with



JEUILE

individual cells drifting eastward. Movement of the cell near Eniwetok to a position southeast of Majuro gave strong southwesterly flow in the local area on 28 February and west southwest flow on 1 March, replacing the light variable winds which existed on 23 - 25 February. Continued southward notion and approach of following cell from west brought west to northwest flow on 2 and 3 March.

(4) At 50,000 feet, the slow movement of the clockwise circulation from a position northwest of Eniwetok to south of Majuro changed the flow gradually from north northwest to west southwest. After 1 March this circulation continued its movement to the southeast and weakened, with flow in the local area becoming west northwesterly.

(5) At 60,000 feet, no clearly defined trend was present. Whinds were quite light and erratic at this level, with easterly flow at higher levels. These variable winds persisted through ERAVO plus 3 days.

d. Differences between forecast and observed winds: In regard to the differences between forecast and observed winds, reference is made to a report of Project 4.5, JTF THEE, by Palmer, Miller, and Stopinski. This report stated that studies of the observational errors in upper wind observations taken by GED/1 equipment indicated that for wind speeds above 10 knots, approximately 15 per cent of the observations varied by more than 20 degrees, even when the observations were made by several different ground units, but with the same airborne unit. The differences between forecast

PEUHLI


winds and observed winds during the period of ERAVO were within these observational limits. On the last specific wind forecast issued at H minus 8 hours, there were ten (10) winds forecast to be above 10 knots which can be compared with the observed winds taken from the CURTISS at HRAVO hour. Of these ten (10) winds, six (6) differed by 10 degrees, two (2) by 20 degrees, one (1) by 30 degrees and one (1) by 40 degrees. 80 per cent of the forecast winds which can be checked in the immediate locale were within observational limits imposed by the equipment itself. The errors of 30 degrees and 40 degrees were the forecasts for the levels immediately above and below 10,000 feet.

6. <u>CONCLUSIONS</u>:

a. Weather conditions during the five days prior to HRAVO day were indicative of a favorable trend for HRAVO, involving on an average, easterly winds below 15,000 feet, with winds of a southerly component above. HRAVO minus 2 and HRAVO minus 1 days were especially favorable from a fallout point of view. The weather situation presented at H minus 6 hours for a 24 hour period was satisfactory; however, an unfavorable trend was predicted to occur during the following 24 hours since northwest winds were forecast for the 10,000 to 20,000 foot

b. The forecast wind directions were well within the normal forecast error, which must be anticipated. Forecasts of the same precision as those made in areas of dense observation networks cannot be expected in this area. The forecasts of winds aloft for ERAVO were, nevertheless, approaching the limits of human ability which the art at present OFOREE.

allows.

Lt. Colonel, USAF Staff Weather Officer

21 Inclosures

- 1. Area Forecast Valid Oll5001 to C21500M March 1954.
- 2. Rongerik and Eniwetok Observed Winds Aloft for 0121001 March 1954.
- 3. Msg DTG 202016Z Feb 1954, to CUTF SENEN, re: Meather Outlook for Shot Day.
- L. LSg DIG 2705442 Feb 1954, to CJIF SEVEN, re: Weather Outlock for Shot Dar.
- 5. Lemo for Record, B-1, 1100M Weather Briefing w/2 Incl.
- 6. Meno for Record, B-1, 1600M Weather Briefing.
- 7. Memo for Record, 3-1, 2400M Weather Briefing W/2 Incl.
- E. Mero for Record, 0400M ERAVO Day Meether Briefing.
- 9. Symbols for Weather Distribution Chart.
- Chart #1 1500 foot Streamlines and Teather Distribution Chart for 2300002 February 1954.
- 12. Chart #2 10,000 foot Streamlines for 2803002 February 1954.
- 12. Chart #3 20,000 foot Streamlines for 280300Z February 1954.
- 13. Chart #4 40,000 foot Streamlines for 280300Z February 1954.
- 14. Chart #5 Prognostic 1500 foot Streamline and Weather Distribution valid for 05451, 1 March 1954.
- 15. Chart #5 Air Particle Trajectories for H to H plus 72 hours, prepared at H minus 9 hours.
- Chart #7 1500 foot Streamlines and Weather Distribution Chart for 2318002 February 1954.
- 17. Chart #3 10,000 foot Streamlines for 281800Z February 1954.
- 18. Chart #9 20,000 foot Streamlines for 281800Z February 1954.
- 19. Chart #10 40,000 foot Streamlines for 281800Z February 1954.
- 20. Chart fil Revised Air Particle Trajectories prepared at H plus 9 hours.
- 21. Chart #12 Completed Air Particle Trajectories prepared at H plus 36 hours

APEA FORECAST VALID 011500 TO 0215001 MARCH 1954

General Situation: Easterly flow extends to 7000 feet. Testerlies prevail to tropopause. Eniwetok Terminal: 18 sold 180 sold 350 brkm vsby 10 sic wind 0715. <u>Bikini Terminal</u>: 18 brkm 180 brkm 380 brkm vsby 10 sfc wind 0715. Eniwetok winds and temps: 00715 20820 40822 60718 83604 03302 F11C 152511 P2C 202917 17C 302830 126C 402632 150C 502628 603308 700718 800230 900740. Bikini Winds and Terrs: 00715 208204 40822 60718 63604 93508 P110 152612 P2C 202722 17C 302628 126C 402636 150C 502730 60330 700718 800830 900740. Further outlook for Eniwetok and Bikini (Planning Only). Decreasing clouds Bikini area. Little change elsewhere. Briwetok Clouds: 2/8 CU bases 13 tops 40 with sold isolated tops to 60. 4/8 as bases 130 tops 200. 7/8 CI near 380. <u>Bikini Clouds</u>: 6/6 CU bases 180 tops 40 with sold isolated tops to 60. 5/8 as bases 186 tops 200. 7/8 CI near 350. ECHE 3/8 CU bases 18 tops 40 with sold isolated tops to 60. 2/8 as bases 160 tops 200. 5/8 CI near 380 after 21004. Freezing level 170. Tropopause height 550 temp & 820. Height of contrail formation 350. Sea and swell 6 feet from EE Eniwetok, Bikini. <u>Kwajalein Terminal</u>: 15 sotd vsby 10 sfc wind 0510 tempo 10 brkn vsby 5 sfc wind 0615 in 1gt sctd shwrs.

RONGERIK OBSERVED WIEDS ALOFT

1 Earch 1954, 21004

(Last observation made prior to enameridan) 00510 11013 21015 31012 41011 50711 60908 70703 81803 92706 99991 02839 23314 43212 62110 83007 99992 03014 52827 97773 02734 52534 99996 0283 52737 99995 02836 52815

ENIMETOK OBSERVED HINDS ALOFT

1 March 1954, 2100

00715 10914 21015 31013 41010 51010 61104 72805 82806 92810 99991 03829 22706 43305 63410 83315 99992 03030 52728 99993 02525 52526 99994 02534 52532 99995 02818 5 calm 99996 00906 52703 99997 0 calm 50920 99998 00928 50942 99999 00946 50946 70954

Ţ

CTRAINER FROM

10

CJIP SEVEN

. .

.

CJTF SEVEN (ADLIN)

WEATHER OUTLOOK FOR SHOT DAY POLLOES CLE CLOUDS AND MEATHER CLE SCATTERED CUMULUS CAM SCATTERED CLEREDS CAM WHIDELY SCATTERED SHERE FD THOS CLE SID-FACE CAM EASTERLY ONE FIVE TO THO ZERO KTS SACLE ONE ZERO ZERO ZERO ZERO CAM EASTERLY ONE ZERO TO CHE FIVE KTS SACLE THE ZERO ZERO ZERO CAM SOUTH-EASTERLY CHE FIVE TO THO FIVE KTS SACLE FOUR ZERO ZERO ZERO CAM SOUTH-EASTERLY CHE FIVE TO THO FIVE KTS SACLE FOUR ZERO ZERO ZERO CAM SOUTHERLY CHE FIVE TO THO FIVE KTS SACLE FIVE ZERO ZERO ZERO ZERO CAM SOUTHERLY CHE ZERO TO THO FIVE KTS SACLE FIVE ZERO ZERO ZERO ZERO CAM SOUTHERLY CHE ZERO TO THO FIVE KTS SACLE FIVE ZERO ZERO ZERO ZERO CAM SOUTHERLY CHE ZERO TO THO FIVE KTS PD RADSAFE OUTLOCK VEST FAVORABLE

LTCOL C. D. BOIST, USAF

ON CONTIAL

II:CL 3

2705442 FEB 54 OPERATIONAL INVEDIATE

ICLD CNTI

14

COLFIDENTIAL OPERATIONAL INSTEDIATE OM

EO

TO: CJTF SEVEN EXTREMON ATOLL (ADETE)

IEFO: CTG 7.3

FROM: CJTF SEVIE

THATHER OUTLOOK AS OF THO SEVEN ZERO FIVE ZERO ZERO ZERA FOR BRAVO DAY FOLLOIS CLE CLOUDS AND WEATHER CLN SCATTERED CUMULUS GAM SCATTERED CIRCUS CM: VERY MIDELY SCATTERED SHOLERS FD WINDS CLN SURFACE TO FIFTEEN THOUSAND FEET CM: EASTERLY CHI FIVE TO THO ZERO KIS SAMIN FIFTEEN THOUSAND TO THER. IT FIVE THANSAND FEET CM EASTERLY CHE ZERO TO CHE FIVE KIS SAMIN THEN FINS TO EMERT THOUSAND FEET CM: SOUTHERLY FIVE TO CHE ZERO KIS SAMIN THIRTY THOUSAND FEET CM: SOUTHERLY FIVE TO CHE ZERO KIS SAMIN THIRTY THOUSAND FEET CM: SOUTHERLY FIVE TO CHE ZERO KIS SAMIN SUITH THOUSAND FEET COMMENSATERLY CHE ZERO TO CHE FIVE KIS FD RAD-SAFE OUTLOCK FOR SAMINETY AND UNELAND VERY FAVORABLE SAMIN OUTLOOK FOR EIKIET FAVORABLE FD REQUEST YOU TAKE ACTION ON COMPTRIATION OF ELECUTE CHEER PARES ITEM THERE KIGHT CHECK LIST PAREN FD OGLE AND GRAVES AND EXAMPLE OUTLOCK FOR SAMINET CHECK LIST PAREN FD OGLE AND GRAVES AND EXAMPLE PARES ITEM THERE KIGHT CHECK LIST PAREN FD OGLE AND GRAVES AND EXAMPLE PARES INFOLMED CONFIRMATION FD BRUTON INTERFOSES NO OBJECTION

LTCOL C. D. BUILOT, USAF



IIIL 4

28 Tebruary 1954

E DRANDUL FOR RECORD

SUBJECT: B-1,1100K Command Briefing

1. Persons present at the B-1, 1100H Command Briefing were:

Kajor General Clarkson Dr. Graves Brigadier General McGinley Rear Admiral Bruton Mr. Reeves Mr. Gibbons Colonel Cowart Lt. Colonel Harbour Lt. Commander Madden

The briefing was conducted by:

Lt. Colonel Bonnot Captain Maynard, USN Lt. Colonel House

2. The 2700002 February Weather Distribution Chart, with the 2703002 February streamline - superimposed thereon was used for the present situation briefing. The undersigned mentioned the frontal system funnel just below Wake, the deep trough associated with it which was reflected through all levels to 60,000 feet; the present high aloft to the east of Bikini which was also remaining rather stationary. The undersigned further amptioned the easterly flow from the surface to 10,000 feet; the south easterly becoming southerly and south westerly below aloft associated with this deep high east of Bikini. Next presented at the briefing was the forecast for shot time: 3/8 cumulus bases 2,000 feet, top 5,000 feet: 3/8 cirrus 39,000 feet with widely scattered showers. The winds were given as per the attached forecast (Incl 1). Mentioned tropopense as. 55,000 feet also attached is the area forecast valid 280300K February to 010300K March (Incl 2). The undersigned described the air particle trajectory forecast stating that all of the trajectories were to the EE emcept the 10,000 feet one and the 60,000 ones which were westerly.

3. Colonel House at this point continued with the Radsafe portion of the briefing.

Lt Colonel, USAF Staff Weather Officer

2 Incls a. Wind forecast b. Area forecast

SEGNET

II:CL 5

FUERCAST CHARLE 3/8 CU 2000 foot () OI (),000 feet widely scattered showers.

55,000 L SEC C77 5000 C22 10,000 070/11 £0,03**0** 050/10 160/15 15,000 20,000 230/25 70,000 070/14 25,000 2:0/15 30,000 250/14 80,000 090/12 260/22 35,000 90,000 100/18 250/24 40,000 240/20 45,000 50,000 230/14

ICL 54

Ξ,

E.

(

271310Z FEB 54 UNCLASSIFIED OFERATIONAL PRICEITI OL NO ED

FROM: TG 7.4 (AFICAT) TO: HEA CENT ENLISTOR

WEEP ESTES 031 FOR DISSECTIVATION X AFEA FORECAST 280300 TO 0103004 FEB 54 X GENERAL SITUATION I NOT TRADES FERSIST LOVER LEVELS I HIGHER LEVELS DONNI-TED BY STRONG CLOCKLISE CIRCULATION EAST OF MAJURO X RUTHETOKTER/INAL X 20 SCID 390 EEX VSBI 10 MI TELEO 20 ERKN VSBI 6 MI SFC VIND 0918 VITH GUSTS TO 30 KTS IN LGT SCTD SELPS & BIKINI TERPTEAL 20 SCTD 380 BEKN VSEY 10 MT SFC TIND 0920 TELPO 20 BEEN VSBY 3 MI SFC TIND 0920 HITE GUSTS TO 30 KTS IL LOT SOLD SHEES I EXTLETOR JUNDS AND TELPS 00918 20922 40924 60926 81020 01116 P15C 151414 P6C 202410 12C 202320 120C 402230 102C 502424 600414 700612 800711 900811 X FURTHER OUTLOOK FOR ENTWETOK AND BIKINI (FLANNING ONLY) X SLIGHT INCHRASE IN SHOULE ACTIVITY & SOUR DECEPASE IN CIFFUS & ENTWETON CLOUDS 3/8 CU BASES 20 TOPS 40 WITH SCTD ISOLATED TOPS TO 60 I 5/8 CI NEAR 380 I TEAPO 5/8 CU BASES 3 TOPS 60 WITH SCID ISOLATED TOPS TO 70 I 2/8 AC EASES 120 TOPS 140 I 6/8 CI NEAR 380 I BIKINI CLOUDS SAME AS ENTRETOK I SIGNIFICANT CLOUD AREAS I PATCH SC EXISTS OVER EASTERN MAPSHALLS I CIRCUS COVERS ENTIRE MARSHALL AND GILBERT ISLAND AREA X FREEZING LEVEL 175 X TRO-POPLUSE HEIGHT 550 TELP LEAC HEIGHT OF CONTRAIL FORMATION 370 I SEA AND STELL 8 FT FROM ENE AT EXTRETOK AND EIKINI X KWAJALEIN TERMINAL X 15 BEEN VSEY 10 MI TELPO 10 BEKE VSEY 5 MI SFC WIND OB16 WITH GUSTS TO 25 KIS in 100 SHORERS

HERSCHEL H SLATER, LTCOL, USAF

· ``-

SCOR

28 February 1954

MEMORANDUM FOR RECORD

Itacurated in the

۰,

(

(<u>)</u>

SUBJECT: B minus 1, 1800 Hour Weather Briefing

1. The following personnel were present for this briefing: Major General Clarkson, Brig. General McCinley, Dr. Greves, Dr. Ogle, Mr. Reeves, Dr. Duane Sewell, Colonel Cowart, Captain Maynard and Lt. Colonel House.

2. This was a very short briefing wherein mainly the light winds at 10,000 feet were mentioned as the only significant change to the 1100 hour briefing given this morning. It was pointed out that there was a tendency for southerly winds, which were very light, at 10,000 feet. It. Colonel House continued with the radsafe briefing.

BONKOZ

It. Colonel, USAF Staff Teather Office

JEUNE



1 March 1954

LEVORANDUM FOR EECORD

SUBJECT: 2400L Teather Briefing Prior to ERAVO Shot (B-0645 hours)

1. The following personnel were present for this briefing: Lajor General Clarkson, Dr. Graves, Brig. General EcGinley, Brig. General Estes, Dr. Ogle, Dr. Duane Sewell, Mr. Reeves, Capt. Knickerbocker, Col. Cowart, Capt. L'aynard and Lt. Col. House.

2. The weather briefing was opened with a general statement on the synoptic situation over the Pacific area. It was mentioned that there was no significant change from the briefing this morning, and that the weather in general was holding up for shot time.

3. The attached forecast was presented (Incl a) which stated that at shot time there would be 2/8 of cumulus, bases 2,000 feet, tops 5,000 feet; 2/8 of stratocumulus, bases 6,000 feet, tops 7,000 feet; 4/8 thin cirrus at 35,000 feet; widely scattered light showers in the area, but none over shot site at shot time; and contrail formation level at 36,000 feet. The tropopause height was 55,000 feet; temperature -78° C. Hention was made of the 40 knot winds at 40,000 feet, and that the cirrus was caused by the flow around the high, located at 30 to 40 thousand feet over this area.

4. That there might be some locally induced weather caused by the detonation itself was mentioned, as well as the light and variable winds at 10,000 feet; that these winds would likely have a westerly component; and that the level from 7,000 feet through 14,000 feet was also very light and variable, the tendency being for a westerly component at all these levels.

5. The terminal forecast for Kwajalein and Wake was given next, stating that Wake had been quite poor, with low ceilings and frequent showers but would improve for shot time and would have broken cumulus and cirrus with scattered showers in the area. For Kwajalein it was stated that there would be continuous showers activity, and that they had been having broken to cocasional overcast lower clouds with occasional showers. These showers reduned visibility to three miles. The latest winds aloft used in this briefing were the 2100% winds from the USS CURTISS (Incl b). These winds, along with the discussion of the air particle trajectories completed the items discussed in the weather briefing.

6. It. Col. House followed this with the radsafe briefing. It was decided to have an additional look at the latest finds at 0400%.

linn

C. D. BUINDT Lt. Colonel, USAF Staff Teather Officer

2 Incls a. Area forecast b. Wind forecast

OF ODE

IIII 7

 \mathbf{C}

 (\cdot)

FROM: TG 7.4 (AFLOAT)

TEN ASSTETIC

TO: SEA CERT RELIGETOR

C

5

FORECAST FOR 0645M, 1 MARCH 54. FREPARED 2200M, 28 FEB 54.

HURP ESTES 04.0 FOST BASI I 2/8 CU 2000 TOPS 5000 FT 2/8 SC BASE 6000 TOPS 7000 FT 4/8 THIN CIRRIS BASE MEAR 38000 WIDELI SCATTERED LIGHT SHUTERS I CONTRAIL LEVEL 36000 I TROPOPAUSE HEIGHT 55TESD 1780 I WINDS 00720 50716 10147 152518 202712 252320 302326 402338 452440 502638 550510 60 147 700809 800825 900720

H. H. SLATER, LTCOL, USAF

BIKINI OBSERVED WINDS ALOFT TAKEN ABOARD THE USS CURTISS

28 February 1954 - 2100M

00714 10722 20722 30814 40514 50608 60905 71302 82805 93011 99991 03311 20107 43117 62515 82714 99992 02716 52627 99993 01830 52430 99994 02537 52244

A I I

 \bigcirc

 $\boldsymbol{\zeta}$

1 March 1954

VELSEANDUM FOR RECORD

SUBJECT: Teather Briefing at 04001, HRAND Day

1. The following personnel were present for this briefing: General Teyland, Major General Clarkson, Dr. Graves, Brig. General McGinley, Brig. General Estes, Dr. Ogle, Dr. Duane Sewell, Mr. Reeves, Colonel Cowart, Captain Maynard and Lt. Colonel House.

2. The CURTISS winds aloft for midnight and 030000 were discussed in view of the relationship to the shot site, the levels 7,000 fest through 11,000 fest being scrutinized the most (Incl a). The area forecast for shot day is also inclosed (Incl b).

3. It. Colonel House further discussed the radsale situation after which it was decided to go ahead with the shot as scheduled.

D. BONNOT

It. Colonel, USAF Staff Teather Officer

2 Incla

(...

ſ

. .

> a. CURTISS winds aloft for 0000M and 0300M 1 March 54.
> b. Area Forecast for Shot Day



BIKINI OBSERVED WINDS ANOTE MATEN APPLARD THE USS CURELSS

1 Mamin 1.024 - CCCCH

00518 10726 20820 30819 40815 50508 60505 73503 83311 93215 99991 03215 22217 42809 62915 82916 99992 02002 50517 95993 02121 52324 99994 02343 52441 50 13 seing 55 Missing 99995 02736

1 March 205% - 0300M

00921 10818 20719 30718 40815 50510 60707 70504 80205 93410 99991 03312 22514 42715 62415 82917 99992 02728 52224 99993 02231 52238 99994 02238 52538 99995 02631 53211 90221

í

281440Z TEB 54 USEL METTY

FROM: TG 7.4 (AFLOAT)

TO: WEA CENT ENTRETOK

THEP ESTES OLL FOR DISSELENATION X APEA FORECAST VALID 0103001 TO 0203001 MAR SL X GEVERAL SITUATION X SHALLON LAYER OF EASTERLIES EXTERDS TO 5000 FT OVER AREA X VESTERLY WINDS ALOFT PREVAIL & ENDWETCK TERMINAL & 20 ERMN 380 SCTD VSBY 10 MI SPC TIND 0720 TELPO 15 BRKN VSBY 5 MI IN LGT WIDELY SCTD SHIRS X BIKINI TEMMAL X 20 SCID 380 SCID VSBY 10 HI X SFC KIND 0720 TELPO 15 EREN VSBY 5 HI in LGT TIDELY SCID SHIPS I ENTHETCH WINDS AND TELPS I 00720 20723 40816 63012 83012 03212 FLC 152410 P3C, 202314 145C 302325 126C 402335 148C 502638 600413 700512 800725 900718 EIKIKI TIEDS AND TEEPS I 00720 20822 40818 63210 83210 03312 FILC 152518 P3C 202612 115C 302325 126C 402342 148C 502735 6014V 700809 800825 900718 I FURTHER CUTLOOK FOR EXIMETON AND BIKINI (PLANNING CHLY) SLIGHT INCREASE IN LOS CLOUDI-LIESS AND SHOHER ACTIVITY I ENTHETOK CLOUDS 2/8 CU BASES 20 TOPS 40 JITH SCTD ISOLATED TOPS TO 60 2/6 SC BASES 50 TOPS 60 4/8 CI NEAR 380 I BIRINI CLOUDS I 2/8 CU BASES 20 TOPS 40 KITH SCTD ISOLATED TOPS TO 60 2/8 SC BASES 50 TOPS 60 4/8 CINEAR 380 X SIGNIFICANT CLOUD AREAS X BROKEN CIRRUS SOUTH AND EAST SITE-SIVE SHEET OF LIDDLE AND LOX CLOUDS EXTERDING EASTREST THROUGH WARE I FREEZOEG LEVEL 170 X TROPOPAUSE HEIGHT 550 TELP 1780 HEIGHT OF CONTRAIL FORMATION 360 X SEA AND SMELL 6 FT FROM ENE AT ENTRETOK AND BIKINI X KWAJALEDI TERLIDIAL 18 EEKE VSEY 10 MI SPC TELD 0520 TELPO 12 HELL VSEY 5 MI SPC TELD 0522 TITH GUSTS TO 30 KTS IN LGT SCTD SHIPS X

H. H. SLATER, LTCOL, USAF

I:I 8P

Έ















-







•









SECRET

TAB B

RADSAFE NARRATIVE SEQUENCE OF EVENTS

By the morning of B-I day, the wind patterns (forecast and actual) were favorable but the trend of the observed resultant wind patterns was toward an unfavorable or marginal condition. No transient shipping was reported on the B-2 day P2V sweep centered on a significant forcast cloud movement on true bearing of 300° out to 800 miles from CZ. The B-1 day search by P2V out to 375 miles on a forecast significant cloud movement on a true bearing of 330° disclosed no transient shipping except the General Patrick, whose course and speed would take her outside the bazardous area by shot time. At the midnight briefing, the forecast offered a less favorable comition in the lower levels (10-25 thousand feet). Resultant winds at about 20,000 feet were forecast in the direction of Rongelap and Rongerik (Figure 1); however, it was considered that the speeds and altitudes did not warrant a conclusion that significant quantities and levels of debris would be carried out so far. TARE Site was forecast to be well in the fall-out area and NaN Site to be in a fairly high intensity area. Since the B-1 day forecasts gave winds tending significantly toward KNE, a decision was made at the midnight briefing to search on B day ahead of the cloud, i.e., centered on true bearing of 65° out to 600 NM and to warn ships out of the 450 NM minimum radius.

The routine H-18 hour advisory to CINCPACFIT indicated no significant fall-out forecast for populated Marshall Islands, and no safety problems on air or surface routes except surface routes between 275° clockwise to 80° out to a radius of 450 NM with possible significant fall-out in this area. No known shipping was in the forecast fall-out area. The surface radex was forecast for shot time to shot plus six hours to be oriented in a narrow sector to the northeast and a wide sector to the south, with an additional circular radex area around GZ of radius 15 miles (Figures 1 and 2). The sector pointing at Rongelap was considered insignificant due to the low altitudes from which fall-out could occur and due to the very light winds acting on the levels involved.

At the 0430, 1 March briefing, no significant change had been observed in the midnight winds received, however, a radsafe recommendation was made to move the task force ships radially further out from the minimum of 30 NM to a minimum of 50 NM in the SE quadrant. The low level cloud was forecast to overrun the TARE camp and move on to the east with a strong possibility of overrunning NAN. The resultant winds pointing at Rongerik and Rongelap were light and were not forecast to transport significant debris to these atolls.

At 0645 the BRAVO detonation was accomplished without hazard to task force personnel. The bunker firing party reported in safe, but by 0715 the radiation levels were reported rising at the bunker. These levels continued to rise to about 25 r/hr. The firing party was con-

(

SECRET

RESTRICTED DATA

CEONLI

sidered to be in a reasonably safe position since the personnal were able to get into a well protected area deep in the bunker, reading approximately 35 mr/hr.

Ë,

The cloud tracking (by NB-29, Wilson 2) during the morning of B Day indicated no contamination of consequence moving toward Eniwetok or Ujelang at ten thousand feet. The afternoon upwind mission for Wilson 2 was a 30 degree sector out to 500 NM centered on true bearing of 70° from GZ. During the upwind portion of the cloud tracker's mission, readings were from 100 to 500 mr/hr at ten thousand feet. During the morning the B Day P2V sweep in front of the cloud encountered contamination early in his mission. This P2V was replaced by another, which completed the search out to 600 NM.

A report was received about 1500 on B Day that the AEC/NIOO instrument in the hands of the weather detachment on Rongerik had gone off scale. These instruments had a full scale reading of 100 mr/hr. The off-scale report was not viewed with concern since task force ships were experiencing readings (while steaming south) of more than 100 mr/ hr (The BAIROKO going as high as 1.0 r/hr on the flight deck). Considering the distance (133 NM) and a cloud tracker at about 1945M, 1 March, reporting of zero contamination over Rongerik, it was generally believed that Rongerik and the task force ships were caught in a general pattern of finely divided (95% less than 5 micron by cascade impactor) particles over a wide area moving ENE to E. The weather detachment was advised of this condition. Nevertheless, the AEC/MICO Iwejalein Flight ABLE was requested at midnight to be run the following morning. Kwajalein Flight ABLE pattern covers all Marshall Islands north of Kwajalein, and up to Taongi as the northern-most turning point. Aerial readings taken on the flight are extrapolated to the ground.

About 2000 the task force commander was briefed on the overall situation as was known at this time. This included the results of some imitial damage and radsafe survey information taken about noon by helicopter, reports from the sampling aircraft (F-84, B-36 Featherweights and B-36 Control), the first twelve hour cloud tracking mission (Wilson 2) and the first few reports on the H/12 to H/24 hour cloud tracker (Wilson 3). The upwind mission for Wilson 3 was a 30 degree sector out to 500 NV centered on true bearing of 65 degrees from Rongerik, followed by a vectored mission to 17N 163E to base at Eniwetok. The task force commander was advised that fairly heavy contamination had been encountered by Wilson 2 in the sector portion of his flight and that one P2V aircraft had been contaminated in about the same region. The task force commander was advised that the readings taken at ten thousand feet by the cloud trackers were thought to be on the order of magnitude of that encountered by the task force ships. Since the only significent com-



tamination was found in regions which confirmed the forecast cloud trajectories, the requirement for H^{2} 4 through H^{4} 8 hour tracking coverage was cancelled.

As a result of the report from Rongerik, the advisory to CINCPACFLT at 2000 hours B Day included mention of fall-out at Rongerik plus ginor fall-out at Rongelap and other northern Marshall Islands. The fall-out was attributed to an H-12 hour change in the forecast air particle trajectory for the twenty thousand foot level. This trajectory, formerly moving toward the ENE, was reforecast to move toward SE in a circular clockwise path through south to west. The CINCPACFLT advisory included no health hazard problem for surface and air routes, but that fall-out on Bikini-Atoll, as well as damage to structures, would delay reentry several days.

During the trip back to Eniwetok on the night of B Day, the fleet encountered a wide area of finely divided (apparently less than 5 micron) particles which caused top-side intensities as high as 350 mr/hr. Appropriate measures were instituted by the Navy Task Group Commander to the affect that all personnel not essential to open deck duties would remain indoors. Ship's weather doors were closed and the washdown systems operated intermittently to hold down the levels.

Based on the advisory from the weather detachment that their instrument was off-scale, the Air Task Group, on the morning of B/1 day, sent a monitor by amphibian aircraft to check the Rongerik situation. This monitor upon arriving over Rongerik, reported the atoll contaminated and requested permission to start evacuation of personnel. At about 1300 the monitor reported readings on Rongerik of 240 mr/hr at 250 feet and 3.2 r/hr one inch off the ground. The Rongerik evacuation was completed in two segments, the first group (8 people) coming out by H/30 hours and the second group (20 people) by H/35 hours.

The special AEC/NYOO Kwajalein Flight ABLE, requested the previous night, had been instructed to make an in-flight report upon reaching Taongi. (A preliminary Flight ABLE report indicated 1350 mr/hr on the ground at Rongelap, 445 mr/hr on the ground at Ailinginae and zero for Wotho.) Accordingly, it was decided to start a destroyer on the way to Rongelap isrediately and to set up a Sk-16 amphibian with monitors to check the surface conditions at Rongelap before dark. The destroyer was directed to be off Rongelap ready to start evacuation at dawn the following day. Trust Territory representative with interpreter was requested to move by FBY from Kwajalein to arrive at Rongelap at the same time. The SA-16 was set up, two responsible monitors were especially briefed to make readings at waist height, use several meters of the same type for comparison and to use different types for cross-check. An average reading of 1.4 r/m made in the living area of Rongelap Island by these monitors was used in the decision the same night to order the destroyer to commence evacuation operations at dawn. Evacuation operations began about 0730, 3 March and

-3-

and were completed by 1030 the same date. Interrogation of natives disclosed that all were present except 17 who were fishing at Ailinginae. Following the Rongelap operation, the destroyer proceeded to Ailinginae, removed the remaining 17 and proceeded to Kwajalein. A total of 17 males, 20 females, 15 boys and 14 girls were removed by destroyer and disenbarked at Ewajalein. 16 old and sick were moved at about 0930 by FBM to Kwajalein. Decontamination of all natives was accomplished during the trip to Ewajalein.

The full report from ABC/NIOO Flight ABLE indicated Utirik ground contamination at 240 mm/hr at 1651M, 2 Kerch and 76 mm/hr at about 1716M. 2 Earch at Ailuk, the nearest populated island to the south. Bikar, the nearest island to the north was determined to be unpopulated and contarinsted to about 750 mr/hr at about 1600M, 2 March. Taongi, the next nearest island to north was 1.5 mr/hr and unpopulated. Based on these facts a decision was made to start another destroyer to Utirik to anticipate an order to start evacuation at dawn on 4 March. In the meantime a PEN was set up to ground survey Utirik on 3 March while the destroyer was on the way. The infinity dose of the Utirik natives was computed at 58 r. The decision to evacuate was made and the destroyer ordered to start evacuation the following morning, 4 March. A total of 47 males, 55 females, 26 boys and 26 girls were removed, decontaminated on the destroyer enroute to Kwajalein and disembarked on 5 March. Questioning of natives disclosed that all had been removed. The destroyers which evacuated Rongelap and Utirik were directed to obtain drinking water samples from these atolls. A check of the water samples indicated from 2 to 28 times the task force standard for full time usage.

With the decision to evacuate Utirik made and the machinery set in motion to accomplish this operation, the status of Ailuk was put up for consideration. This stoll has a population of 401. The infinity dose was determined at less than 20 r, i.e., less than the minimum standard used by the task force for its sampling aircraft crews. This was the major factor in the decision not to evacuate Ailuk.

During the afternoon of 2 March a directive was issued to execute Kwajalein NYOO Flights BAKER and CHARLIE. These flights cover all Marshall Islands south of Kwajalein. The flights were set up on the assumption that the twenty thousand foot trajectory could have brought contamination around to the south and west and contaminated some of the southern Marshalls. The flights were executed on 3 March. No significent ground contamination was found. An additional NYOO type flight was performed over the Gilbert Islands for the same reasons. No significent contamination was found.

On the basis of Flights ABLE, BAKER and CHARLIE, it was determined that no further atolls would need to be evacuated. The affort was there-





fore concentrated on those populated atolls indicating more than 10 mr/hr at H plus 24 hours and which were not evacuated. For this purpose a special survey was set up under the technical direction of Dr. Thomas N. White, H Div, LASL, assisted by Major Robert Grea, Hq JTP SEVEN, to start from Kwajalein on 5 March by FBM.

Following the survey under Dr. White and Major Crea, the next effort was directed toward acquiring data on the evacuated atolls in order that the effects of the radiation could be better evaluated. The investigation included ground monitoring and the taking of soil and water samples from living areas. Secondary purposes were efforts to reduce the adverse impact on real and personal property of the hasty departure, to determine radiation data of scientific interest and to evaluate the time of reoccupancy by the former inhabitants. This effort was assigned to a destroyer in order that working parties would have a floating base for operations ashore and decontamination facilities afloat. The technical direction of the effort was placed under the supervision of Dr. Eerbert Scoville, Technical Director, Armed Forces Special Weapons Project, assisted by representatives of CASTLE Project 2.5a. The rehabilitation portion of the effort was placed under the supervision of the commanding officer of the destroyer. The party with equipment departed 7 March for Kwajalein to join with the Trust Territory representative in a FBN rendezvous with the destroyer at Rongelap early morning of 8 March.

Arrangements were made to air ship soil and water samples to Health and Safety Laboratory, AEC, New York Operations Office, Attention Mr. Merril Risenbud. Mr. Risenbud was requested to provide the task force with decay information and activity per unit area on the soil samples and activity per unit volume on the water samples. He was also requested to make such other analysis as he thought necessary considering the unusual circumstances and interest in ERAVO Event.

Detailed reports by Dr. White, Dr. Scoville and Major Crea have been distributed separately to interested agencies. Continuing surveys of the evacuated atolls have been made for picking up of animals for medical studies, rehabilitation and for studies of marine life. Reports on these activities will be included in the above distribution as they become available.

RICEARD A. HOUSE

Lt Colonel, USAF

Chief, Tech Ops Branch, J-3

6 Incl

- 1. RadSafe factors Considered at the Wea RadSafe Command Briefing.
- 2. Memo for Record: with 6 Incl covering RadSafe Briefing material as presented at Cmd Briefings for BRAVO.
- 3. Discussion of Off-Site Fall-out.
- . 4. Cloud Tracking Operations.
 - 5. MR: Protection of Transient Shipping During Operation CASTLE.
 - 6. Analysis of Fall-out Following BRAVO Event with 4 Incls -5-



RADSLIFE FACTORS CONSIDERED AT THE WEATHER/RADSAFE COLMAND BRIEFINGS

1. Resultant wind diagram, forecast winds for HOW Hour.

2. Surface <u>RADEX</u>: H to H plus 6 hours, bearings, radius, hot areas, cool areas.

3. 72 hour cloud trajectories given by WI Officer to advise British and CINCPACFLT. Sampling area and BU sampling area; need for penetration authority.

4. Air RADEX: Not used at briefing unless requested. Air RADEX plotted and displayed in RadSafe Office. Does not basically affect decision. Sampling region given by hodograph or 72 hour trajectories.

5. Outlooks: (Based on).

- a. Bikini (hodograph)
- b. Eniwetok (hodograph)
- c. Ujelang (hodograph)

d. Netive atolls in SE quad (hodograph)

e. Control destroyer (hodograph)

f. ATF for YAGs (hodograph)

E. Air Routes

- (1) Thru Wake (72 hour trajectory)
- (2) Thru Kwajalein (72 hour trajectory)

h. Surface routes inside 500 miles (approx 1 day cloud travel). Plot of transient ship chart at briefing (hodograph)

i. CINCPAC advisory (72 hour trajectory), native outlcok; fir and surface routes.

ALL BOVE INCLUDED IN CLOUD TRACEING PLAN

6. Summary: Evaluation as (very favorable) (Favorable) (Favorable except) (Unfavorable) RadSafe condition for shot time.

NT NNT I


5 March 1954

MEMORANDUM FOR RECORD:

SUBJECT: RadSafe Material Presented at Weather/RadSafe Command Briefings for BRAVO

Attached hereto are copies of material presented to CJTF SEVEN during the period H minus 48 hours through H minus 2 hours for BRAVO Event.

R. A. HOUSE Lt Col., USAF RadSafe Officer

6 Incls:

 $\langle \cdot \rangle$

- 1. L/R: E-48 hrs TWX Briefing for CJTF SEVEN 2. M/R: E-36 hrs TWX
- 2. M/R: E-30 hrs TWI Briefing for CJTF SEVEN
- 3. K/R: Command Briefing, 1100X, 28 Feb. 1954.
- K/R: Command Briefing, 1800K, 28 F.b. 1954.
 M/R: Command Briefing,
- 5. M/R: Command Briefing, 0000M, 1 March 1954.
- 6. K/R: Final Weather and RadSafe Check, 0430M, 1 March 1954.

CONFIDENTIA

1 March 1954

OP.

I

MEMORANDUM FOR RECORD:

SUBJECT: H-48 Hour TWX Briefing for CJTF SEVEN

CJTF SEVEN

262016Z FEB 54 0

PRIORITY

I

I

CJIF SEVEN (ADAIN)

Weather outlook for shot day follows: Clouds and weather: Scattered cumulus, scattered cirrus, widely scattered showers. Winds: Surface, easterly 15 to 20 knots; 10,000, easterly 10 to 15 knots; 20,000, easterly 15 to 20 knots; 30,000, southeasterly 15 to 25 knots; 40,000, southerly 15 to 25 knots; 50,000, southwesterly 10 to 20 knots, 60,000, southeasterly 5 to 15 knots. RadSafe outlook very favorable

Lt COL C.D. Bonnot

1

"t Col Bonnot

M/R: Compiled by Bonnot, House and Maynard.

CERTIFIED TRUE COPY: /s/ R.A. Eouse Satour /T# R.A. HOUSE Lt Col. USAF



1 March 1954

MELORANDUM FOR RECORD:

SUBJECT: H-36 Hour TWX Briefing for CJTF SEVEN

CJTF SEVEN

270544Z FEB 54 OPERATIONAL INSEDIATE

F

UPTRATIONAL DOCDIATE

10

OY

CJTF SEVEN ENTRETOK ATOLL (ADAIN)

CTG 7.3

Weather outlook as of 270500Z for Bravo day follows: Clouds and weather: Scattered cumulus, scattered cirrus, very widely scattered showers. Winds: Surface to 15,000 feet, easterly 15 to 20 knots; 15,000 to 25,000 feet, easterly 10 to 15 knots; 25,000 to 30,000 feet, southerly 5 to 10 knots; 30,000 to 50,000 feet, southwesterly 20 knots; 60,000 feet nortneesterly 10 to 15 knots. Radsafe outlook for Eniwetok and Ujelang very favorable; outlook for Bikini favorable. Request you take action on confirmation of execute order (Item 38 Checklist). Ogle and Graves and Reeves recommend confirmation. Bruton interposes no objection.

1

LT COL C D BONNOT

LTCOL BONNOT

CERTIFIED TRUE COPY:

Work R.A. House Lt Col., USAF

L/R: Lembers present: Graves, Ogle, Reeves, Bonnot, House, Marmarc, Dr. Graves called Adm Pruton. RAH

T 4.

CONTRA