401933

BEST COPY AVAILABLE

Health Survey of the Trust Territory of the Pacific Islands

ALICE M. HETZEL, B.S.

IN THE SUMMER of 1948 the U.S.S. Whidbey began the voyage from island to island that, before its completion, was to take it through the Caroline Islands, the Marshall Islands, and the Northern Mariana Islands. Under the auspices of the United States Navy, a survey staff aboard the Whidbey was undertaking a survey of health and sanitary conditions. The ultimate goal of the survey was a physical examination of every inhabitant.

These islands, with the exception of Guam in the Marianas, then designated as the Trust Territory of the Pacific Islands (fig. 1), were the former Japanese mandated islands. Invaded and occupied by American military forces during World War II, responsibility for civil administration of the islands was placed with the Navy, pending the enactment of legislation by Congress designating the permanent governing authority. Under direction of the Secretary of the Navy, the Commander in Chief of the Pacific Fleet served as High Commissioner of the trust territory.

The trusteeship agreement of the United Nations directed the United States to care for and improve the health of the inhabitants. It was recognized immediately that a broad general health-service policy must be established to meet the need for improvement of the health and hygiene in the islands. As a result of the war the natives were in a state of emotional shock. They had been displaced from their homes, their food had been confiscated, trade and industry had been abandoned, and schools were nonexistent. Medical needs had long been neglected. The islanders, a primitive people who had been exposed to the diseases of civilization against which they had no racial immunity, were unable to support doctors, dentists, and nurses in private practice or to maintain hospitals, dispensaries, or departments of public health.¹

From Medical Statistics Division, Bureau of Medicine and Surgery, Department of the Navy, Washington, D.C.

OCTOBER 1959



One of the main difficulties encountered in developing a publichealth program for the trust territory was the lack of reliable information concerning health and sanitation conditions. The most reliable reports came from the naval dispensaries where natives living in the immediate vicinity were given medical care and treatment. There was definite indication from these reports that certain diseases were affecting the health of the islanders and that sanitation in general was on a very low level. Without a comprehensive survey, however, a correct appraisal of disease prevalence, morbidity, and sanitation problems could not be made.



Figure 1. The Trust Territory of the Pacific Islands.

Motivated by these considerations, the Navy undertook a health survey of the trust territory to determine the public-health problems of the indigenous population. The U.S.S. Whidbey, converted especially for this survey, was a twin-screw diesel-driven 177-foot vessel displacing 935 tons. The Whidbey was equipped with a photofluorographic unit and facilities for complete physical and dental examinations. The survey was begun in the Marshall Islands area on 1 August 1945. Four officers and 21 enlisted men made up the ship's company. The survey staff was comprised of the following personnel: 2 Medical Corps officers, 1 Dental Corps and 1 Medical Service Corps officer, and 9 hospital corpsmen serving in laboratory, x-ray, dental, epidemiologic, and clerical activities.

1200

VOLUME 10, NO. 10

encompletion and all the second states and the second of the second second second second second second second s

Ł

٢

Traveling from island to island the survey group examined every native present and investigated and evaluated the food, water, sewage, and garbage facilities, as well as the general living conditions and native habits and customs that had a bearing on the subject of health and sanitation. Among other things the survey included a physical examination, photofluorographic examination of the chest, Kahn test, tuberculin skin test, and stool examination. This report concerns those islands surveyed by the U.S.S. Whidbey during the period from August 1948 through June 1950. Three of the trust territory civil administrative districts, those of Palau, Saipan, and the Marshall Islands, are represented. All of the western Caroline Islands, the northern Mariana Islands (except Alamagan and Agrihan Islands), and the eastern islands of the Marshall Islands group are included.

Subsequent to June 1950 the U.S.S. Whidbey also surveyed the Truk and Ponape districts. However, data for these districts corresponding to those for the three districts mentioned before are relatively incomplete. As a consequence, it is impossible to present a comprehensive description of the health status of the population in the trust territory as a whole. The fragmentary data available for Truk and Ponape, while not included in this article, are contained in a detailed study in the Bishop Museum, Honolulu, Hawaii, and the Bureau of Medicine and Surgery, Department of the Navy.

POPULATION CHARACTERISTICS

The total native population of the trust territory was placed at approximately 54,300 in 1950. About 28,600 of these were in the three districts of Palau, Saipan, and the Marshall Islands. The population figures presented in the accompanying tables, totaling 22,146 for these three districts, are based upon the number of identification cards distributed during the survey. One card was distributed to each inhabitant present on the island at the time of the visit of the U.S.S. Whidbey. Although it was intended that every inhabitant be included, the length of time involved in making a survey of this nature, coupled with the fact that there was a great deal of interisland visiting, resulted in some individuals being examined at places away from their home island and others not receiving any examination at all. The area of the island, the scattering of the population, the terrain, and the amount of control exercised over their people by the local chiefs or administrative heads were all factors affecting the extent to which the population was brought into the survey.

Density. Population density varied from island to island. Out of 39 islands or atolls, 5 had a population of less than 100, 24 had a

OCTOBER 1959

1201

population between 100 and 499, and 10 had a population of 500 or more. In this latter group were Saipan, Babelthuap, Yap, Koror, Majuro, and Kwajalein, all with over 1,000 inhabitants.

Area was no indication of population. In terms of size, some of the larger islands were very sparsely populated. Tinian and Rota, for instance, while nearly as large as Saipan, had much smaller populations. On some of the tiny islands with an area of less than 1 square mile, the population density was far greater than on the larger islands. Population density in terms of over-all area was misleading in numerous instances for, on many of the larger islands, only a very small area was inhabitable. This was especially true on the volcanic islands where much of the island was too rocky to be inhabited. The result was a concentration of population in a very small area with an extremely high population density, while the rest of the island had few or no inhabitants at all.²

Table 1 shows by district the number of females visibly pregnant at the time of the health survey. Estimated annual birth rates based on these figures are also shown. For comparative purposes, in the United States in 1949 birth rates per 1,000 estimated female population, between the ages of 15 to 44, were as follows: all races, 105.2; white, 102.6; and nonwhite, 126.4.³

Age distribution. The median age for the three districts combined was 23.0. Over one third of the population was under 15 years of

Population examined	Total	Saipan district	Palou district	Marsh Island distric
		Nuo	ab er	
Total	22, 146	4, 999	10, 575	6, 572
Females 15 to 44 years of age Females pregnant on day of	4, 138	1, 036	2, 416	. 666
strvey	406	119	152	135
		nnnai birth	rate per 1,000	ŋ•
Total	24. 4	31.7	19. 2	27.3
Females 13 to 44 years of age	130.8	153.2	83, 9	262.4

 Table 1. Females 15 to 41 years of age risibly pregnant on day of surrey

"Estimated on basis that all pregnancies will terminate in live births. Rates computed by relating total pregnancies on day of survey to population and multiplying by 4/3.

1202

5013254



VOLUME 10, NO. 10

というないである ちょうちょうちょうちょう ちょうしゅちんしょう

.

「「「「」」

age, and approximately one fifth was 45 years of age or over. It should be borne in mind that most ages were estimated by an interpreter, as very few natives knew their own age.

The age distribution in the Saipan district presented the most youthful pattern of population found in the territory. The median age was 17.3 as compared with 25.5 for the Palau district and 23.6 for the Marshall Islands district. Only part of the Marshall Islands district is included in this study. Statistics were available for about two thirds of the inhabited islands of the district, accounting for over 70 percent of the population.

Age distributions for the islands or atolls of the Marshall Islands and Saipan districts were relatively consistent. In the Palau district, however, there was considerable variation. Merir, Sonsorol, and Tobi had the highest proportion of old people. There was a strong tendency on the part of the young people of these islands to leave for homes elsewhere while the older people remained. Of Merir's 11 inhabitants, only 3 were under 45 years of age. Approximately 70 percent of Sonsorol's population of 111 was 45 years of age and over, and almost one third of the total had reached or passed the 65-year mark. On Tobi, 60 percent of the 128 inhabitants had reached or passed 45 years of age. At the other extreme were Koror, Eauripik, and Rota, each of which has only 10 percent in the 45-year-and-over age group.

The median age of 22.8 years for Tinian Island appeared high as compared with Saipan and Rota in the same district. There were two groups of inhabitants on Tinian, the Yap-Chamorros and the patients at the leprosarium. The latter group was composed mainly of middle-aged and old people. There were no small children at the colony. Infants born at the leprosarium were sent to Saipan immediately and cared for at the hospital until they were adopted. The median age for the leprosarium patient was 35.4 years, while that for the Yap-Chamorros was 16.6 years, the second lowest in these districts.

An unusual situation existed on Pulo Anna. Fourteen of the sixteen inhabitants were members of one family: the chief; his mother; two brothers; two sisters: two young Indonesian girls, the wives of his two brothers; and six children.

Marital status. The traditional marriage customs with their large gatherings, feasts, and exchanges of property were still practiced throughout the territory. Missionary influence had made its imprint on marriage customs and many marriages were being performed with church sanction.

OCTOBER 1959

5013255

Earlier marriages for women than for men resulted in a much higher proportion of married females than males in the 15- to 19-year age group, 30.2 percent for females as compared with 4.7 percent for males. In all age groups up to 45 years of age, with the exception of Palau's 35- to 44-year age group, there were proportionately more married females than males. The reverse was true without exception for those $\frac{15}{2}$ years of age and over.

Height and weight. The inhabitants of the trust territory were of medium stature and weight. In the adult age groups the average height ranged from 5 feet 2 inches to 5 feet 6 inches for males and from 4 feet 11 inches to 5 feet 1 inch for females. The average weight for males was from 133 to 145 pounds and for females from 112 to 135 pounds.

Height and weight varied by district. For a given height the inhabitants of Saipan and Palau were heavier than those in the Marshall Islands, with Saipan having a slight edge over Palau in this respect (table 2). The difference was more marked for females than for males. It may be noted in table 3 that the inhabitants of Saipan were slightly taller than those of the other two districts. In Palau males were noticeably shorter than in Saipan and the Marshall Islands.

Blood-pressure readings. Blood-pressure readings were taken for over 60 percent of the population. Only one reading was taken for each individual and all readings were taken routinely during the course of the examination.

Table 2. Average weight, by height and sex, of 44-year age group

Height (inches) and sex	Balpan district	Palso district	Marshall Islands district				
-	Average weight (pounds)						
6 0 to 62							
Male	130.6	128.5	122.4				
Female	134.4	176.2	123.5				
🖼 to 65							
Male	137.9	137.7	135.3				
Female	149.3	135,9	133.1				
66 to 68			ł				
Male	150.0	147.6	142.9				
Female	() ()	(7)	0				
🗰 to 71]				
Male	164.4	159.2	160.9				
Female	(*)	(C)	ര				

"No lemales over 65 inches in height.

1204

5013256



VOLUME 10, NO. 10

いっていい しているいちょう とうちょう しょうしょう ちょうちょう

a standard offer to see

Age group (years)	Salpan district		Palau	district	Marshall Islands district		
	Male	Female	Male	Female	Male	Female	
10 to 14	53.4	ક્ર 7	52. 4	\$3.3	54. 5	કાર	
15 to 19	63.4	61.1	61. 5	50 5	G 1	j 59. L	
20 to 24	65.8	61.2	64. 6	602 İ	65.4	59.8	
25 to 34	65.4	60.9	6-L 7	60.1	63.8	59,6	
33 10 44	65.6	60.8	64.2	59.0	65, 1	60.0	
45 to 64	64.8	60.4	63.6	59.2	64.6	59.8	
55 and over	63.1	59.1	62. 2	57.9	63. 5	58.7	

Table 3. Average height by age group and sex

No generalizations may be made for the trust territory as a whole with respect to variations in median blood-pressure readings. Variations followed no consistent pattern in the districts. As normally expected, median blood-pressure readings increased with age. This increase was most rapid in the Saipan district. The median systolic and diastolic readings were generally higher for males than for females, with the smallest difference between sexes occurring in the Marshall Islands (table 4).

Systolic	Distolic	Systolic	Disstolic	Systolic	Diastolle
	-		Contraction of the local division of the loc		
119	60	124	76	135	80
113	· 70	115	73	132	79
118	70	122	74	125	75
117	60	119	70	122	73
	[]				
116	76	118	Π	125	78
113	74	117	76	. 130	70
	110 113 118 117 116 113	119 69 113 70 118 70 117 69 116 76 113 74	119 60 124 113 70 115 118 70 122 117 69 119 116 76 118 113 74 117	119 69 124 76 113 70 115 73 118 70 122 74 117 69 119 70 116 76 118 77 113 74 117 76	119 60 124 76 133 113 70 115 73 132 118 70 122 74 125 117 60 119 70 122 116 76 118 77 125 113 74 117 76 130

Table 4. Median systolic and diastolic blood pressure by age group and sex

DISEASE PREVALENCE

American occupation forces found the health of the islanders generally poor and their needs for medical and surgical care extremely pressing. The conditions under which the native population had been forced to live during World War II, wartime diets of low calorie content, and the inadequacy of medical treatment received prior to and during the war had all left their mark.

OCTOBER 1959

1205

By the time the health survey was launched, however, health conditions were much improved. The islanders had received medical care on an emergency basis immediately following American occupation in 1945, until July 1947 when the territory passed from military government to civil administration. In 1947 the health service policy of the Navy for the Trust Territory of the Pacific Islands, outlining a health program for the trust territory, was adopted for the purpose of controlling preventable disease and rendering medical and dental care. By 1 August 1948, the date the health survey started, this program was in full operation and rapid strides had been made in the control of preventable disease. Measures to correct sanitary practices of the islanders were being enforced; treatments, inoculations, and vaccinations had been administered; and the food supply had been improved to the extent that malnutrition no longer existed except in a few areas. Diseases that were previously the most serious threats had been brought under control. In many respects the health survey served as a means of evaluating the health program.

A marked reduction in the prevalence of intestinal parasites was accomplished in the interval prior to the health survey. It is believed that at the time of the American occupation nearly 100 percent of the indigenous population was infested with hookworm. Ascaris and Trichuris were also extremely prevalent. As a result of a mass deworming program, the control of flies, and education regarding health habits, sanitation and particularly the use of sanitary latrines, the prevalence of intestinal parasites had been noticeably reduced throughout the trust territory.

Yaws is another disease that was brought under control with effective treatment. The large number of inhabitants exhibiting multiple scars and the high proportion of positive Kahn tests bear testimony to the widespread prevalence of yaws. Very few primary lesions were noted during this survey, however, as these had been virtually eradicated by extensive treatment with penicillin administered by Navy medical officers on field trips prior to the survey.

The findings of the health survey revealed, however, that certain sanitation problems still remained. Most islands had community latrines that, in general, were clean and in good repair. It was evident, however, that use of the beaches was still a common practice throughout the islands. In many places, leaves and coconut fiber used in lieu of toilet paper were thrown about promiscuously. Maloelap Atoll was the only place where no latrines were available but many islands had clean and well-constructed latrines that were seldom used. Even on Saipan Island, where virtually every household had its own

1206

VOLUME 10, NO. 10

_

the second second

ł

:]

latrine, use of the beach was evident. Garbage was usually buried in pits but on some islands it was burned or carried out to sea and dumped. The most inadequate methods of 'garbage disposal were noted on Utirik Atoll and Namorik Atoll where the garbage was allowed to accumulate over long periods of time.

In general, the fresh-water supply for the islands was furnished by rainfall. This was supplemented by wells wherever possible and, on the high islands, springs, streams, ponds, and reservoirs were also utilized. Rainwater was collected in old oil drums by means of spouts and drains on houses and coconut trees. The water was then stored in the drums, often without covers or other safeguards against debris. There were indications that the water supply on the following islands or atolls was contaminated: Babelthuap, Elato, Lamotrek, Yap, Tabal, Ebon, Kwajalein, Majuro, Mili, Maloelap, Namorik, and Ormed. Most wells were contaminated and the natives were warned to boil water before using it. On Saipan Island all water was chlorinated.

FINDINGS OF HEALTH SURVEY

Stool examinations. Stool examinations were made for 2,252 inhabitants of the Saipan district, 4,584 of the Palau district, and 3,495 of the Marshall Islands district, approximately 45 percent of the combined population. As shown in table 5, 52.4 percent of all stools examined were positive for intestinal parasites. One out of every five positive stools showed evidence of multiple species. The highest proportion of positives occurred among children from 5 to 14 years of age.

Comparison of the three districts reveals that Saipan had the highest proportion of positive examinations, 79.3 percent as com-

Table 5.	Percentage of stools positive for intestinal
	parasites by age group and sex

	Percentage positive				
Age group (years)	Total	Saipan district	i Palau district	Marshall Islands district	
All ages	52.4	79.3	64. 5	19.3	
Male	53, 5	79.5	64. 9	18.9	
Female	51, 4	<u>ເ</u> ສ.າ	64. 2	19.5	
Under 5	47.7	74.6	51.6	20.9	
5 to 14	60.3	88.7	68.8	23.6	
15 10 24	48.8	76.4	65.3	16.1	
25 16 44	51.4	77.2	64.9	16.0	
45 and over	51. 5	74.4	65.7	21.4	

OCTOBER 1959

5013259

pared with 64.5 percent for Palau and a much lower percentage of 19.3 for the Marshall Islands. On the following islands or atolls less than 5 percent of the stool examinations were positive: Ngulu, Lib, Aur, Mejit, Likiep, Utirik, Tabal, and Kwajalein. Over 75 percent of the stool examinations were positive on Eauripik. Namorik, Saipan, Elato. Tinian, Kayangel, Faraulep, Woleai, Satawal, Ifalik, and Lamotrek (table 6).

Table 6.	Distribution of islands by percent of stools
	positive for intestinal parasites

Percent of positive stools	Total number of islands	Saipan district	Palsu district	Marshall Islands district
Under 5	8		2	6
što 9	4 }		-	4
10 to 24	4	-	1	3
25 to 49	5	1	4	- (
50 to 74	7	-	5	2
75 and over	11	2	8	1

The most prevalent intestinal parasites were hookworm, Trichuris or whipworm, and Ascaris or roundworm. Hookworm was found in 23.1 percent of the stools, Trichuris in 22.3 percent, and Ascaris in 17.2 percent. The relative importance of the three species varied from district to district. In the Saipan district Ascaris had the greatest prevalence, with Trichuris ranking second and hookworm third. The rank order in the Palau district was just the reverse, while in the Marshall Islands district Trichuris was the most prevalent, hookworm ranked second, and Ascaris was practically nonexistent (table 7).

Table	7.	Percentage	of	positire	stools	bŗ	parasitic
		in	(e:	station			

	Percent positive					
Parasitic Infestation	Tota!	Baipun distrut	Palau district	Marshail Islands district		
All parasites	32.4	79L J	64. 5	19.3 -		
Hookworm	23.1	26.0	34, 0	6.8		
Trichuris	22.3	40.6	22.0	11.0		
Ascaris	17.2	44.2	16.8	:3		
Other	2.8	16	1.4	29		

VOLUME 10, NO. 10

-

1208

5013260

ł

Variations of prevalence rates by age and sex were slightly different for each species. Hookworm and Ascaris show opposite tendencies with respect to age. The former increased with age while the latter decreased. Trichuris showed little change by age. Hookworm was more prevalent among males, while Trichuris and Ascaris had slightly higher rates among females.

The actual procedure of obtaining the sample of stool specimens resulted in a haphazard rather than a representative cross section of the population. Each individual examined was given a stool box and requested to submit a fecal specimen. The number of specimens obtained varied greatly. After the specimens were received, some had to be rejected because of contamination from the specimen wrappings and others because of inadequacy of the specimen. The remaining specimens were examined for ova of the helminth or platyhelminth group. On some of the specimens, examinations were made for protozoa. Stools were examined by the simple flotation technic with a concentrated sugar solution used for ova of helminths and flukes. Direct normal saline smears were used to determine the presence of protozoan cysts. When protozoan cysts were indicated by the smears, iron-hematoxylin stains were made for positive identification.

Kahn tests. Kahn tests were given to 16,320 inhabitants, nearly three fourths of the population of the Saipan, Palau, and Marshall Islands districts The Kahn-test technic was not uniform throughout the portion of the survey reported here. Originally, the three-tube test was utilized. Later, using this same Kahn antigen, only the second dilution was used. Approximately 50 percent of the Kahn tests were positive, with little difference between the sexes (tables S and 9). The Palau district had by far the highest proportion of positives-65.4 percent as compared with 35.1 percent for the Marshall Islands district and 30.4 percent for the Saipan district.

Wide variation occurred among the islands, paralleling the variation between the districts. Tabal Island in the Marshall Islands

 Table	8.	Distribution	of	islands	Ьŗ	percent	of	
		positive K	ahn	reaction	14			

Percent of positive Kahns	Total num- ber of islands	Saipan district	Palau district	Marshall Islands district
Under 25	2	1	- 1	I
25 to 49	17	-	6	11
50 to 74	13	2	7	4
75 and over	7	-	7	- 1

1209

OCTOBER 1959

Table 9. Percentage distribution of examined population with positive Kahn reactions by age group

	Percent positive					
Age group (years)	Tota)	Saipan district	Palau district	Marshall Islands district		
All ages	50. 1	20.4	6 5. 1	38.1		
Under 5	37.6	41.0	61.3	18.9		
5 to 14	42.9	18.3	65.0	25.9		
15 to 24	49.5	33.7	66.4	37.4		
25 to 44	53.8	36.7	66.3	43.0		
45 and over	52.9	32.0	63.9	44.0		

district had the lowest percentage of all, only 3.6 percent positive. The next lowest was Saipan Island, with 22.8 percent. At the other extreme was Elato Atoll in the Palau district, the only place where every individual tested had a positive reaction. Other islands or atolls where over 75 percent of the population had positive reactions were Kayangel (95.4), Satawal (92.9), Woleai (89.1), Peleliu (87.0), Lamotrek (84.2), and Faraulep (77.3), all in the Palau district.

Generally speaking, the proportion of positive reactions increased with age up to the age group of 45 and over, where a slight decline was noted. The same rate of increase from age group to age group, however, was not observed in each district. In the Saipan district a marked increase occurred between the 5- to 14- and 15- to 24-year age groups, with little change in the age groups thereafter. The conspicuously high rate for Saipan's under-5-years-of-age population, however, is based on only 39 tests. Very little variation is observed between the age groups in the Palau district. On the other hand, the Marshall Islands district exhibits definite increases with each age group.

In this survey a positive Kahn test was considered likely evidence of the presence of yaws infection. It is recognized that the Kahn test is far from being an infallible index for the prevalence of yaws, for a positive reaction to the test may result from many diseases and conditions. Because malaria was not noted in this area and because syphilis, leprosy, and infectious mononucleosis seldom occurred in proportion to the number of positive Kahn tests, it would appear to be a reliable guide in this instance.

Yaws is not, as commonly believed, a venereal disease, but is a contagious disease that may be transmitted from person to person by

VOLUME 10, NO. 10

1210

5013262

÷

direct contact or carried by flies. The spirochete causing yaws, *Treponema pertenue*, enters through open lesions such as scratches and minor cuts. The primary lesions appear most frequently on the lower extremities and may last from 3 months to 3 years. It is not unusual for old healed-over lesions to break down. Tertiary-stage lesions often cause severe scarring and, in the form of gangosa, sometimes completely destroy the features.

Many inhabitants had multiple scars as evidence of a history of yaws. Very few diagnoses of active yaws were established during the course of the health survey. As shown in table 10, most of these cases were found in the Marshall Islands district. Numerous cases of active yaws were suspected, but dark-field examinations of the lesions were negative, possibly due to local treatment of ulcerative lesions with merthiolate and mercuric ointments by the native health aides.

Table	10.	Results of survey for active yates, Saipa	n,*
	Pe	lau, and Marshall Islands districts	

District and island or atoll	Number of cases	District and island or atoli	Number of cases
Total	73	Marshall Islands.	
Palau	2	Majuro Atoli	T
		Alluk Atoll	1
Babelthuap Island	1	Aur Island	3
Elato Atoll	1	Liklep A toll	3
	J	Wotje Atoli	3
Marshall Islands	- 71	Kwajalein Atoli	2
		Namorik Island	2
Ine Island	12	Tabal Islaud	2
Ebdu Atoli	111	Utirik Atoli	2
Maloelap Atoll	11	Lib Island	· 1

"No active cases of yaws reported for the Saipan district.

During the survey (table 11) urogenital examinations were made only of the male population, as the women were reluctant to permit examination. Little or no acute urethritis from gonococcus was noted but, although cases were not reported on the health record, medical officers at some of the hospitals stated that among the women there were many cases of "salpingitis, gonococcic" and "cervicitis, acute, due to gonococcus." The presenting symptoms of the reportedly widespread gonococcal infection among the men were those of a nongonococcic urethritis. Syphilis was almost nonexistent, a condition found to parallel the prevalence of yaws. Very few cases of primary chance were found and secondary skin manifestations oc-

OCTOBER 1959

5013263

. بىلىلە

and the second second of the second
.

÷

ŗ

÷

District and island or stoll	Syphills	Gonorrhes
Total number of cases	6	12
Palau	-	2
Koror Island	-	1
Yap Island	-,	1
Marshall Islands	6	10
Alluk Atoli	-	1
Arno Island	1	1 -
Ebon Atoll	1	-
Kwajalein Atoll	- 1	3
Majuro Atoll	4	4
Namorik Island		1

Tuble 11. Results of survey for syphilis and gonorrhea, Saipan,* Palau, and Marshall Islands districts

. "No cases of syphills or gonorrhea reported for the Saipan district.

curred infrequently among both men and women. Several cases of syphilis and gonorrhea were reported from the Majuro and Kwajalein Atolls but only an occasional case was reported from any of the other islands or atolls.

Chest x-ray examinations. Chest roentgenograms were made of approximately 82 percent of the population of the Saipan, Palau, and Marshall Islands districts (tables 12 and 13). Many of those not

X-ray findings	Total		Salpan district		Palau district		Marshall Islands district	
,	Number	Percent	Number	Percent	Number	Percent	Number	Percent
X-ray examinations All chest defects	18.094 817	100. 0 4. 5	4, 226 230	100. 6 5. 4	8, 733 388	100.0 4.4	5, 135 199	100, 0 3, 9
Tuberculosis, pul- monary	267	1, 5	85	20	141	1.6	41	.8
Moderstely and far-advanced Other active Arrested	94 147 26	.5 .8 .1	24 44 17	.6 1.0 .4	33 51 7	, 6 . 9 . L	17 22 2	.3 .4 .0
Chest tumor Cardiovascular ab-	31	.3	7	. 2	17	. 2	7	.1
normality Bone deformity Other	48 32 439	.3 .2 2.4	11 3 124	.3 .1 29	11 2 217	.1 .0 2.5	26 27 96	.5 .5 2.0

Table 12. Results of chest x-ray examinations

1212



VOLUME 10, NO. 10



i

5.3

1

ł

Table 13. Results of chest x-ray survey for active pulmonary tuberculosis by age group

	Active by x-ray								
Age group (years)	Total		Salpan district		Palau district		Marshall Islands district		
	Number	Percent	Number	Percent	Number	Percent	Number	Percent	
All ages	241	1.3	68	1.6	134	1.5	39	0.7	
Under 5	5	. 5	3	.8		0	2	. 5	
5 to 14	15	.3	5	.4	ि हो	.4	2	.2	
15 to 24	30	1.1	16	20	17	LI	6	. 5	
25 to 44	67	1.2	17	L.S	. 43	. 1.5	7	. 5	
45 and over	115	3.2	27,	4.4	66	3. 5	22	2.0	

x-rayed were aged and infirm. The initial examination consisted of a 35-mm photofluorographic roentgenogram that, if the x-ray shadows were suspiciously abnormal, was followed by a 14- by 17inch chest roentgenogram. The findings reported in this survey were made from the 14- by 17-inch films without further clinical study.

On the basis of these examinations 4.5 percent of those x-rayed were found to have chest defects, and 1.3 percent were suspected of having active pulmonary tuberculosis. The number of chest tumors, cardiovascular abnormalities, and bone deformities was negligible, none being over 0.3 percent of the total x-rayed.

The prevalence rates for active pulmonary tuberculosis increase with age, showing but little variation by sex. The proportion of active-pulmonary-tuberculosis suspects among those x-rayed is greatest for those 45 years of age and over, 3.2 percent as compared with the next highest rate of 1.2 percent for the 25- to 44-year age group. Only slight differences were noted between rates for both sexes, the overall rate for females exceeding that for males by 0.1 percent.

Of the three districts Saipan has the highest percentage of chest defects, with Palau ranking second and the Marshall Islands lowest of all, 5.4, 4.4, and 3.9 percent, respectively. Of the persons x-rayed only 0.7 percent in the Marshall Islands, 1.5 percent in Palau, and 1.6 percent in Saipan were recorded as being suspected of having active pulmonary tuberculosis.

On the basis of the chest roentgenograms performed (table 14), 10 islands had no active pulmonary tuberculosis. On the other islands the percentage of positive films for the x-rayed population ranged from Kwajalein's 0.1 to Faraulep's 6.8. On six of these islands or

OCTOBER 1959

Table 14. Percenti putmonary tube	reul osis b	tx-raysindu r rank orde	ntite of act	ice of
Islands or atoll Iands districts	s. Saipan,	Palnu, and	Marshall	14-

Island or atoll and district	Percent sctive	Island or atols and district	Percent active
Faraulan Atoll*	1	World Arall**	17
Tara toll	1 1 1	I Maiuro Atoli	
Entered Island*	3.6	Malit Ish-des	
Tomotrek hight	1 3.0	Babalthran Island	5
Tabal Island**	1	Satamal Lind	1 6
The Islands	3 9 1	Ebon Ardise	
Ane Lyland	1 3.1	Liber Areles	
Mar Alon -	1 2.3	Autox Aros: Nomenik Liberd 77	
FRIS ISLAND	20		
sap Islands	10	Kwajsien Aton	
Am Elono.	21	Angaur Bling	
Koror Island	2.4	Eauriph Atoll	1 0
Maloelap Atoli"	2.2	Lib Island**	{ 0
Kayanget Atoll*	21	Likiep Atol)**	1 0
Ngulu Atoli"	21	Merir Island*	1 0
Tinian Island***	\$ 5.8	Peichu Island*	1 0
Halik Atol!	1.7	Pulo Anna Islawd*	5 0
Saipan Island***	1.6	Tobi Islas J.	0
Wolesi Atoll	1.3	Clithi Atoli"	} 0
Arno Island**	1.2	Utirik Arel!**	{ o
Rota Island***	1.2		

*Palou district.

**Marshall Islands district.

***Salpan district.

atolls, Ine, Tabal, Lamotrek, Sonsorol, Elato, and Faraulep, over 3 percent of the x-rayed population had active pulmonary tuberculosis.

An interesting experience was reported from Tabal Island. The U.S.S. Whidbey visited this island in November 1948 but, because heavy seas rendered the x-ray machine inoperable, the ship was forced to return at a later date to complete the examinations. During the initial visit one 39-year-old woman on the island showed clinical evidence of pulmonary tuberculosis and had positive sputum for tubercle bacilli. At that time nine people were living in the same house with this active case of pulmonary tuberculosis. Eighteen months later when the survey ship returned the woman had died of tuberculosis. The chest roentgenograms for the other nine members of the household were still negative for tuberculosis.

Table 15 shows the diagnoses of tuberculosis that occurred among inhabitants who were not x-rayed. These diagnoses had been established prior to the health survey and, in some cases, the individuals were receiving treatment for the disease.

Nearly 90 percent of the population were tuberculin-tested. Table 16 shows the distribution of islands by percent of positive tuberculin

VOLUME 10, NO. 10

5013266

.

1

22

u,

ŧ

li L

· · · · · · · · · · · · · · · · · · ·			• • · · · •	Tuber	culosis			
District and island	Total	Pulmonary		Skin	Ismph	Male	Bronchus	Other
	CillSes	Active	Arrested		nodes	genitalia		
Total	н	9	3	6	14	1	1	10
Salpan: Salpan		. 	:					
Island	6	1	- 1	2	1	· _	[_]	2
سلد	33	4	3	3	13	1 1	1	8
Babelthuap		1 ·						
Island	4	1	1 - 1	1	1.	- 1	-	1
Fais Island	1	-	-	-	-	- 1	- 1	1
Koror Lind	4	- 1	1 - 1	-	3	- 1		1
Neulu Atoll	1	- 1	1 - 1			1	-	-
Pelebu Island	2	<u>-</u>] _]	-	· -	-	-	2
Satawal Lind	I	-	- 1	-	1	-	-	-
Sonsorol Island	1	1	1 - 1		-	- 1	9 - 1	-
Yap Islands	19	2] 3]	2	8	- 1	1 1	3
Marshall Islands	5	4	-	1	-	-	-	-
Atluk Atoli	2	1	-	1.	-	- 1	-	-
Kwajalem Atoll	1	1.	_]	-		-	-	-
Majuro Atoli	1	1	- 1	-	-	-	-	-
Tabal Island	1	1	-	-	-	- 1	1 - 1	-

Table 13. Diagnoses of tuberculosis for natives not x-rayed

tests. Two test-strength dosages of tuberculin, 0.0001 mg PPD and 0.00002 mg PPD, were used for the tests, the latter dosage being used early in the program and occasionally when the 0.0001 mg PPD dosage was not available. On most of the islands in the Marshall Islands district the 0.0001 mg PPD dosage was used, while primarily the 0.00002 mg dosage was used in the Palau district. Both dosages were used in the Saipan district. From Babelthnap Island, where the 0.0001 mg PPD tuberculin dosage was used on about one half of the tested population and the 0.00002 mg PPD tuberculin dosage was

Table 16.	Distribution of islands by percent	of
	positive tuberculin tests	

Percent of positive tuberculin tests	Total	Saipan district	Palau district	Marshail Islands district
Under 9	. 3	-	1	2
10 to 19	6	-	-	6
20 to 29	7	- 1	3	4
3 0 to 39	8	1	3	4
40 to 49	3	1	2	- 1
30 to 59	6	1	5	1 -
40 to 69	4	-	4	-
70 to 79	1	-	1	i -
80 and over	1		1	-

OCTOBER 1959

5013267

!

ì

1215

ί

E

used on the remainder of the population, it was reported that there was no significant variance between the number of positive reactors to the two dosages.

The findings for the Yap Islands presented in this article are all based on dosages of 0.00002 mg PPD tuberculin. In 6 of the 10 districts of Yap, however, all natives with negative reactions to these first tests were given a second test of 0.005 mg PPD tuberculin. Out of 304 tests there were 251 positive reactions. In other words, of the negative reactors to 0.00002 mg PPD test-strength tuberculin, who were retested with 0.005 mg PPD test-strength tuberculin, approximately 83 percent reacted positively. The practice of employing second test-strength dosages of tuberculin was discontinued. It was found that most individuals reacted to the second dose and the reactions were regularly so severe that they were alarming the natives.

Of the tuberculin-tested population 46.5 percent were positive reactors. The Marshall Islands district had much the lowest percentpositive, 19.2 percent as compared with 54.6 percent for the Saipan district and 60.0 percent for the Palau district.

There is wide variation among islands in the proportion of positive reactors (table 17). The widest range occurs in the Palau district,

Island or atoll and district	Percent positive	Island or stoll and district	Percent positive
Yap Islands"	BL. 5	Lib Island	34.0
Fais Island*	79.0	Tabal Islami***	32.9
Tobi Island"	66.4	Aur Island***	30.0
Koror Island*	63.8	Eauripik Atoll"	23.9
Merir Island*	63.6	Mill Aron-	27.9
Pelaliu Island"	62.4	Namorik Island***	27.5
Angaur Island*	59.7	Majuro Atal	25.6
Salpan Island**	58.4	Pulo Anna Liland*	23.1
Sonsorol Island*	57.9	Kwajalein atoil-**	22.3
Ngulu Atoli"	36.3	Satawal ILand*	20.0
Wolesi Atoll"	54.6	Wotje Atat	19.2
Ulithi Atoll*	52.5	i Meht Islam!	18.0
Babelthuap Island*	47. 5	Maloelap atell***	16.4
Rota Island**	42.1	Allok Atazert	13.3
Kayangel Atoll*	40.2	Utirik Atedata	12.6
Lamotrek Atoll*	38.6	Likiep Atiit***	10.1
Ualik Atoll"	3%.5	* Eluto Att2= *	6.1
Timan Island**	36.4	Arno Istant	4.3
Ine Island***	36.0	Ebon Attd***	1 2
Faraulep Atoll*	34.5		

Table 17. Percentage of positive tuberculin tests by rank order, selected islands or atolls

*Palan distruct

**Salpun district

*** Marshall Islands district.

1216

<u>5013268</u>



VOLUME 10, NO. 10

C

from 6.1 percent for Elato Island to 84.5 percent for the Yap Islands. All of the islands in the Marshall Islands district had less than 40 percent positive reactors. The two lowest percentages in this district were those for Ebon Atoll and Arno Island, 3.2 percent and 4.3 percent, respectively. Most consistency was shown in the Saipan district, where the percent positive was 36.4 percent for Tinian Island, 42.1 percent for Rota Island, and 55.4 percent for Saipan Island.

The proportion of positive reactors progresses steadily with age (table 18). The only exception is the Saipan district where the peak is reached in the 15-to-24-year age group and a slight decline occurs after the age of 44. The percentages for both sexes in each district are very close. Where substantial differences exist the higher percentage is for males.

Table 18.	Percentage of positive reactions among
tuberculln	-tested population by age group and sex

	Percent positive					
Age group (years)	Total	Saipan district	Palsu district	Marshall Islands district		
All ages	46. 5	54.6	60.0	19.2		
, Male	47.6	54.4	61. 2	22.1		
Female	45. 3	54.8	58.8	16. 2		
Under 5	9, 1	8.0	16.3	1.7		
\$ to 14	32.7	41.3	40.1	6.9		
15 to 24	53. 3	78.7	65.4	20.9		
25 10 41	60. 3	73.0	73.5	27.7		
45 and over	58.9	71.1	76.2	27.7		

Skin diseases. Skin diseases are prevalent throughout the territory. The low standards of hygiene among the islanders and the humid, hot climate of the area are all conducive to these conditions.

Dermatophytosis is the most common skin disease. The prevalence rate for the three districts combined was 295.5 per 1,000 inhabitants. The Saipan district had the highest rate for this disease, 379.9 as compared with 305.7 for the Marshall Islands and 253.7 for Palau. Over one-half the population of Tabal, Aur, Faraulep, Satawal, Rota, Wotje, Lamotrek, and Maloelap had dermatophytosis. In this class tinea versicolor is in such common occurrence that it is ignored by the natives. Tinea cruris and tinea circinata are also prevalent.

The prevalence rate for diseases of skin and cellular tissue was 149.2 per 1,000 persons. This group of diseases includes a high proportion of local skin infections. These include ecthyma, which is especially

OCTOBER 1959

prevalent in this area, and such conditions as diseases of the sweat glands, dermatitis seborrheica, and molluscum contagiosum. Rates were much higher in the Saipan and Marshall Islands districts than in the Palau district—246.2 and 214.2 per 1,000 inhabitants as compared with 62.9, respectively.

MISCELLANEOUS DISEASES

Prevalence rates for diseases which occurred with greatest frequency are presented by district in table 19. These diseases were diagnosed from the general physical examinations rather than from special tests and examinations.

Table 19.	Morbidity for selected diagnoses
(D	release rates per (000 examinations)

Belected diagnosis	Ťotal		Saipan district		Palau district		Marshall Islands district	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Examinations	22, 146 ,	(*)	4, 999	(*)	10, 575	(*)	6, 572	(*)
Dermatophytosis Chronic tonsillitis and nasopharyngi-	6, 611	296. 5	1.899	379. 9	2, 683	253. 7	2, 029	308.7
tis Diamantalia	3, 326	150. 2	1, 433	286.6	555	52.5	1, 338	203.6
and cellular tissue.	3, 304	149.2	1, 231	246. 2 80. s	665 675	62. 9 63. 8	1.408	214. 2
Desentrative joint	~	P. 4		QU. 0	0,3	0.0	· ~	110.0,
disease	1, 563	70.6	262	52.4	900	85.1	401	61.0
Conjunctivitis	909	41.0	273	54.6	443	41.9	193	29.4
Cataract	784	35. 4	87	17.4	519	49.1	178	27.1
Acute respiratory infections	- 632	28.5	426	6 5. 2	123	11.6	83	12.6
Neoplustic diseases	325	14.7	60	12.0	74	7.0	191	29.1
Informatory diseases			1 1		1	•		
ofear	217	9.8	132	26.4	10	. 9	75	11.4
Opacity of cornea	211	9.5	82	16.4	5G	8.1	43	6.5
Vitamin deficiency	195	8.8	101	20. 2	28	2.6	66	10.0
Anemia	185	8.4	64	12.8	4	. 4	117	17.8

Not applicable.

The most prevalent diseases in the Saipan, Palau, and Marshall Islands districts were skin diseases, diseases of the eye, and respiratory diseases. Leprosy was also of major importance. Other diseases of wide prevalence were degenerative joint disease, neoplastic diseases, rickets and vitamin deficiency, anemia, and inflammatory diseases of the ear. Diseases carried by mosquitoes and other arthropods do not assume the importance in the trust territory that is usual in other tropical areas. Malaria is not an indigenous disease at the present

1218

VOLUME 10, NO. 10

2

e

time because of the absence of anopheline mosquitoes. None of the rickettsioses were present. Although dengue typhoid fever, and amebic dysentery were formerly regarded as widespread in the territory, no cases were reported present in these three districts during the survey. Several cases of filariasis were reported from the Palau district and one case from the Saipan district.

It should be borne in mind that some of the variations occurring among the islands may be due to the fact that examinations on the various islands were performed during different seasons of the year and by different personnel.

Diseases of the eye. Over 4,000 cases of diseases of the eye and adnexa oculi were reported from this area; the prevalence rate was 198.4 per 1,000. Pterygium, conjunctivitis, and cataract were greatest in occurrence. The intense sunlight, fine coral sands blowing about, the use of open fires, and frequent submersion in salt water are considered factors possibly contributing to the wide prevalence of pterygium and conjunctivitis. Cataracts were mostly of the senile type. Eighty-five cases of blindness were reported: 17 of these were bilateral and 68 were unilateral. It is believed that most of these were either congenital or resulted from physical injury.

In comparison with the other districts, the Mershall Islands had a strikingly high rate for pterygium, 146.5 per 1000 persons as compared with 89.8 for Saipan and 63.8 for Palau. Conjunctivitis was most prevalent in the Saipan district, while cateracts occurred most frequently in the Palau district.

Acute respiratory infections. Acute respiratory infections were in common occurrence throughout the three districts. The over-all prevalence rate was 28.5 per 1,000 persons. This rate was much higher in the Saipan district than in the Marshall Islands and Palau districts, 85.2 per 1,000 persons as compared with 12.6 and 11.6 respectively. The changeable humid climate, crowded living conditions, sleeping on floors, low levels of nutrition, and poor standards of sanitation were undoubtedly factors contributing to these diseases. Saipan Island had an unusually large number of acute respiratory infections for its examined population. Chronic tonsillitis and nasopharyngitis were widespread in this area. The prevalence rate for the 3 districts combined was 150.2 per 1,000 persons. The rate for the Palau district, 52.5, was far below those for Saipan and the Marshall Islands, 286.6 and 203.6, respectively.

Leprosy. Eighty-six lepers from various parts of the trust territory were under treatment at the leprosarium on Tinian Island at the time of the health survey. Of these cases, 25 were thought to be lep-

OCTOBER 1959

50132

romatous, 59 tuberculoid, and 2 mixed. As shown in table 20, 83 additional cases clinically suspicious for leprosy were found in the Saipan and Palau districts during the course of the survey. None were reported from the Marshall Islands district.

Table 20. Results of survey for clinically suspicious cases of leprosy, Saipan, Palau, and Marshall Islands^a Districts

District and island or atoll	No. of cases	District and island or atoli	No. of cases
Total	83	Palau-Continued	3
Saipan: Saipan Island	65	Koror Island	2
		Setawal Island	2
Palau	18	Wolesi Atol!	2
Lamotrek Atoli	4	Elato Ato!1	1
Babelthuap Island	3	Fais Island	1 T

* No clinically suspicious cases of leprosy for Marshull Islands district.

The fact that the true macular lesion is not readily recognized by other than specialists in the field may have reduced the discovery rate for leprosy in this survey. Recognition of leprosy was especially difficult because of the prevalence of tinea versicolor and traumatic contracture, both of which required differentiation from the disease. Personnel conducting the survey were reluctant to make a diagnosis of leprosy where there was any question of its certainty.

Diseases of ear. Inflammatory diseases of the ear occurred at the rate of 9.8 per 1,000 inhabitants. The majority of cases were in the Saipan district where the rate was 26.4 per 1,000 persons. The rate in the Marshall Islands district was 11.4 and only 0.9 in the Palau district.

The most prevalent disease of the ear was "infection, diffuse, external auditory meatus." It is an interesting fact that of the 161 cases reported, 101 were on Saipan Island. It is not known whether personnel conducting the survey were more thorough in making examinations on this island than they were elsewhere or whether an epidemic was in occurrence at the time of the survey.

Other conditions. Degenerative joint disease was rather prevalent. showing a rate of 70.6 per 1,000 inhabitants in the three districts combined. Rates were highest in the Palau district and lowest in the Saipan district: This disease was most prevalent among the aged.

A total of 325 neoplastic diseases was reported, establishing a prevalence rate of 14.7 per 1,000. Only 11 neoplasms were proved malignant, although other malignancies were suspected. The malig-

VOLUME 10, NO. 10

1220

nant neoplasms were distributed among the islands or atolls as follows: Saipan, 3; Babelthuap, 2; and 1 each on Tinian, Ulithi, Woleai, Yap, Kwajalein, and Wotje. Among the 312 benign neoplasms there were 114 melanomas of skin, 63 lipomas, and 29 hemangiomas and lymphangiomas. The bulk of these were on Majuro, Ebon, Saipan, Yap, Babelthuap, Maloelap, and Mili.

The prevalence rate for vitamin deficiency was 8.8 per 1,000 inhabitants. Of the 195 cases reported, 98 were on Saipan Island where the prevalence rate was 24.2. Other islands where the rate was high were Ine, 72.8; Arno, 57.6; and Mili, 2S.2.

Anemia, with a prevalence rate of 8.4 per 1,000 persons, was reported in substantial numbers from Saipan, Majuro, Ebon, and Mili. According to reports of this survey anemia was practically nonexistent in the Palau district.

It was reported that a large number of cases of leukoplakia buccalis were seen on Saipan and Babelthuap Islands. These occurred principally among betel-nut chewers. A comparatively small number of cases were reported on the health record cards used in this study for tabulation purposes. In the summary reports accompanying the cards, however, it was estimated that on Saipan alone over 200 cases of leukoplakia buccalis were observed. Many of these were verified by biopsy. It was further pointed out in the reports that, although betel-nut chewing was also a common practice on the Yap Islands, leukoplakia buccalis was not prevalent there. It was suggested that this might have resulted from the oral mud packs used on Yap to hide the staining effect of the betel nut. The mud packs were not used on Babelthuap. The reports from Saipan did not state whether mud packs were used there.

SUMMARY

In the summer of 1948 a survey staff aboard the U.S.S. Whidbey began a health survey of the Trust Territory of the Pacific Islands. its objective being a physical examination of every inhabitant of the area. This report covers the civil administrative districts of the Saipan, Palau, and Marshall Islands. Fifty-two percent of all stools examined were positive for intestinal parasites. The most prevalent intestinal parasites were hookworm, Trichuris, and Ascaris. Approximately 50 percent of the Kahn tests were positive. This was judged to be evidence of the yaws infection on the islands. Of those receiving chest x-ray examinations, 4.5 percent were found to have chest defects and 1.3 percent were suspected of having active pulmonary tuberculosis. The number of chest tumors, cardiovascular

OCTOBER 1959

5013273

sbnormalities, and bone deformities was negligible. The incidence was 0.3 percent or less of the total number of inhabitants x-rayed.

The general physical examinations disclosed that the most prevalent diseases were skin diseases, diseases of the eye, and respiratory diseases. Leprosy was also of major importance. Other diseases of wide prevalence were degenerative joint disease, neoplastic diseases, rickets and vitamin deficiencies, anemia, and inflammatory diseases of the ear.

ACKNOWLEDGMENT: This report was made possible by many persons within the Bureau of Medicine and Surgery, Department of the Navy, who contributed to the initiation, development, and accomplishment of the health survey itself and, more specifically, by the personnel of the medical statistics division who so painstakingly processed the data. The author is especially indebted to Dr. W. V. Charter, director of the division, for sponsorship of the study; to Mr. Louis P. Hellman, deputy director, for his guidance and invaluable suggestions; and to Mrs. Helen Rod for the manuscript preparation. The professional advice relating to the medical aspects of the data that was so generously provided by personnel of the preventive medicine division is gratefully acknowledged.

REFERENCES

1. HANDBOOK ON THE TRUST TERRITORY OF THE PACIFIC ISLANDS. Navy Department, Office of the Chief of Naval Operations, Washington, D.C., 1948.

- 2. HANDBOOK ON THE TRUST TERRITORY OF THE PACIFIC ISLANDS. Navy Department, Office of the Chief of Naval Operations, June 1950.
- 3. BIRTHS BY AGE OF MOTHER, RACE, AND BIRTH ORDER, UNITED STATES 1949. Federal Security Agency, U.S. Public Health Service, National Office of Vital Statistics, 15 Oct. 1951, p. 145.

PROGRESS IN DIABETES

Diabetes continues to maintain an unchallenged position in the study of clinical conditions chiefly because it represents all aspects of medicine. The number of diabetic patients increases each year, not because it is as such on the increase, but rather because the thousands of undiscovered cases are now being discovered. . . . The future of diabetes is brighter now than ever in the past. Research may bring about fewer complications, oral management and even eventual cure, but our present knowledge. when applied, makes the future more hopeful. Early diagnosis, better management, control of disabling conditions and efforts in the prevention of diabetes are all important steps toward the goal we are all striving for, namely, make the two million known diabetic patients and the one and one-half million undiscovered cases a valued part of everyday living because they are quite like their nondiabetic neighbor .-- I. W. WILKENS: The Future of Diabetes. The Journal of the Indiana State Medical Association, October 1958.

2222

j |

1

VOLUME 10, NO. 10

. . .

1

;

ŧ.

5