408283

R

May 4, 1966

MEMORANDEM FOR CHAIRMAN SEABORG

COMMISSIONER PALTREY

COMMISSIONER RAMEY

COMMISSIONER TAPE

THROUGH GENERAL MANAGER

SUBJECT: CUREENT STATUS OF THYROID MODULES AMONG THE MARSHALLESE EXPOSED TO FALLOUT FROM THE BIKINI TEST OF MARCH 1, 1954

In our last memorandum on this subject, October 11, 1965, it was noted that a cumulative total of 11 instances of nodular thyroid glands had been found in the 69 surviving Marshalless who had been exposed to fallout from the subject test while on Rongelap and on Ailingnae. All but one of these cases were in young people who were less than 10 years old at the time of exposure. The single exception was in a 41-year-old woman (approximately 30 at time of exposure) who was found to have a carcinoma of the thyroid at operation.

Br. Robert A. Conard of the Medical Department of Brookhaven National Laboratory, and leader of the annual (or more often) medical surveys of these people, has just returned from the latest survey conducted during February and March of 1966. He reports that an additional five cases of nodular thyroics have been diagnosed in this group. The present cumulative incidence is shown in the following table.

Age at Exposure	Skyosed	Total Incidence of Hodules Through March 1966	
		Mumber	Percent
1-5	±6	ŷ	56
6-10	¥	4	44
11-15	7	Q.	-
16-20		₩	•
- 20	3.7		
/	Q -3	3. %.	N 25

BEST COPY AVAILABLE

of 14 K plus proportionately smaller internal thyroid doses were examined for nodular thyroids; none were found.

Among 194 unexposed controls who are now living on these islands, three cases of thyroid nodules were found, all in adults more than 47 years. Other forms of thyroid disease were not seen.

1 10 TAG

You will be kept informed of the progress of these studies; the next noteworthy item will be the clinical and possibly surgical findings in the five Marshallese coming to Brookhaven.

> C. L. Dunkem, M.D. Director Division of Biology and Medicine

> > 140 F. S.

39 1

- 第3章 1788年 (1982年 - 1982年 -