(Health and Safety - HC-4500)

UC-41

410200

A RECONSTRUCTION OF CHRONIC DOSE EQUIVALENTS FOR RONGELAP AND UTIRIK RESIDENTS - 1954 TO 1980

E.T. LESSARD, N.A. GREENHOUSE, AND R.P. MILTENBERGER

October 1980

BEST COPY AVAILABLE

SAFETY AND ENVIRONMENTAL PROTECTION DIVISION

BROOKHAVEN NATIONAL LABORATORY UPTON, NEW YORK 11973

Coded AS 4350 /

888 51257 86-41 (H-alth and Safety - 11(-1120)

A RECONSTRUCTION OF CHEOMIC DOSE LOFT ALL LET > OR RONGELAP AND UTICHC RECIDENTS - LOVE TO LEMO

FT ELSARD IA GREENMARY AND R.P. BULLET CO.

DISCLAIMER

11

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency, contractor or subcontractor thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency, contractor or subcontractor thereof.

> Printed in the United States of America Available from National Technical Information Service U.S. Department of Commerce 5285 Port Royal Road Springfield, VA 22161 Price: Printed Copy \$7.00; Microfiche \$3.50

A RECONSTRUCTION OF CHRONIC DOSE EQUIVALENTS FOR RONGELAP AND UTIRIK RESIDENTS - 1954 TO 1980 E. T. Lessard, N. A. Greenhouse, R. P. Miltenberger

ABSTRACT

From June 1946 to August 1958, the U.S. Department of Defense and Atomic Energy Commission conducted nuclear weapons tests in the Northern Marshall Islands. BRAVO, an aboveground test in the Castle series, resulted in radioactive fallout contaminating Rongelap and Utirik Atolls. On March 3, 1954, the inhabitants of these atolls were relocated until radistion exposure rates declined to acceptable levels. Environmental and personnel radiological monitoring programs were begun in the mid 1950's by Brookhaven National Laboratory to ensure that dose equivalents received or committed remained within U.S. Federal Radiation Council Guidelines for members of the general public. Body burden and dose equivalent histories along with activity ingestion patterns post return are presented. Dosimetric methods, results, and internal dose equivalent distributions for subgroups of the population are also described.

1170s. Your persons visited Argonne from Rongelap and, in addition, INTRODUCTION entry of the sector ibar teachers and the

On March 1, 1954, at Bikini Atoll, BRAVO, the first of six nuclear weapons hinne**un** yne ee bei te A War and tests in the Castle series, was detonated. The BRAVO device caused substantial a Frazil Character and Validada a 200462 surface contamination on inhabited stolls within a 2,000 square mile area. The I he mine to a familiant contaminated region was cigar shaped and included Ailinginae, Rongelap, Rongerik, and Utirik Atolls which lay east of ground zero at distances from 60 provide the comparent of a constrained of the second of the second s to 300 miles. The fallout on Rongelap, initially visible at H+6 hours, had thinned out to the extent that it was no longer seen at H+10 hours (G162).

1.6

the net odd mode of the two odd of the manual some of On March 3, 1954, the 64 residents of Rongelap Atol1 and 18 residents of and the Headmerel way to reat as as to for the same Sifo Island, Ailinginae Atoll, were evacuated. On March 3 and 4, evacuation of 91611-0-0 alder report and two diff and the 157 Utirik Atoll residents also took place. During the first few weeks and at HIRRARIS AG. 11 least once every year from 1957 to the present, a Brookhaven National Laboratory K SE medical team, organized by the Department of Defense and by the Atomic Energy THE SECTION ALL FRAME i sare e 1. 1. 1. 1. Commission and its successor organizations, has provided medical examinations to monitor the health of the persons initially affected by the fallout from the nuclear testing program, plus a comparison population. Reports of their findings are given in Cr56, Co58, Co59, Co60, Co62, Co63, Co65, Co67, Co70, Co75, and Without Bride W Co80.

The Utirikese and Rongelapese returned to their home atolls in June 1954 and in June 1957 respectively. The earlier repatriation of Utirik Atoll Wag based on the low level of external radiation exposure measured after the initial 3 month observation period (March to June 1954). The Utirik population was not examined by a Brookhaven medical team until March, 1957, when 144 people received comprehensive physical examinations. Following the 1957, medical survey, two men, removed from Utirik for medical reasons, were whole body counted at Argonne National Laboratory and provided urine samples for radiochemical anal-

ł

MAN TRANSPORT - - LENG 1

1. F. (1994) (1) N. ()

The aforementioned body burden tables illustrate adult mean values for drammed social social relation of an entry of a static static structure of a static

Because of the paucity of measurements at Utirik, information on ⁶⁰Co, ⁶⁵Zn, and ⁵⁵Fe was in some instances derived from the ratio of adult mean body burdens between Rongelap and Utirik. A mean ratio of 2.6 was observed in body burdens for ⁶⁵Zn, ⁹⁰Sr, and ¹³⁷Cs after they reached their maximum values. The standard deviation of this ratio was 15%.

the state of the second

In the following analysis, personal body burden histories and residence intervals, in conjunction with contemporary dosimetric models, are used to estimate internal dose. Dosimetric distributions were constructed from the results and a summary of the derived activity ingestion rates and dose equivalents was provided for various subgroups of the population. Additionally, exposure rate history curves were constructed for each atoll for the period following the

2

Table 1 Rongelap Body Burdens Adult Freeles Adulte Adult Hales Body Body Burden ilusber Days Post Return Jody Burden Number 4.02 of ~ Burden of . of HCi HCL Persone HCI Days 711400 Person 60_{Co} 1.7x10-5 7.8x10-3 7.0x10-3 . 1.1.8 : es 1 2.3=10-5 NA 37 45 **MA** 74 2.9×10-5 144 37 1 1 . 1 4 9.0x10-3 2.2x10+3 aistes 1.0x10⁻² 2.5x10⁻³ 1320 -54..... 45 99 . . . - 2831 ⁶⁵Zn 4.3x10-2 3.8x10-2 HA 4.1=10-2 1 NA 30 32 38 6. . . . 4.3x10⁻¹ 6.2x10⁻¹ 9.5x10⁻² 3.8x10-1 5.0x10-1 12 - alize . 1 4.1=10⁻¹ 5.6=10⁻¹ 304 639 2 27 8.5x10-2 23 9.0x10"2 61 . Sy lac. I 1370 55**7**4 4-1±10-1 4.0x10~1 4.3x10-1 28 32 60 - G. K. K 4626 1.11 90_{Sr} 1.9x10-4 3.7x10-3 1.4m10-4 1.7=10-4 **XA** 4 NA 15 NA 1 3.4x10-3 4.8x10-3 2.8x10-3 3.5x10-3 304 11 1. 6. 4. 4 11. 1 3.7x10-3 5.7x10-3 3.7x10-3 8.8x10-3 7.9x10-3 24 16 639 40 1.6x10-3 7.9x10-3 7.4x10-3 3.0x10-3 8.4x10-3 7.7x10-3 13 1370 2100 9 100 2466 3561 3927 ñ 1 4.6x10-3 3.1x10-3 3.3x10-3 2.8110-3 3.7110-3 12 12 3.7x10-3 3.5x10-3 3.6x10-3 2.5x10-3 1.6x10-3 1.6x10-3 3.9x10-3 4.1x10-3 11 11 ü į3 4292 1.3±10-3 3.3+10-3 ij 8 19 4657 5022 3.1x10-3 2.0x10-3 6.6x10-3 2.8#10-3 à 1.4#10-1 12 13 14 \$ 1 5388 4.2x10-3 1.7x10-3 4.3#10-3 •••• 5753 6118 4 3.3x10-3 4.4x10-3 6.3x10-4 10 2.8=10-3 4 нь 21 с. – р 16A 43 23 NA ò MA 7579 4.6x10-4 5.5=10-4 24 19 \$097 137 C. 1.4=10-2 8.4110-3 NA NA 49 37 1-1=10-2 NA NA 47 NA 164 96 74 89 66 51 37 304 639 1370 2831 5.2x10⁻¹ 4.1x10⁻¹ 6.8±10⁻¹ 5.7±10⁻¹ 6.7±10⁻¹ ····· 8.7×10-1 7.9x10⁻¹ 9.5x10⁻¹ 37 4.7+10-1 9,4x10-1 4.9x10-1 6.8x10-1 45 3.0x10-1 1.9x10-1 24 21 3.9x10⁻¹ 2.5x10⁻¹ 1.7x10⁻¹ 6118 4.8x10-1 3.0x10-1 30 1.8x10-1 1.5x10-1 19 8097 18

NA * Not available

14

5

3

, ,

 $1 \le k_{1} \le 1$

5 · • • · · · 5 • 5

.

大学が大学のないと

の子白い たいしょう

Sales and

.....

-		1.1			- 4 - 3	- 1	
-	-		-	-		0.1	1.

			t-n t	Del	rik Body Burdens	the strength	At Mout -	
* · * ;	Adult (Body Burden uCi	tales Humber of Persons	- 1-24 - 5 + 1 - 5 - 5 + 0 - 1 - 5 + 0 - 1	- Adult Pin Body Burden uGi	thinker of Diates	Body Burden Alf NCi St	Munbar of Ultra	Days Popi Return Days
1 ⁰ co	4.0x10-3 9.7x10-4		0141,7 0141,7 0141,7 0141,7 0141,7	3-1=10-3 7.4=10-4	「「「」」「「」」」、 「「」」」、 「」」」、 「」」」、 「」」」、 「」」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」」、 「」、 「	3.5x10-3 Ad 8.7x10-4 Gr	4 ² Ωiςπέςa ¹² Πίγκει 12 Πίγκει 3 12 Ιαζίας	2464 3924
652m D	3.5x10 ^{-1*} 2.7x10 ⁻¹ 3.7x10 ⁻²	2 16	11 aQ.P	1.6x10 ⁻¹ 3.3x10 ⁻²	15 a area o a	2.1x10-1 3.5x10-2 81	n der e Markene	1734 2464
¹⁵ F¢ D 90 ₂₀	1.7=10-1		28547 29445 8844 2804	1.6x10-1		1.6x10~1	Martin Carlo San Data Karlota Karlota	6114
26	l.4x10 ⁻³ l.2x10 ⁻³ M4 i.5x10 ⁻⁴	5 5 12 14	8,441 9,41,5 7,4 7,4 7,4 7,4 7,4 7,4 7,4 7,4 7,4 7,4	2.4x10 ⁻³ 1.3x10 ⁻³ MA 1.5x10 ⁻⁴	2	1.7x10 ⁻³ 1.3x10 ⁻³ MA 1.5x10 ⁻⁴	7 137,54 8 11 21 x 1 3 26 1 1 3 3 1 21 21 x 1 4 31 21	1734 7213 8669 9225
¹³⁷ C#	4.1x10-1 2.9x10-1 2.6x10-1 1.2x10-1 4.2x10-2	NA 15 9 27 19		2.7x10-2 2.0x10-1 1.3x10-1 7.8x10-2 4.3x10-2	13	3.3±10 ⁻¹ 2.5±10 ⁻¹ 1.8±10 ⁻¹ 1.0±10 ⁻¹ 5.3±10 ⁻²	Colored 30 1000 - 1 30 1000 - 1 22 1000 - 1 36 1000 - 1 36 1000 - 1	1004 1734 7213 8303 9225

flatio derived body burden

Not available Nessured at Argonne National Laboratory

the film and the state of

A Patier was

25 -----

1. 1. . .

1.0

to test. These data, together with appropriate conversion factors and fivily

1.41

are used provided to and the art being , state at

eet.at

 e_{i} point to the decision of the transfer of the transfer

en el Britaneza de la belta el compositor el Christelle en contra en termanita de la





A Second

1. 5. 14 W HILLIAM

BRAVO test. These data, together with appropriate conversion factors and living pattern models, provided an estimate of external dose equivalent.

METHODS

Exponentially declining activity concentrations have been observed in surface soil for ¹³⁷Cs, ¹²⁹I, and ⁹⁰Sr from 1954 to the present on Rongelsp and Utirik Atolls. Declining activity concentrations have also been observed in vegetation at a rate greater than that predicted by radioactive decay. Thus exponential decline in dietary activity was assumed and the following general equations were derived.

 $\lambda P^{\circ} = \frac{U_{0}U_{g}/E_{i} - q^{\circ} (\Sigma_{i} K_{i}X_{i} e^{-(\lambda+K_{i})t})}{\sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{i=1}^{n} \frac{\chi_{i}K_{i}}{K_{i} - K_{E}} e^{-(\lambda+K_{i})t} = e^{-(\lambda+K_{i})t}$ (1) $\lambda P^{\circ} = \frac{\mathbf{q} - \mathbf{q} \circ [\mathbf{L}_{i} \mathbf{X}_{i} \mathbf{e}^{-(\mathbf{\lambda} + \mathbf{K}_{i})t}]}{f_{1} [\mathbf{L}_{i} \frac{\mathbf{X}_{i}}{\mathbf{K}_{i} - \mathbf{K}_{E}} (\mathbf{e}^{-(\mathbf{\lambda} + \mathbf{K}_{E})t} - \mathbf{e}^{-(\mathbf{\lambda} + \mathbf{K}_{i})t})]$ (2)

 $D = f_{1}\lambda^{po} \Sigma_{i} \frac{\chi_{i}}{K_{i}-K_{E}} \left[\frac{K_{i}-K_{E} - (\lambda+K_{i}) e^{-(\lambda+K_{E})t} + (\lambda+K_{E}) e^{-(K_{i}+\lambda)t}}{(K_{E}+\lambda) (K_{i}+\lambda)} \right]$ + q° $\Sigma_i = \frac{\chi_i}{\lambda + K_i} (1 - e^{-(\lambda + K_i)t}),$ (3)

where

or

and

whole body retention at any time t-T of the fraction of initial radioaccivity t E time post onset of uptake, days, λ \equiv instantaneous fraction of atoms decaying per unit time, day $1^{(2)}$ 14/11/1 P° E initial atom ingestion rate, stome day 1, $K_i \equiv$ instantaneous fraction of atoms removed from compartment. i by suidelist common tadi on omit ta vivitus enusphinsteri odo . physiological mechanisms, day", $\chi_i \equiv \text{compartment i deposition fraction},$ $\chi_i \equiv$ the number of Latons in compartment (i relative to the number in all compartments at the onset of declining continuous uptake, (t=0), U = instantaneous urine activity concentration, Bq 2, ,aime the shae an Tani an an $U_{\pm} \equiv$ subject urine excretion rate, \pounds day⁻¹, $f_1 \equiv fraction from GI_3$ $fract, to blood_{f(\lambda+a)}$ f = fraction excreted by the urine pathway, $K_{\rm E} \equiv$ instantaneous fraction of atoms removed or added to the atom uptake per unit time, day 1, due to factors other than radioactive decay, is defining to En sympthy -1. q Ξ instantaneous body burden, Bq, q° \equiv body burden at the onset of uptake, Bq, D = the number of disintegrations in all compartments occurring during the uptake interval, Bq days. 11690

The development of Eqs. (1), (2), and (3) was based on the following convolution integral. At some variable time, τ , defined during a fixed uptake interval, T, the daily activity ingestion rate crossing the gastrointestinal tract to blood is given by

 $\lambda f_1 P^{\bullet} e^{-(k_E + \lambda)\tau}$,

7

* * State Cardina Carlor Baller

1.19.6.14

The whole body retention at any time t-r of the fraction of initial rediosctivity
(ave), ave one of the section of initial rediosctivity
inputed at time r is
yeb, suil in a section of atom is to solve a low on the section is
(
$$T_{i}$$
 (T_{i} (T_{i}) (T_{i}

$$\frac{\lambda P^{o} f_{1} E_{i} \chi_{i} (e^{-(\lambda + K_{E})T} - e^{-(\lambda + K_{i})T} - (\lambda + K_{i}) \emptyset}{K_{i} - K_{E}}$$

in the same pateroph and

As previously stated, Eq. (2) applied at Rongelap and Utirik, it was for the situation that variable time t was the uptake interval. Additionally, persons who returned to the stolls in June 1954 and June 1957 did so with an initial body burden, q°. The behavior of this contribution to body burden, q, was embodied in the q° term of Eq. (2). A similar model was used to relate

urine activity concentration to body burden. Equation 3 was obtained by integrating Eq. (2).

Equations (1) and (2) were used to determine the instantaneous fraction of atoms removed or added to the atom uptake per unit time, $K_{\rm g}$, and then the initial daily activity ingestion rate required to produce the measured or derived body burden. Equation (3) was used to determine the number of disintegrations that occurred in the body during the residence interval of an individual living on Rongelap or Utirik Atol1.

If the mean residence time in the diet is much much longer than the residence interval, then constant continuous uptake is achieved. Equations (1) and (2) can be converted to the constant continuous equations by replacing K_g with $-\lambda$. Single uptake expressions are obtained by setting **P** equal to resort. In some cases only radioactive decay may remove the nuclide from distance items; for these cases K_g would equal zero. In the case of the former Bikini residents, the maturing of coconut trees during residence on Bikini Atoll caused a continuously increasing distary uptake of ¹³⁷Cs. Thus, K_g was found to have a negative value. In the case of Rongelap and Utirik, K_g was found to have a positive value for ¹³⁷Cs, ⁶⁵Zn, ⁶⁰Co, and ⁹⁰Sr. This indicated that in addition to radioactive decay, some other removal mechanism decreased the radipactivity in dietary items during the residence interval. For the nuclide ⁵⁵Fe, only one measurement was published by the BNL Medical Program (Be72); thus an estimate of K_g was not possible.

 K_E was determined by using Eq. (1) or (2) and the population subgroup mean body burden or urine activity concentration. Portions of these bioassay data are illustrated for adult males and females in Figures 2 to 6. Two consecutive urine or body burden data points were used to eliminate the unknown ingestion tivity concentration to body hurden. Equation a was obtained by AST APE DE re million as a maximum since the determined the inclusion of the contract of the rest of would be added to the atom uptobe per ant time, $\xi_{\rm p}$, and have the line by attrivity ingestion cate required to produce the measured or derived () dev. Equation (3) was used to determine the number of it integrations the budy during the levisitence interval of has video with the bost ALDIA SITUDE THE ADD $\phi(x)$ means the event of the diet is much much construction of the theorem in the two the the terms of ter MALES nor ADULTS a gade , feener . 1: X 9 2 1 3 FOPTLOD & COUNT STFEMALES անների heat where is

ംപ

BODY .

102

1954

56

a a

aboi

58 60 62 64

alls optant expressions are obtained by secting 1°

spiles , establishe vom vorab evrandetber vier b

ini kala shikayar yaranga ayar markozî te mar

70 72

CHARLES A MICLORAN PRACE DEDAY OF BOA

78

1014

16.3 15

113

...d

់តផង

-06

a ɗ h

ie

538

10.1

113

11

113

-55

 ~ 1 is

Fig. 2 Mean Adult ¹³⁷Cs Body Burden History at Rongelap Atoll a a de

66 68

YEAR





- ALLES



Fig. 4 Mean Adult ⁹⁰Sr Urine Activity Concentration History at Rongelap Atoll





Generations

ţ

n 12 i e ni

CARANTAL AND AND AND A STATE OF

rate from the equation. This method yields n-1 estimates of K_E where n was the number of data points. An average value of K_E was assigned for each nuclide, and the results for the Rongelap and Utirik populations are given in Table 3. For the evaluation of K_E from Eq. 1 and 2, radiological and physiological parameters were obtained from the open literature (ICRP59, ICRP68, ICRP69, ICRP79, Ki78). A representative sample of these parameters is presented in Table 4.

S	ummary of Dieta	ry Rate Cons	tants (K _E , d	1)
	60 _{Co}	90 _{Sr}	65 _{Zn}	137 Cs
ongelap Adults			· · · · · · · · · · · · · · · · · · ·	
Males	1.5x10 ⁻³	1.8x10 ⁻⁴	3.1x10 ⁻³	1.4x10 ⁻⁴
Females	1.6x10 ⁻³	4.1x10 ⁻⁴	3.5×10 ⁻³	1.4x10 ⁻⁴
Adults	1.5x10 ⁻³	1.9×10 ⁻⁴	3.1×10 ⁻³	1.4×10-4
irik Adults				
Males	N.D.	4.6×10^{-4}	N.D.	1.4x10 ⁻⁴
Females	N.D.	4.0x10 ⁻⁴	N.D.	1.4x10 ⁻⁴
Adults	N.D.	4.2x10 ⁻⁴	N.D.	1.4x10-4

The values of K_E were similar for males and females and for residents of Rongelap and Utirik. For 90Sr on Rongelap a factor of 2 difference between K_E values was observed for males and females. The female parameter for Rongelap Atoll compares with that obtained from the Utirik data. A paired t-test of the Rongelap male and female data indicates that the male/female difference was highly probable and therefore not significant. This difference leads to a

60 17^{Co}

55 26⁷*

0.5 0.3 0.1 0.1

1.0

300

2045 - C. 1 1	ang taon ang sa	···· smit (44)	a a a a a a a a	· Francis and a
	·	ter et de later en ser en se		с. С. , у
		· ·		
			:	· (• · · ·)
,		(and the quite of
		111		:

Contraction of the second second

1.0

60*#1 28

3.6#10-4

7,0#10-3

		**************************************		Table 4			
		т	otel Body Dosi	metric and Physicl	ogic Dete		
Nuclide	Compartment Deposition Fraction	Compartment / Removel Rate Constant	GL Trect to Blood Transfer	Fraction Excreted in Utine	Decay Constant	Significant Progeny	Branching Ratio
A. Z ¹	×i	Kill	' f _l	fu	ک d-۱	AX .	
137 55Ce	0.13 0.87	0.50 0.0051	1.0	0.90	6.3=10 ⁻⁵	137 m 56 Ba	0.945
65 30 ^{Zn}	0.25 0.75	0.058 0.0022	0.35	0.25	2.8#10-3	65*Cu 29	0.49
90 38 ^{Sr}	0.89	0.21 7.1±10-4	0.20	0.85	6.5x10 ⁻⁵	90 39 90*	1.0
	0.031	L'ORIO -				40 ²⁸	0,0007

0.70

0.0

0.05

0.1

i.4 0.12 0.013 8.7x10⁻⁶

).5#10⁻⁴

bimodal activity ingestion rate distribution for ⁹⁰Sr in the Rongelap popular tion.

Data for 60 Co and 65 Zn were not sufficient for analysis for the Utirik Atoll residents. Values for K_g observed at Rongelap were assigned to Utirik males and females and body burden histories for population subgroups were reconstructed using Eq. 1 or 2. Figures 7 and 8 illustrate the derived mean adult body burdens for all significant nuclides studied on Rongelap and Utirik. This method provides a best fit of the data shown in Figures 2 through 6, and provides a body burden history during the early years post return at Utirik, a time when body burden measurements were not made. Actual data points are also plotted to demonstrate the fit,

The curves shown for 55 Fe in Figures 7 and 8 were obtained by setting K_E equal to zero. This underestimated the initial body burdens and overestimated future ones. Since 55 Fe contributed less than 1.0% to the total dose equivalent, an arbitrary assignment of K_E based on observed values for the other and clides was not attempted. During 1974, another series of blood samples was obtained from Rongelap and Utirik (Co75). Analysis for 55 Fe has yet to be reported. A recalculation of 55 Fe body burden and its impact on early dose equivalent rates will be conducted when the data is made available. A substantial change in dose equivalent is not to be expected.

Figure 4 and Figure 6 illustrate the observed adult histories of 90 Sr and 137Cs mean urine activity concentrations. Mean values for adult males or all adults were plotted. Measured values for 137Cs body burdens were also shown in Figure 7. A much smoother curve was plotted in Figure 7 and it was determined that the collection and analysis technique for urine samples introduced the additional variations. On the basis of this observation for 137Cs, a smooth body



日本の言語



.

,

- ef 1

608 8 11 1



いますし

Fig. 8 Composite Nuclide Body Burden History For Adults at Utirik Atoll

19

burden curve for ⁹⁰Sr, reconstructed from raw data and Eq. 1, was considered a more accurate history. A detailed presentation of the greater variation in radiochemical analysis of urine versus direct body burden measurements can be found in Mi81.

Figure 9 illustrates the variation exhibited in the body burden of 5 randomly chosen subjects over the 25 year monitoring period. These individual variations may have had a dramatic impact on the mean data. In Figure 2, which illustrates the adult male, edult female, and edult population mean ¹³⁷Cs body burden for the 25 year exposure period, a decrease followed by an increase whe seen during the years 1958 through 1963. Although the Castle BRAVO test initially contaminated Rongelap in March 1954, it had been proposed that the Hardtack Phase I series added to this an amount of contamination equal to that responsible for the Figure 2 body burden pattern (Co63). Figure 9 suggests that most individuals counted in those years had body burdens, which remained the same or declined; however, one individual's burden (#881 M) rose and fell quite differently from the others. Several factors could have contributed to this variation from the mean such as departure and return to the atoll, sickness, the dietary contribution of imported foods, etc. Since the mean values are based on small numbers of persons who were chosen at random, it is conceivable that individuals like 881 M influenced the mean body burdens to a greater degree than recontamination of the inhabited atolls. The impact of the individual body burden pattern on the true mean value is moot since body burdens of all individuals were not monitored consistently throughout their residence intervals except in the few cases exhibited in Figure 9.





Ref. 1. And States of Barry

RESULTS AND DISCUSSION

Daily Activity Ingestion Rates

Daily activity ingestion rates were calculated for dosimetrically significant nuclides post return. An exponential decline was proposed for the ingestion rate within a population subgroup and initial reference values are given in Figures 10 through 14 (June 1, 1957, was assigned as a return date to Rongelap). Figure 10 demonstrates the differences in ingestion of 137 Cs for various population subgroups. This undulating pattern was exhibited by 137 Cs, 90 Sr, and 65 Zn, nuclides for which sufficient data existed for analysis.

Differences in ingestion rates of the stable element at the same geographic location have been shown to occur among members of a population (ICRP 23). Age dependent diet studies for ingestion of Cs for urban Japan have values varying from 11 μ g d⁻¹ for adults to 8.6 μ g d⁻¹ for children, Sr in a western type diet rose from 600 μ g d⁻¹ for infants to 690 μ g d⁻¹ for 5 year olds to 3,600 µg d⁻¹ for 13 year olds and fell to a mean of 1,900 µg d⁻¹ for adults. Zn in the United Kingdom rose from 2 to 40 mg d⁻¹, the higher value of Zn being observed in adult tea drinkers. Fe ingestion in a western type diet has a minimum at age 3 and maxima at ages 1 and 20 years. Co is ingested at a rate of 20 ug d⁻¹ for Japanese adults and half this amount for children. The Marshallese population also exhibits dietary changes as a function of age. The authors of the Marshall Islands Diet and Living Pattern Study (Na80) observed coconut sam being used as a major food supplement for infants, and later in adult life each major source of daily fluid intake. Since coconuts and coconut tree sap provided the major source of ¹³⁷Cs on Bikini Atoll (Le80, Mi80), the shape of Fig. ure 10 was in agreement with the observed diet pattern.

「「なれる」となったが、「





TEN

23

and the second of the second of the second second









TY I COM

2.00

1.8





「「「「「「「「」」」」

A CARLES AND A CARL AND A CARL AND A CARL



Figure 11 shows the individual data calculated for ¹³⁷Cs for all Rongelap residents and is referenced to June 1, 1957. The individual maximum ¹³⁷Cs deily activity ingestion rate was approximately 4 times the population mean value. The standard deviation observed for the adult activity ingestion rate distribution was 41% of the mean value, 39% of the mean value for young adults, 48% for adolescents, 38% for children, and 54% for infants. Adolescents and infants exhibited a broader distribution than adults, while children showed a fractional variation in activity ingestion rate similar to that of adults. Breast feeding versus coconut sap supplements would have contributed to the greater variation observed in infants. Adolescents and young adults were the population subgroups which have been observed to move frequently between atolls. This mobility would lend to greater variations in the daily activity ingestion rates relative to those observed in the more stationary population subgroups.

Figure 12 also exhibited q wave pattern; however, a distinct difference between males and females was indicated. This difference arose from the use of values for K_E listed in Table 3 which were derived from urine data for male and female residents at Rongelap Atoll. Its major impact was on the dose equivalent rate, not on the total dose equivalent; and its effect was to cause the dose equivalent rate for males to rise and decline more rapidly than for females.

Figures 13a and 13b summarize the individual data for 90 Sr for all Rongelap residents and were referenced to June 1, 1957. A bimodal shape was observed for the distributions which contained both sexes, again reflecting the difference in the 90 Sr dietary rate constants. Data from urine bioassay indicated that the observed difference between the male and female values for K_g was not significant. A t-test was performed for consecutive urine measurement data during the 23 year residence interval. The results indicate that because

of urine activity concentration variability, there was a 60% probability that the male value for K_E would be different from the female value by the factor observed. Thus differences in the derived activity ingestion rates and dose equivalents were not significant.

Figure 14 shows a semi-log plot of the ⁶⁵Zn and ¹³⁷Cs activity ingestion rate histories for adults on Rongelap. A curve was drawn between points, and the appearance of an increasing 137 Cs ingestion rate during the 1960's indicated the possibility of another contaminating event. The Hardtack Phase I series was conducted just prior to the observed increase in the curve and fallout from the Cactus, Yellow Wood, and Hickory experiments detonated at Bikini and Enewetak would have reached Rongelap. However, several observations fail to support the conclusion that recontamination was significant. These are as follows: 1) the increase in ¹³⁷Cs ingestion rate was not in conjunction with an increase of ⁶⁵Zn; however, since ⁶⁵Zn is an activation product it may have not been produced in the same proportions. 2) The peak Cs body burden at Utirik occurred nearly three years after the initiating event, Castle BRAVO, while the peak body burden at Rongelap followed six years after the potentially contaminating experime ments of the Hardtack series in 1958. 3) The activity ingestion rate at Utirik demonstrated a continuously declining pattern versus the humped pattern observed at Rongelap. This occurred even though there was an equal external exposure rate history following the Hardtack series as measured by the U.S. Public Health Service on both Rongelap and Utirik (Un59). 4) The peak exposure rate on Rongelap following the Hardtack series was 10,000 times less than the peak exposure rate following BRAVO. These facts suggest that the Hardtack series was non a major factor influencing the Rongelap body burden patterns. Thus it is postulated that body burden variations were caused by travel away from the atol

29

NO STREET

or sickness and other factors. Regardless of the cause of individual differences from the mean, a smooth description of the body burden and activity ingestion rate for the population could be adopted. On this basis a declining continuous uptake model was used.

(1)

i fre i

(4)

Internal Dose Equivalent Rates

The approximate instantaneous dose equivalent rates for the total body were determined from the body burden data illustrated in Figures 7 and 8 and from the following equation

where

- $H \equiv$ the total body dose equivalent rate, mRem y⁻¹,
- I = equilibrium dose equivalent rate to the total body per unit body burden, mRem y^{-1} μci^{-1} ,
- $q \equiv$ instanteous body burden, μ Ci.

The approximate nature of the estimate was due to the assumption that the radioactive atoms were distributed among the body tissues as they would be following constant continuous uptake for periods of time much greater than the means residence time for the total body. In the case of 90Sr, 86% of equilibrium was assumed. These assumptions were not used in the estimate of the total dose equivalent. In addition, since mean adult body burdens were computed, a factor of 1.2 was needed to adjust for differences in body mass relative to a 70 kilo gram adult. Table 5 lists values of I which were determined from information given in ICRP59 and corrected for body mass differences.

A _X z ^X	per unit	mRem y ⁻¹ µCi ⁻¹ "
⁵⁵ Fe 26		2 x 10 ⁰
60 27	ο το Κ ρητη Το Το Το	6×10^2
65 _{Zn} 30		1×10^2
90 <mark>s</mark> r 38		3×10^2
137 55		2×10^2

Table 5

Figure 15 illustrates the relative contribution to the composite dose equivalent rate for each dosimetrically significant internally deposited nuclide. For the average Rongelap adult, the residence interval begins June 1, 1957; however, many adults were reported to have resettled during the next 3 to 6 months (Co80b). The composite dose equivalent rate indicated that a bread maximum of approximately several hundred millirem per year persisted for several hundred days. Most of the dose rate is attributable to the 137Cs component Cesium dominated over the entire post return period and would be of prime concern for populations returning to a contaminated environment years after a fination type initiating event.

Figure 16 illustrates two possibilities for the Utirik dose equivalent rate resulting from the ⁶⁵Zn body burden history during the first three years post-return. The higher body burden resulted from use of the two measured 52n

医小脑静脉的 新来来了了

11,526 9 11





0.44

the second that we are a property and good

.,



いのないではある

の語名の

語言

Developments and the state of the state of anti-



body burden means for adults on Utirik and the observed K_E rate constant from Rongelap. It was observed on Rongelap that .031% of 65 Zn was removed from the diet pathway each day in addition to radioactive decay. Additionally, reduction in dietary radioactivity on Rongelap had been observed for $^{(137)}$ Cs, 90 Sr, and 50 Co to be greater than that predicted by radioactive decay alone. Instantaneous reduction fractions very similar to those at Dirick for the 90 Sr, and 137 Cs nuclides. The lower curve on Figure 16 reflects the dose equivalent, dose equivalent rate, and body burden which would have occurred had radioactive decay alone accounted for the removal of 65 Zn from the Utirik environment. Since additionalismechanisms could be measured for other nuclides at Utirik and for the 65 Zn nuclide on a nearby stoll, the upper curve was chosen as the most likely body burden history for sdults post return to Utirik Atoll.

Figure 17 indicates the Utirik adult mean total body dose equivalent rate for each nuclide. An obvious difference relative to the Rongelap history exists; ⁶⁵Zn not ¹³⁷Cs was the major nuclide contributing to the dose equivalent rate. This was due to the Utirik population returning 3 to 4 months after the initial contaminating event, and the Rongelap population returning after 3 years. The age of the fallout had a dramatic influence on the importance of each nuclide contributing to the internal dose equivalent. In fact ⁶⁰Co and ⁶⁵Zn played major roles during the first 3 years, a time interval that

corresponded to the period during which field whole body counting facilities were being developed at Brookhaven National Laboratory and when medical examinations for people on Utirik Atoll were not done. Additionally, pooled and/or individual radiochemical analysis of urine was not performed during this period. The impact of 65 Zn and 60 Co was such that even if the least conservative rate

34







: ::

And the second second

The second se





1.1

•,•

10

35

- e

constant ($K_E \approx 0$) was used for Zn, the dose equivalent rate for the average adult was in excess of Federal Radiation Council Guidelines for the first 2 years following the return to Utirik.

Internal Dose Equivalents

Disintegrations occurring in the total body of an individual during residence following repatriation were determined by several methods. Equation (3), together with personal body-burden histories and stoll specific K_g rate constants from Table 3, provided an initial estimate of disintegrations between consecutive body burden measurements. The second method used was a log-log plot of the subject's body burden history and an algebraic determination of area between to consecutive measured points. The third method used a linear plot of the subject's body burden history. The area under the curve was cut and weighed and compared to a standard weight of known area. Quality control procedures required that all three methods agree within ±10% before a subject was assigned his or her total body disintegrations during residence post return. In general, the methods compared to within ±5%.

After the total number of disintegrations occurring in a subject's body was assigned, they were apportioned among the body organs according to the following equation

$$F = \frac{E_2' E_i A_i B_i (E_i C_i D_i + \ln 2/\lambda)}{E_i C_i D_i (E_i A_i B_i + \ln 2/\lambda)}, \qquad (5)$$

STREE BOLDER STREET SO VERSON WITH STREET STREET

where

F ≡ the fraction of total body disintegrations occurring in the organ of interest,

 $A_i \equiv organ$ compartment deposition fraction for the element,

 $B_i \equiv \text{organ compartment biological half time for the element,}$ $C_i \equiv \text{total body compartment deposition fraction for the element,}$ $D_i \equiv \text{total body compartment biological half time for the element,}$ $f'_2 \equiv \text{fraction of the element from blood to organ of reference.}$

Equation (5) applied where significant decay occurred at the deposition site, and not during transit or re-transit to the organ of interest. Values for compartment deposition fractions and compartment half times were obtained from Ki78. Values for the remaining quantities were from ICRP59.

The dose equivalents to a specific organ or the total body were determined by using the source to target dose equivalent per unit cumulated activity parameters from Ki78. The total target dose equivalent was obtained by summation of the dosimetric contributions from all source organs. Several important modifications to the general procedure were made in order to compute individual dosimetric results. For each person, the source to target dose equivalent per unit cumulated activity was weighted by the ratio of a standard man's body mass relative to the actual mean body mass during the interval for which the dose equivalent was determined. In the case of ¹³⁷Cs, the long term biological removal rate constant for the Marshallese population was highly dependent upon body mass (Mi81). Appropriate modifications to Eq. (2), (3), and (5) were made to reflect this dependence. Finally, for ⁹⁰Sr deposition in bone, 28% of the source to target dose equivalent per unit cumulated activity was assumed from cancellous bone and 72% from cortical bone.

Figure 18 demonstrates the mean dose equivalent from ¹³⁷Cs for various age and sex groupings. The residence interval was from 1957 to 1980 for this population. The adolescents and persons above 50 years of age in 1957 maintained the lowest dose equivalent. Persons who died during this period were not included

: 8 110 11 1.11 6-mi (10 100 1 11 1 6 111 smi 1 frai \overline{a} (Lassi 6.200 ad 217 11 EMALES 11 MEAN DOSE EQUIVALENT, rem 100 MALES Ū. ğ





Contraction of the

38

+1444Y2

and the second second

and the second se

in the figure nor were they included in any dosimetric distributions for any of the nuclides. Thus all persons considered, regardless of initial age in 1957, experienced a 23 year exposure interval.

Figure 19 shows dose equivalent distributions according to age and the for ¹³⁷Cs among the Rongelapese. The shape or the population distribution was skewed with a mean of 1.7 Rem and a maximum of 9.0 Rem. Thus the maximum was 5.3 times the mean value for ¹³⁷Cs on Rongelap. An examination of the subgroup distributions reveals that persons who were infants at the time of rehabilition at Rongelap also were the recipients of the higher doses. This was due to the combined effects of lower average body mass, a higher average ingestion rate, and more rapid turnover of ¹³⁷Cs than that for adults or even children. The parameter having the greatest impact on the infant dose equivalent was body mass. The standard deviation for the adult male distribution was 49% of the mean dose equivalent, for adult females 43% of the mean dose equivalent, and for adules-cents 47%. Within a subgroup, the maximum observed dose equivalent was approximately twice the mean value for all distributions considered here.

Figure 20 shows mean dose equivalents as a function of returning age groups for 65 Zn on Rongelap. Adolescents, young adults, and adults 50 and up were the groups receiving lower total dose equivalents, while children and middle aged persons received higher dose equivalents during the residence interval. Measured 65 Zn data for persons who were infants at the return date were not reported in the publications by Conard et al. Figure 21 shows the dosimetric distributions observed for members α (the Rongelap population for ⁶⁵Zn. Again the population overall exhibited a sidewed distribution of dose with a maximum value nearly three times the mean. (h) dren demonstrated higher doses than persons who were adults during the entire 23

the tigure nor wave they in a decis are in any sontime to a short of the second sont

5.50

DIBROW DEPARTORMENT STATE SHELL BADITOR V

16 . 3.1.

いたので、「「「「「「」」

こうちょう していたいない いちょう していたい シート



(H) Infants on Rongelap



and a series



41



and the set of the second

and the second

year period. The standard deviation was in general 30% of the mean value for all age and sex subgroup distributions. This less pronounced variation may be due to the fact that 65 Zn measurements took place over a 3 year interval while 90 Sr and 137 Cs occurred over a 23 year interval and thus was contained in a more homogeneous population than were the longer lived nuclides.

Figures 22 and 23a and 23b summarize the ⁹⁰Sr dose equivalent results for individuals at Rongelap.

In this analysis, only the ingestion pathway was considered important. Some radioactivity would enter the body via the resuspension and direct inhalation pathways. It is known that for a given soil concentration of the stable naturally occurring analogs to the radionuclides considered here, the ration of food and fluid intake to blood relative to airborne intake to blood, are as follows:

Co	>	3000	Zn.>	>	130
Fe	>	550	Sr >	>	10,000
Cs	>	400			

Thus, dietary intake of radioactive material is the principal pathway leading to internal deposition. This applies to most nuclides in the environment, however, there are notable exceptions including I. U. and Pu. 「「「「「「「」」」」

and the second se

External Exposure

A value of .73 rads in tissue of interest per röntgen, measured in all at one meter above the surface, was used to convert exposure in air to absorbed dose in tissue. The source was assumed to be an exponential distribution of 137 activity with depth in soil, typical of aged fallout (Be70). Because of the multidirectional nature of the source, variation of absorbed dose with depth of organ was minimal. Additionally, external doses were adjusted for living pate





and the state of the



1.11.111

tern variations since the stolls present a heterogeneous exposure rate environment (Gr77).

External exposure calculations are based on Figures 24 to 26 which were derived from data listed in Cr56, Sh57, Un59, and Gr77. The area under straight line portions of the curve was determined by

$$\mathbf{x} = \frac{\mathbf{R}_2 \mathbf{t}_2 - \mathbf{R}_1 \mathbf{t}_1}{n+1} , \qquad (6)$$

where

X \equiv external exposure during straight line interval, mR, R₂ \equiv exposure rate at the end of the interval, mRh⁻¹, R₁ \equiv exposure rate at the beginning of the interval, mRh⁻¹, t₂ \equiv time post detonation at the end of interval, hours, t₁ \equiv time post detonation at the beginning of interval, hours, n \equiv slope of a straight line.

Data from 11 detonations during May, June, and July of 1958 (Sh57) indicated a mean fallout deposition exponent of 18.8. This mean value was observed at Utirik, Rongelap, Parry, and Wotho and was applied to early time post detonation of BRAVO to obtain the initial increasing exposure rate history shown on Figures 24 and 26. This method yielded a fallout deposition period of 5.5 hours on Rongelap and 12 hours on Utirik. This time compares well with the original observations reported by the Marshallese and by U.S. Navy personnel stationed in the area (Sh57). Initial dose equivalents on "acute doses" are developed in greater detail in another report.





ッチ イズモ

Fig 25 Rongelap External Exposure Rate History Post Cactus

I BELLE ADDRESS A MURILIM MET TOMA HE ELECTION AND ADDR

.

The rear and



. 2 }

13

đ

51

ţ



Fig. 26 Utirik External Exposure Rate History Post Bravo

Figure 25 demonstrates the external exposure following the 1958 testing are ries. Since raturn to Rongelap followed 3 years after the BRAVO contamination, this series contributed in large part to the external exposure post return.

SUMMARY

The Castle BRAVO shot of March 1954 caused the contamination of the inhabited atolls Rongelap and Utirik. Evacuation from Rongelap commenced 50 hours after detonation and from Utirik 55 hours after detonation. During June 1954 and June 1957 the return of the Utirikese and Rongelapese occurred respectively. Body burden data for dosimetrically significant nuclides were obtained throughout the residence interval post return primarily by direct in vivo gamma spectroscopy and by indirect radiochemical analysis of urine and blood.

The dosimetric models used in this analysis were representative of a declining continuous uptake regime. Dietary decline of radioactivity included radioactive decay of the source and a conglomerate of other factors which might have included increased use of imported foods and weathering of the source. Dietary loss rate constants were estimated from sequential body burden data and were comparable for both atolls. Variation in body burden history data for a particular nuclide on a particuular atoll was observed in whole body counting data and urine bioassay results. This was attributed principally to the statistical variation encountered when small groups are sampled from a heterogeneous group of body burdens in people, and in the case of urine bioassay additional variation was introduced during the laboratory analysis of samples.

Daily activity ingestion rates were determined for all measured radionuclides. In general, infants, children, and adults between 20 and 40

50

- EMPLOY - AMPLITY - PROVIDED AND A STATE

years of age ingested more activity each day than did adolescents and persons greater than 40 years of age. Maximum deviation from the average value of the daily activity ingestion rate for members of an age subgroup was no greater than a factor of 3. However, the population distributions illustrated a maximum factor of 5 times the mean activity ingestion rate value.

Dose equivalent rates post return were determined for members from both atolls. For Rongelap Atoll, the residents received approximately 100 to 200 mRem per year during the first 5000 days post return from internal emitters. The principal contributing nuclide was ¹³⁷Cs. For Utirik Atoll, the remidents received up to 15 Rem per year during the first 400 days post return. The major contributing nuclides were ⁶⁵Zn and ⁶⁰Co. Dose equivalentizates to the Utirikese from internal emitters fell below 500 mRem per year at approximately 1200 days post return.

The dose equivalent for population subgroups and for individuals was determined. Table 6 summarizes the results for the total body, thyroid, red marrow, testes, ovaries, lower large intestine wall, and liver. The catenary compartment model of Bernard and Hayes (Ber70) was used to determine doses to various segments of the gastrointestinal tract. The Utirikese received significantly more radiation dose from 65 Zn, 60 Co, and 55 Fe than did the Rongelapese because of short mean residence times of these nuclides in the environment. 90 Sr doses to the Rongelapese were 2.5 time greater and 137 Cs doses 1.5 times greater than doses received by persons at Utirik. This occurred even though Utirik residence to which Utirik was less contaminated than Rongelap.

- <u> </u>	bib	.14	:11	<u>.</u> 1	, , ((1)	1	1	ņ	;	<u>}</u> .	5		:
				Tal	le	6		-				_		
	mos;					į D			• •					

Chronic	Phase
Dose Equivalen	t Summary, Ren

•

• • • •

	T	otal Body	Thyroid			
Nuclide	Utirik Adults	Rongela Adults	p Utirik Adults	-Rongelap Adults		
90	010	10-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-	00075	0017		
508r	.012		.00073	.0017		
137	.033	.023	.054			
600-	1 + 1 K 1	1.1	1.0	2.4		
65~	12	+014	• 30	•010		
" 'Zn	13.	.0/0	11.	.00/		
Incernal	14.	1.9	13.	2.5		
External Total	3.2 17.	2.0	3.2	4.5		
	Red Marrow	•	<u>Testes-Ova</u>	ries		
9037	. 054	.12	.0007500075	.0017+.0017		
20 Fe	.060	.042	.058062	.074043		
137:s	1.7	2.6	1.5-1.7	2.3-2.6		
0 70	.63	.018	.44-1.8	0.12050		
0 - · n	17.	.10	1116.	.069099		
In: nal	20.	2,9	1320.	2.5-2.8		
External	3.2	2.0	3.2	2.0		
Total	23.	4.9	1723.	4.5-4.8		
	Low	er Large				
	Inter	stine Wall		Liver		
90 _{Sr}	. 23	. 57	.00067	.0015		
55Fe	.067	,047	.12	.080		
13/ _{Cs}	. 59	.90	1.8	2.7		
60 _{Co}	4.7	.13	.79	.022		
65 _{Zn}	15.	.091	17.	.14		
Internal	21.	1.7	19.	3.0		
External	3.2	2.0	3.2	2.0		
Total	24.	3.8	22.	5.0		

1810 N. 194

しょうしょう **(1989) 1**1 (1997) たいしょう きわられた (1997) 日本 (1997) たいしょう かんしょう かんしょう しんしょう たちゃく

Be70	Bennett, B.G., 1970, Estimation of gonadal absorbed dose due to environ
	mental gamma radiation, Health Phys. 19, 757-67.
Be72	Beasley, T.M., Held, E.E., and Conard, R.A., 1972, Iron-55 in Rongelap
	people, fish and soils, Health Phys. 22, 245-50.
Ber70	Bernard, R.S. and Hayes, R.L., 1970, Dose to various segments of the
	gastrointestional tract, Pro. Symp. Med. Radionuclides: Radiation Dode
	and Effects.
Co57	Conard, R.A., et al., 1958, March 1957 Medical Survey of Rongelap and
	Utirik People Three Years After Exposure to Radioactive Fallout, BNG
	501.
Co59	Conard, R.A., et al., 1959, Medical Survey of Rongelap People, March
	1958, Four Years After Exposure to Fallout, BNL 534.
Co60	Conard, R.A., et al., 1960, Medical Survey of Rongelap People Five and
	Six Years After Exposure to Fallout (With an Addendum on Vegetation),
	BNL 609.
Co62	Conard, R.A., et al., 1962, Medical Survey of Rongelap People Seven
	Years After Exposure to Fallout, BNL 727.
Co63	Conard, R.A., et al., 1963, Medical Survey of Rongelap People Eight
	Years After Exposure to Fallout, 3NL 780.
Co 65	Conard, R.A., et al., 1965, Medical Survey of the People of Rongelap
	and Utirik Islands Nine and Ten Years After Exposure to Fallout Radia-
	tion (March 1963 and March 1964), BNL 908.

i,

53

ួត ម

• • •

1.14 . .

· •

. ۰.

ر ال 1.14

, -, -, -

Co67 Conard, R.A., et al., 1967, <u>Medical Survey of the People of Rongelep</u> and Utirik Islands Eleven and Twelve Years After Exposure to Fallout Radiation (March 1965 and March 1966), BNL 50029.

Co70 Conard, R.A., et al., 1970, <u>Medical Survey of the People of Rongelap</u> and Utirik Islands Thirteen, Fourteen and Fifteen Years After Exposure to Fallout Radiation (March 1967, March 1968, and March 1969), BNL 50220.

- Co75 Conard, R.A., et al., 1975, <u>A Twenty-Year Review of Medical Findings in</u> <u>a Marshallese Populations Accidentally Exposed to Radioactive Fallout</u>, BNL 50424.
- Co80a Conard, R.A., et al., 1980, <u>A Twenty-Five Year Review of Medical Find-</u> ings in a Marshallese Population Accidentally Exposed to Radioactive <u>Fallout</u>, BNL, in press.

Co80b Conard, R.A., private communication.

- Cr56 Cronkite, E.P., Bond, V.P., and Dunham, C.L., 1952, <u>A Report on the</u> <u>Marshallese and Americans Accidentally Exposed to Radiation from Fall</u> <u>out and a Discussion of Radiation Injury in the Human Being</u>, U.S. Atomic Energy Commission, TID 5358.
- Gi72 Glasstone, S., Editor, 1957, <u>The Effects of Nuclear Weapons</u>, Defense Atomic Support Agency, Department of Defense.
- Gr77 Greenhouse, N.A., and Miltenberger, R.M., 1977, <u>External Radiation Sur-</u> vey and Dose Predictions for Rongelap, Utirik, Rongerik, Ailuk and Wotje Atolls, BNL 50797.
- ICRP59 Recommendations of the International Commission on Radiological Protection, 1959, ICRP Publication 2, Report of Committee II on Permissible Dose for Internal Radiation.

54

- ICRP68: Recommendationsofther International Commission on Madiological Protection, 1968, ICRP. Publication: 10, Report of Committee IV on Evaluation of Radiation Doses to Body Tissues from Internal Contamination due to Occupational Exposure. Softher of the Advances of Advances
- ICRP69 Recommendations of the International Commission on Radiological Protection, 1969, ICRP. Publication 10A, Report of Committee (IV on the Assessment of Internal, Contamination Resulting from Recurrent or Prolonged Uptakes.
- ICRP74 Recommendations of the International Commission on Radiological Protection, 1974, ICRP Rublication 23, Report of the Task Group on Reference Man.
- ICRP79 Recommendations of the International Commission on Radiological Protection, 1979, ICRP Publication 30, Report of Committee II on Limits for Intakes of Radionuclides by Workers.
- Ja64 James, R.A., 1964, Estimate of Radiation Dose to Thyroids of the Rongelap Children Following the BRAVO Event, UCRL-12273.
- Ki75 Killough, G.G., Dunning, D.E., Bernard, S.R., and Pleasant, J.C., 1978, Estimates of Internal Dose Equivalent to 22 Target Organs for Radionuclides Occurring in Routine Releases from Nuclear Full-Cycle Facilities, NUREG/CR-0150, ORNL/NUREG/TM190.

- Le80 Lessard, E.T., Miltenberger, R.P., and Greenhouse, N.A., 1980, Dictary Radioactivity Intake from Bioassay Data: A Model Applied to ^{13/}0s Intake by Bikini Island Residents, Health Phys. 39, 177-83.
- Mi80 Miltenberger, R.P., Greenhouse, N.A. and Lessard, E.T., 1980, Whole Body Counting Results from 1974 to 1979 for Bikini Islands Residents, Health Phys. 39, 395-407.

55

14,

- Mi81 Miltenberger, R.P. golessard, E.T., and Greenhouse, N.A., 1981, <u>60 Co and</u> 137 Cs Long Term Biological Removal Rate Constants for the Marshallese Population, Health Phys. 39, in press.
- Na80 Naidu, J., Greenhouse, N., and Knight, J., 1980, <u>Marshall Islands: A</u> Study of Diet and Living Patterns.
- Sh57 Sharp, R., and Chapman, W., 1957, Exposure of Marshall Islanders and American Military Personnel to Fallout, Naval Medical Research Institute, WT-938.
- Un59 United States Public Health Service, 1959, <u>Report of Public Health Serv</u> vice Off-Site Radiological Monitoring Data, Operation Hardtack Phase 1 1958.

1.8.40

1. 1. 1.

HTT:

111 18 18433A