

~~CONFIDENTIAL~~

R

A

152

John C. Jugher, M. D., Director
Division of Biology and Medicine

May 4, 1954

Gordon M. Dunning, Health Physicist
Biophysics Branch, Division of Biology and Medicine

DATA ON SOIL SAMPLES COLLECTED ON THE ISLANDS OF THE PACIFIC FOLLOWING THE FIRST DETONATION AND ALSO THE DOSE RATE READINGS AT THE SAME LOCALITIES.

NO COPY AVAILABLE

Attached (Annex I) are data on soil samples collected on the islands of the Pacific following the first detonation and also the dose rate readings at the same localities. One method of evaluating such data is to try to establish relationships between different units if possible. One useful relationship would be the conversion of disintegrations per minute per gram of soil to milliroentgens per hour of gamma radiation at a three-foot height or vice versa.

Larson's work with soils around NPG during the spring 1953 tests indicated the following relationship: 10 $\mu\text{c}/\text{sq. ft.}$ beta counts of soil (after absorption and geometry corrections) \rightarrow 1 mr/hr gamma at 3 feet. We found that essentially all of the activity was in the first one inch of top soil.

In collecting soil on the Pacific Islands good care was taken to collect one square foot of surface (in fact, templates were made for this purpose). It was impossible, however, to scoop up the soil to a uniform depth so the rule followed was to collect to one inch or greater. If the fallout activity in the Pacific Islands also was contained in the first one inch, the additional soil below this contributed mass but little activity to the sample. By taking the disintegrations per minute per gram (after a thorough mixing) and multiplying by the total number of grams for each sample one should arrive at the activity per square foot.

The plot of beta disintegrations per minute per gram of soil versus mr/hr of gamma at 3 feet is shown on the attached graph. The correlation is not too good. For references, several curves are arbitrarily drawn on the graph. The data strongly suggest that less than 10 $\mu\text{c}/\text{sq. ft.}$ is equivalent to 1 mr/hr . For lower levels of activity the data are more of the order of 2 $\mu\text{c}/\text{sq. ft.}$ \rightarrow 1 mr/hr and for higher levels of activity less than 1 $\mu\text{c}/\text{sq. ft.}$ \rightarrow 1 mr/hr . If ~~one is~~ in trying to fit a curve to such widely scattered data, ~~then~~ $\mu\text{c}/\text{sq. ft.}$ \rightarrow 1 mr/hr is a very rough approximation. Another possibility but not probability is a more complex curve shown ~~by~~ the solid black line.

10005
CLASSIFICATION CANCELLED
OR CHANGED TO
BY AUTHORITY OF DOE/DPC
L. B. [Signature] 2/1/85
REVIEWED BY DATE

~~CONFIDENTIAL~~

OFFICE \rightarrow BMBP
SURNAME \rightarrow Dunning:MMack.
DATE \rightarrow 5/4/54

MILITARY RESEARCH & APPL.

~~CONFIDENTIAL~~
~~CONFIDENTIAL~~
~~CONFIDENTIAL~~
~~CONFIDENTIAL~~

Dr. John C. Dugher

-2-

May 4, 1954

Based on calculations made in Annex II it would appear that about $10 \mu\text{c}/\text{ft}^2 \rightarrow 1 \text{ mr/hr gamma at 3 feet}$. If one accepts the figure of 2 beta emissions for each gamma photon (Effects of Atomic Weapons) then $20 \mu\text{c}/\text{ft}^2 \text{ (beta)} \rightarrow 1 \text{ mr/hr gamma at 3 feet}$.

Knowing the difficulties of collecting, handling, packaging, shipping and counting the samples, it is probable that some of the data are not entirely valid. The information shown in this memo suggests that further carefully controlled studies must be made before a more firm conclusion may be reached.

Attachments 3

Annex I

Annex II

Graph (w/cy 1A only)

DISTRIBUTION: copy 1A - Dr. Dugher
2A - Dr. Dunning
3A - Bioph. Dr. Reading File
4,5,6A - M&M Files

~~CONFIDENTIAL~~
~~CONFIDENTIAL~~
~~CONFIDENTIAL~~
~~CONFIDENTIAL~~
~~CONFIDENTIAL~~
~~CONFIDENTIAL~~

OFFICE ▶

5001072

~~CONFIDENTIAL~~

Dr. Dugher

May 4, 1954

ANNEX I

Comparison of Soil Activity to dose-rate readings

<u>Location</u>	<u>Date of Collection</u>	<u>Yo. Trees</u>	<u>d/n/gm (same date)</u>	<u>Total Act. (mc/24)</u>	<u>mr/hr (survey meters)</u>
Likiep	6 March	2,400	23,000	2.5×10^1	~3
Jeno	"	1,060	13,000	0.2	~3
Ailuk	"	2,160	23,000	2.2×10^1	3 - 15
Mejuit	7 March	1,360	30,000	1.3×10^1	3 - 10
Ormed	5 March	1,325	15,000	9.0	3.5
Erikub	5 March	1,720	4,300	3.3	1.5
Kaven	6 March	1,335	5,500	3.3	1.8
Wotho	6 March	1,490	2,400	1.6	~0.8
Balap	7 March	765	950	0.4	0.5
Rongelap (Northern)	8 March	703	890,000	9.2×10^2	440.0
(Central)	"	815	1,600,000	5.7×10^2	280.0
(1 mile N. Village)	"	1,680	100,000	7.6×10^1	340.0
(South Western)	"	1,060	140,000	6.6×10^1	220.0
Eriirippa	"	810	9,000,000	3.2×10^3	2,200.0
Eniwetok	"	2,010	780,000	7.1×10^2	900.0
Kabelle	"	1,470	4,500,000	3.0×10^3	2,000.0
Utirik	9 March	1,140	1,100,000	6.9×10^2	40.0
Bikar	"	1,080	85,000	1.1×10^1	140.0
Eniwetok	10 March	1,050	185,000	8.8×10^1	280.0
Bifo	"	1,060	14,000	0.7	100.0

5001073

ANNEX II

Calculations of Dose Rate at Three Feet Above A Plane Surface

For point source (0.3 - 3.0 Mev range)

$$\text{Dose rate (r/hr)} = \frac{6CE}{d^2}$$

where: C = activity (curies)
 E = energy in Mev
 d = distance in feet

$$\text{Dose rate} = \frac{6EA \int_0^R \frac{dx}{h^2 + x^2}}$$

where: A = activity/unit area
 h = height above surface (feet)
 x = distance in feet

$$\begin{aligned} \text{Dose rate} &= 6EA \int_0^{R_2} \frac{dx}{x^2 + h^2} \\ &= 6EA \ln \left[\frac{h^2 + R_2^2}{h^2 + R_1^2} \right] \end{aligned}$$

The mean free path for 0.7 Mev in air is about 360 feet so that essentially all of the dose will be contributed from a surface 1,000 feet in radius.

Let A = 10×10^6 curies/ft²
 E = 0.7 Mev

$$\begin{aligned} \text{Dose rate} &= 6\pi(0.7)(10 \times 10^6) \ln \left[\frac{2 + 10^6}{1} \right] \\ &= 7.2 \text{ mr/hr} \end{aligned}$$

Since this formula assumes no absorption and also a uniform plane surface, an estimate is that

10 $\mu\text{c}/\text{ft}^2$ \approx 7.2 mr/hr.

500 1074