405163

Walter J. Williams, Deputy General Manager

December 14, 1953

John C. Bugher, M.D., Director, Division of Biology and Medicine

MONTHLY STATUS AND PROGRESS REPORT, NOVEMBER 1953 -DIVISION OF BICLOGY AND MEDICINE

SYMBOL: BMA BON

Transmitted herewith is the Monthly Status and I Division covering the month of Hovember 1953.	rogress Report for this			
US DOE ARCHIVE 326 U.S. ATOMIC ENT COMMISSION				
CC: J. H. Burchard	RG DOE HISTORIAN (DBM)			
When separated from enclosure, handle this docum				
as	Box <u>3363</u> Folder <u>#23</u>			

## **BEST COPY AVAILABLE**

This material contains information affect meaning of the \*\* U.S.C., Secs. 793 and 79 mer to an unauthorized pe ated by law.



DOE 5650.2 THE-12 CLASSIFICATION CANCELLED BY AUTHORITY OF DOE/OC

By: Dick KOOGLE 6-10-87

OFFICE >	Adm.	X-0	Director		
SURNAME >	O'Neill:emr	ON 45	9cB	ORGANIZATION & MAI	
DATE >	Kon	70	12/16		- Automate
	2-11				 

	7710III	DIOTED
	=	P

## Collection <u>1132</u>

US DOE ARCHIVES

326 U.S. ATOMIC ENERGY COMMISSION

RG DOE HISTORIAN (DBM)


# NATURAL STATUS AND HOGHESS REPORT

Box 3363

Folder <u># 23</u>

and mutter to

DOS ARCHIVES




The National Radiation Protection Committee recommends a maximum permissible concentration of radium water for continuous use of  $1.\times10^{-3}$  microcuries per cubic centimeter. This is equivalent to  $1.\times10^{-11}$  curies per milliliter or curies per gram. Approximately the same value measured in curies per gram would apply to the total quantity of food consumed. Thus, it is indicated from these studies that tap water and the common foods used in the United States appear to be well within the permissible limits for radium.


## Rediction Effects on Tobacco and Poteto Plants. (UNCLASSIFIED)

A species of tobacco was planted this year in the games field at Brookhavan National Laboratory. Results show that irradiation produced a moderately high rate of spontaneous tumors. In the games field, the tumor-induction rate at fairly high desagn rates (300 r/dsy) was greatly increased over the control lot, not only in numbers of tumors per plant but in size of the individual tumors. This is the first report of a tumor in plants induced by games radiation, although radiation induced tumors have been recognized in animals for many years.

Experiments on the effect of radiation on the "essping" qualities of potatoes were completed by Brookhaven Mational Laboratory recently. A cusntity of potatoes was secured and divided into five lots, four of which were given various radiation doses and the fifth kept as a control. The potatoes later were all stored under conditions compareble to standard potato storage conditions, and semples were taken periodically from each lot and tested for taste and then graded according to standard potatograding procedures. At the final test period on August 15, the control lot was essentially spoiled, whereas those that had been exposed to 20,000 r were still Grade A potatoes in all respects. Fotatoes exposed to about 50,000 r might be classed as Grade B potatoes, and those exposed at 80,000 r were also Grads B potatoes. What apparently happens is that, in late spring, potatoes tend to sprout and these eprouts release an engyme responsible for the breakdown of the starch. Sprouts are inhibited by relatively small doses of rediation, and consequently the starch is not broken down. At the higher doses, apparently, there is enough cellular destruction to cause a general breakdown over a long period of time. Whether these explanations are correct or not is debatable, but there seems to be little doubt now that this is a real phonomenon, since it was observed last year quite accidentally in connection with some other work, and the experiment this year was designed specifically to grove this point.

Typing of Blood Platelets. (UNCLASSIFIED) Discovery of groups and types for blood platelets was announced recently by the New England Hedical Center at Boston, Massachusetts. Platelets are colorless, disk-shaped bodies found in the blood of humans and all other massals. They






play a part in the blood-clotting mechanism, but are still not well understood. Ionizing radiations to the entire body produce smong other effects a reduction in the number of blood platelets from; in some instances, actual total consistence, in this connection, it is interesting that animals have been protected from lothal effects through transfusion of platelets from the blood.

The platelets can new be typed and grouped, much as red blood cells are before normal blood transfusions. There is no correlation, however, between the platelet groups discovered and the familiar groups of red blood cells. Continued efforts may prove of significant value in combatting such maladies as thrombocytopenic purpura, a blood disease in which bleeding occurs under the skin and internally. In adults suffering from this type of homorrhagic disease, the gradual failure of repeated platelet transfusions may be due to incompatibility of the donor's platelets with those of the patient. A similar consition also occurs in newborn babies and this may be due to platelet incompatibility between mother and newborn child, in a way similar to the incompatability.

Effects of Prenatal X-irradiation on Mice. ( In studies of the diffect of radiation on embryos of marmais and other organisms, it is necessary to observe changes induced in one or more variable organs or organ systems. In research in progress at Oak Ridge National Laboratory, the vertebral column and thorax have been used as indicators of radiation damage to the developing mouse. The particular question being investigated is whether radiations cause general damage to the embryo at any exposure time or whether radiations cause specific changes at specific radiation susceptible times. Mice embryos were irradiated at stages of development corresponding to ages of 1/2 day, 1-1/2 days, 2-1/2 days, and so on to 13-1/2 days after conception. At birth the excleton of each embryo was studied in detail. It was found that each skeletal abnormality was, in general, induced by radiation during only a short period of prenatal development and was obtained with high incidence when radiation was applied in that period. The "critical periods" for the characteristics studied fell between 6-1/2 and 13-1/2 days after conception, during which time irradiation produces very little mortality. The abnormalities produced by irradiation resemble, but do not exactly coincide to, some of the abnormalities produced by previously identified mutent genes in the mouse.

## Civil Defense

Dissemination of Weapons Test Information. A recent meeting was held with members of AEC, the Department of Defense, and the Federal Civil Defense Administration. The discussions covered the transmission to FCDA of classified weapons effects information derived from



tests and associated projects under terms of the Joint Chiefs of Staff Directive to the Armed Forces Special Weapons Project controlling such dissemination. The following decisions were reached:

(1) AFSWP agreed to make available height of burst curves and new curves on neutron flux, previously withheld under the JCS directive, and to translate into more readily useable form transmittable data;

(2) the FCDA representatives concurred that no change in the JCS directive would be pressed, since it generally passitted the forwarding of most, if not all, of the data needed for civil defense planning, and that any problem arising in this connection in the future could be discussed on a case-to-case basis.

Puture Tost Programs. (URCLASSIFIED) Heads of the Federal Civil Defense Administration for participation in possible future continental weapons test programs were outlined in a recent informal meeting with FCDA Administrator and the Chairman of the Commission and staff members. AEC willingness to cooperate in all ways possible was expressed by the Commission Chairman. It was agreed that specific FCDA proposals would be submitted for consideration at a later date.

## Technical Assistance Rendered. (UNCLASSIFIED)

- a. At the request of the Federal Civil Defense Administration, design criteria for protective construction standards to be publicly released as guidance to federal agencies and the public in erecting structures in or near designated target areas have been reviewed by the Cosmission. FCDA has incorporated such of the ABC material in its proposed publication "Standards and Criteria for Protective Construction Class I Buildings" which is now undergoing classification review.
- b. The problem of providing shelters in Navy Department Buress of Ships installations at Navy Yards was discussed with the Chief Design Engineer, Public Works, Buresu of Ships. Based on discussion of design criteria, materials, and probability of accomplishing objectives of the Navy, further study will be conducted by Buresu of Ships considering the construction of dual-purpose reinforced concrete buildings to provide the shelter required.

RG DOE HISTORIAN (DBM)
Collection
Box3343
Folder #23

