

RECLASSIFICATION CANCELLED

US DOE ARCHIVES
326 US ATOMIC ENERGY
REG. COMMISSION

BY None DATE 3/26/81

NON-CCFP BY AUTHORITY OF AEC 5650.2, III-12 (ADMIN)
BY Milford J. Dunning DATE 3-26-81

Dr. John C. Bugher, M.D., Director
Division of Biology and Medicine

May 13, 1954

Gordon M. Dunning, Health Physicist
Biophysics Branch, Division of Biology and Medicine

402709

ESTIMATED RADIATION DOSE TO THYROID OF NATIVES FROM RONGELAP

Collection Dead Files
326-78-3 Box 3
MRA 1-S Radiation Exposures
Native and Test Persons
(March-May 1954)

This memorandum is in reply to your request for an estimate of additional doses to the thyroid of the Rongelap natives due to the fact that tellurium, as a precursor to iodine, may be present in the gut after ingestion of fallout material. The tellurium, in turn, might disintegrate into radioactive iodine while in the gut, with subsequent deposition of the iodine in the thyroid.

There are some 17 radioactive isotopes of tellurium but only 7 of these are produced in fission. Of these, 6 are not of interest (4 have too short a half-life, 1 leads to stable iodine-127 and 1 leads to iodine-129 with a half-life of 2.4×10^7 years). The remaining radioisotope is tellurium-132, with a half-life of 77 hours leading to iodine-132 with a half-life of 2.4 hours. (Incidentally there is no tellium precursor that is of interest here.)

Without having the original data of LASL I have accepted their estimate that there were ingested and/or inhaled the products of 5×10^{13} fissions, assumed they were all ingested, and then proceeded to calculate the dose to the thyroid from (a) I^{131} (b) each short-lived iodine isotope of interest and (c) the added dose coming from $T^{132}-I^{132}$. The calculations show that $T^{132}-I^{132}$ will produce an added dosage of about 26%.

The best estimated percentage absorption and deposition of iodine is yet to be determined. The best estimate I can turn up to date is still the 20% quoted in NBS Handbook 52. However, I will continue to search for additional information. In the meantime the table below indicates the magnitude of doses to the thyroid if one assumes 20, 50, and 100% absorption and deposition. Incidentally, it may be noted that the calculations below based on 20% (the number assumed by LASL) show estimated doses to the thyroid from I^{131} and from shorter lived iodine isotopes to be in good agreement with those estimated by LASL, i.e., 50 reps for I^{131} and 80 reps for short-lived iodine isotopes.

BEST COPY AVAILABLE

OFFICE	BHBP			MILITARY RESEARCH & APPL	7-1
SURNAME	DUNNING: Mack	<i>YMD</i>			
DATE	5-13-54				

Form AEC-318

U. S. GOVERNMENT PRINTING OFFICE 16-6910-2

Dr. John C. Bugher

- 2 -

May 13, 1954

DOSE TO THYROID (REPS)

	<u>Assuming 100% Retention</u>	<u>Assuming 50% Retention</u>	<u>Assuming 20% Retention</u>
I ¹³¹	255	255	128
I ¹³²	27*	(54)**	14*
I ¹³³	370	370	185
I ¹³⁵	60	60	30
I ¹³² (Te ¹³²)	185*	(370)**	93*
Total	897*	(1109)**	450*
			(555)**
			169*
			(221)**

* If assume that one-half of the I¹³² (half-life 2.4 hours) present in the gut is deposited in the thyroid.

** If assume all of the I¹³² (half-life 2.4 hours) present in the gut is deposited in thyroid.

Most probable estimate of ratio of doses to thyroid is:

$$\frac{I^{132}, 133, 135 + I^{132}(Te^{132})}{I^{131}} \approx 2.5$$

This document contains information affecting the national defense of the United States within the meaning of Title 10, U.S.C., and is lawfully made available to an authorized person

OFFICE ►					
SURNAME ►					
DATE ►					

Mr. Bugher

- 3 -

May 13, 1954

ANNEX

Calculations of Dose to Thyroid

I^{131}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At D \neq 1 there are 0.017 d/ μ /10 $\times 10^{13}$ fissions

or 8.5×10^7 d/ μ /5 $\times 10^{13}$ fissions

or 38.3μ c intake of I^{131}

or 1.4×10^{12} atoms intake of I^{131}

(Average energy 0.22 Mev)

$$\text{Dose (reps)} = \frac{(1.35 \times 10^{12})^*(0.22)(1.6 \times 10^{-6})}{(20)(93)} = 255 \text{ reps}$$

* Correction for biological decay.

I^{132}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At D \neq 1, I^{132} intake is 1.1×10^{11} atoms

The average mean energy of I^{132} is about 0.55 mev or 2.5 times that of I^{131} .

Thus, the energy equivalent to I^{131} would be

$$(1.1 \times 10^{11})(2.5) = 2.75 \times 10^{11} \text{ atoms of } I^{131}$$

However, due to the short half-life of I^{132} (2.4 hrs) assume that the energy equivalent of 1.5×10^{11} atoms of I^{131} reaches the thyroid.

Thus, the ratio of doses $\frac{I^{131}}{I^{132}} = 9.0$

This material contains neither recommendations nor conclusions of the United States within the meaning of the采

OFFICE	U. S. GOVERNMENT PRINTING OFFICE	16-2761-2
SURNAME	██████████	██████████
DATE	██████████	██████████

Dr. Bugher

- 4 -

May 13, 1954

I¹³³

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At D ≠ 1, I¹³³ intake is 1.24×10^{12} atoms

The average mean energy of I¹³³ is about 0.36 or 1.63 times
that of I¹³¹.

Thus, the ratio of doses $\frac{I^{131}}{I^{133}} \approx 0.7$

I¹³⁵

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At D ≠ 1, I¹³⁵ intake is 2.36×10^{11} atoms.

The average mean energy of I¹³⁵ is about 0.3 mev or 1.36
that of I¹³¹.

Thus, the energy equivalent to I¹³¹ would be

$$(2.36 \times 10^{11})(1.36) = 3.2 \times 10^{11} \text{ atoms of } I^{131} \text{ energy equivalent.}$$

Thus, ratio of doses $\frac{I^{131}}{I^{135}} \approx 4.2$


This document contains neither recommendations nor conclusions of the U.S. Government. It has been reviewed by the U.S. Government and approved for release under the Freedom of Information Act. It contains neither recommendations nor conclusions of the U.S. Government.

OFFICE ►	U. S. GOVERNMENT PRINTING OFFICE	16-661-2
SURNAME ►	[REDACTED]	
DATE ►	[REDACTED]	

Dr. Bugher

- 5 -

May 13, 1954



Assume: inhalation and/or ingestion of 5×10^{13} fissions.

At D/1, Te¹³² intake 100 pc

or 1.5×10^{12} atoms

Assume: the time spent in the gut is 77 hrs.

Then, 7.3×10^{11} atoms of Te¹³² will have disintegrated into 7.3×10^{11} atoms of I¹³².

The average mean energy of I¹³² is about 0.55 Mev or 2.5 times that of I¹³¹.

Thus, the energy equivalent to I¹³¹ would be

$$(7.3 \times 10^{11})(2.5) = 1.8 \times 10^{12} \text{ atoms of I}^{131}.$$

However, due to the short half-life of I¹³² (2.4 hrs), assume that only the energy equivalent of 1×10^{12} atoms of I¹³¹ reaches the thyroid.

Thus, the ratio of doses $\frac{\text{I}^{131}}{\text{I}^{132}(\text{Te}^{132})} \approx 1.4$.

DISTRIBUTION: copy 1A - Dr. Bugher
2A - Dr. Fanning
3A - Biophysics Branch
4,5,6A - R&D Files

Copy 5A destroyed 6/14/54

OFFICE ►							
SURNAME ►							
DATE ►							