

Japan Science Council, Ueno, Tokyo-Wednesday, November 17

FOR HERROTATE KELEASE

The third day of the Japan - United States Conference on Radiobiology was spent in discussion of United States and Japanese methods for measuring the intensity of atomic radiation from various sources. The United States Delegation exhibited and demonstrated a wide range of radiation letection instruments employed for differ nt purposes. Nembers of both delegations presented reports on methods used for analyzing and applying the radiation measurements obtained in laboratory and field.

The chairman for the mermine appsion was Dr. Funio Yamasaki of the chairman delegation; for the afternoon, Dr. Storling B. Hendricks of the United States delegation.

The first subject was equipment employed for sampling air and removing airborne redirectivity particles for measurement in the laboratory. The american air sampling equipment used in atomic plants to help safegrard the health of workers and in outdoor locations to gather the air-borne dusts was whiteful, their specifications given, and their operation explored. There was particular interest on the part of the Japaness delegation in the filter papers and their substances used to extract particles from the dir; also in the lafort squares of samed paper employed at more than a huntral locations in the United States to collect samples of the particles falling freely from the statesphere or brought down by rain or snow.

The discussion next turn dute the instruments used to detect and measure the different types of radiation either in someles collected from plants or catalogs or the radiation which occurs in nature. Associal attention was call to standard methods of collibrating instruments. It was pointed out by Dr. John Harley of the United States Delegation, who opened this discussion, that different readings as between two instruments in the same laboratory or as between instruments in the same laboratory are as between instruments in different southers as on as between instructs in different southers a confirm lack of uniform collibration.

Both deferrations record if their calibration methods. The discussion remode ever not only calibration, but the use and the interpretation of readings on beta-gamma survey instruments, film bedges, ionization chambers, and scintillation counters, and on aix different types of instruments for as curing neutron rediction. There was particular interest in the latter discussion on the part of Dr. 3. Shimizu of Ayoto University, who is building anyeletron and wish a to install the best available neutron instrumentation for use in research s with this machine. Dr. Harley and kr. Legril Eisenbud, who represents the United States delegation in the instrumentation presentations, made extensive as most only of instruments and components but of color slides of the larger types of fixed instruments employed in larger laboratories which could not be transported to the confer nee, and showed action platures of a quirment in operation.

.d.hel

The afternoon session was opened at 1:30 F.M. with Dr. Hendricks presiding. A 16 mm motion picture film and slides were presented to explain in detail the equipment, instruments and methods used on a large scale for surveying the distribution of fallout of radioactive ash when atomic bomb tests are carried out in continental United States. Some of the actual equipment was shown. This showed the procedures for collecting the radioactive dust on gummed paper at the observation points scattered around the country and, mailing to the Atomic Energy Commission, where the many samples are handled mechanically and efficiently, baked into ash, and are automatically measured and recorded for radioactivity.

Afterwards, Dr. Harley used slides and printed matter to explain the method for analyzing minute quantities of radioactive substances included in the material. Interested questions were asked by Dr. Miyake, Dr. Mimura, and others of the Japanese delegation.

Dr. Hendricks followed, using a printed text, giving a detailed explanation of the methods for protecting against various radicactive contamination of workers and the laboratory when carrying out experiments with various radicactive elements. These methods included such factors as the construction of laboratory rooms and the prevention of contamination of the equipment and dask surfaces. A few questions were asked about this subject by the Japanese delegates, such as what should be done for disposing radicactive waste water down the drain, which had been used in the laboratory. Also Dr. Harley was asked what should be done to dispose safely of the samples of radicactive dust, to which question he answered that all of the samples were stored for record.

The day's conference closed at 4 o'clock.