403141

BNL 51958 UC-48 (Biology & Medicine ----TIC-4500)

MEDICAL STATUS OF MARSHALLESE ACCIDENTALLY EXPOSED TO 1954 BRAVO FALLOUT RADIATION: JANUARY 1983 THROUGH DECEMBER 1984

William H. Adams, M.D., John R. Engle, M.D., James A. Harper, M.D., Peter M. Heotis, and William A. Scott



MEDICAL DEPARTMENT

BROOKHAVEN NATIONAL LABORATORY UPTON, LONG ISLAND, NEW YORK 11973

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency, contractor or subcontractor thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency, contractor or subcontractor thereof.

Printed in the United States of America Available from National Technical Information Service U.S. Department of Commerce 5285 Port Royal Road Springfield, VA 22161

.

NTIS price codes: Printed Copy: A04; Microfiche Copy: A01

CONTENTS

	Page
List of Participants in Medical Surveys, 1983-1984	v
Introduction	1
Exposure Groups	2
The Brookhaven Medical Program	2
The Brookhaven Medical Team	4
Laboratory Support	4
Medical Findings	
Recent Mortality	5
Hematology	6
Hepatitis B Serological Survey	6
Tuberculin and Candida Sensitivity	8
Hyperprolactinemia	10
Thyroid Hypofunction	10
Thyroid Neoplasia	11
Individual Laboratory Data	12
Acknowledgments	13
Bibliography	14
Appendices	19

Marshall Islands Survey Participants (1983-1984)

Professional Staff

Name, Participating Survey

Adams, William H., M.D. (Mar. '83, Oct. '83, Mar. '84, Oct. '84)

Arelong, Totha, R.N. (Mar. '83, Oct. '83, Mar. '84, Oct.'84)

Barclay, Paula Jane, M.D. (Mar. '84)

Cheatham, Wayman, M.D. (Mar. '83, Mar. '84)

Dungy, Claibourne, M.D. (Oct. '83)

Engle, John, M.D. (Mar. '84, Oct. '84)

Ferguson, Fred S., D.D.S. (Oct. '83, Oct. '84)

Geller, Paul, D.D.S. (Oct. '83)

Giorgio, Bernard W., M.D. (Mar. '83, Mar. '84)

Harper, James A., M.D. (Mar. '83, Oct. '83)

Jackson, Rebecca D., M.D. (Mar. '83)

Affiliation

Scientist, Brookhaven National Laboratory, Upton, NY 11973 (Currently Principal Investigator, Marshall Islands Study)

Nurse Practitioner Armer Ishoda Memorial Hospital Majuro, Marshall Islands 96960

Kwajalein Hospital APO San Francisco 96555

Department of Clinical Investigation Walter Reed Army Hospital Washington, DC 20012

Assoc. Prof., Dept. of Pediatrics Irvine Medical Center University of California, Irvine Orange, CA 92668

Brookhaven National Laboratory Medical Department Stationed at Kwajalein APO San Francisco 96555

Associate Professor Department of Children's Dentistry School of Dental Medicine SUNY at Stony Brook Stony Brook, NY 11794

Private Practice Brentwood, NY 11717

Obstetrics and Gynecology Private Practice Pearl City, HI 96782

Family Practice Physician Private Practice Portland, ME 04101 Formerly Brookhaven National Laboratory, stationed at Kwajalein

Internal Medicine/Endocrinology Ohio State University Columbus, OH 43210

PROFESSIONAL STAFF (Continued)

Name, Participating Survey

Kabua, Jenuk, R.N. (Mar. '83, Oct. '83, Mar. '84, Oct. '84)

Kaloyanides, George J., M.D. (Mar. '83)

Kehne, Susan, M.D. (Mar. '84)

Kindermann, Reed, M.D. (Mar. '83)

Lerner, Marc, M.D. (Oct. '84)

Malarkey, William B., M.D. (Mar. '83)

McClintock, Claudia, M.D. (Mar. '83, Mar. '84)

Morgan, Beverly, M.D. (Oct. '83, Oct. '84)

Nakasone, Ken, M.D. (Mar. '84)

O'Sullivan, Mary Josephine, M.D. (Mar. '84)

Randell, David, M.D. (Mar. '83)

Affiliation

Nurse Practitioner Brookhaven National Laboratory Stationed at Ebeye, Marshall Islands

Professor of Medicine Director, Division of Nephrology and Hypertension SUNY at Stony Brook Stony Brook, NY 11794 Pediatric Neurology Boston City Hospital Boston, MA 02118

Ophthalmology and Ophthalmic Surgery Private Practice Cherry Hill, NJ 08003

Department of Pediatrics Irvine Medical Center University of California, Irvine Orange, CA 92668

Professor of Medicine Div. Endocrinology and Metabolism Ohio State University Columbus, OH 43210 Internal Medicine/Gastroenterology Boston City Hospital Boston, MA 02118

Professor and Chairperson Department of Pediatrics Irvine Medical Center University of California, Irvine Orange, CA 92668

Obstetrics and Gynecology Private Practice The Honolulu Medical Group Honolulu, HI 96813

Professor of Medicine Dept. of Obstetrics and Gynecology Univ. of Miami School of Medicine Miami, FL 33103

Ophthalmology Private Practice Kaneohe, HI 96744

PROFESSIONAL STAFF (Continued)

Name, Participating Survey

Sherman, Lawrence, M.D. (Mar. '84)

Stone, Martin L., M.D. (Mar. '83)

Affiliation

Professor of Medicine Associate Dean of Academic Affairs SUNY at Stony Brook Stony Brook, NY 11794

Professor/Chairman Dept. of Obstetrics and Gynecology SUNY at Stony Brook Stony Brook, NY 11794

TECHNICAL SPECIALISTS

Adams, Diana (Mar. '83, Oct. '83, Mar. '84, Oct.'84)

Bellu, Will (Oct. '83)

de Brum, Reynold (Mar. '83, Oct. '83, Mar. '84, Oct. '84)

Emos, Helmer (Mar. '83, Oct. '83, Mar. '84, Oct. '84)

Ferguson, Robert (Mar. '83, Oct. '83, Mar. '84)

Heotis, Peter (Mar. '83, Oct. '83, Mar. '84, Oct. '84)

Jacob, Stanley (Mar. '84)

Saul, Joe (Mar. '83, Oct. '83, Mar. '84, Oct. '84)

Scott, William (Mar. '83, Oct. '83, Mar. '84, Oct. '84)

Shoniber, Sebio (Mar. '83, Oct. '83) Medical Department Brookhaven National Laboratory Upton, NY 11973

Ebeye Hospital Ebeye, Marshall Islands 96970

U.S. Department of Energy Majuro, Marshall Islands 96960

Medical Department Brookhaven National Laboratory Stationed at Ebeye, Marshall Islands

Medical Department Brookhaven National Laboratory Upton, NY 11973

Medical Department Brookhaven National Laboratory Upton, NY 11973

Ebeye Hospital Ebeye, Marshall Islands 96970

Armer Ishoda Memorial Hospital Majuro, Marshall Islands 96960

Medical Department Brookhaven National Laboratory Upton, NY 11973

Armer Ishoda Memorial Hospital Majuro, Marshall Islands 96960

Introduction

March 1, 1984, was the 30th anniversary of the Bravo thermonuclear test that resulted in the accidental exposure of the populations of Rongelap and Utirik atolls to radioactive fallout. The chronicling of the medical events resulting from that exposure is continued in this report, which covers the period from January 1983 through December 1984. Humanitarian concern for the exposed Marshallese and for other human populations that might suffer from some future exposure continues to be manifested in the worldwide interest of many individuals and institutions who request Brookhaven National Laboratory reports and other published medical articles describing the medical findings. Therefore, an updated listing of all relevant publications from the Medical Department, Brookhaven National Laboratory, is presented in the Reference Section. Articles not issued by Brookhaven National Laboratory but which also relate to the medical aspects of the Marshallese radiation exposure are included for those desiring further information on the subject. Finally, the listing includes Brookhaven National Laboratory-sponsored articles containing Marshallese data that do not concern radiation. For the most recent comprehensive reviews of the principal medical findings since the fallout exposure, the reader is referred to two reports by Dr. Robert A. Conard, director of the Marshall Islands medical program for many years (Conard et al. 1980a; Conard 1984).

Thirty years of observation continue to show no detectable increase in mortality in the exposed population as a result of that exposure. The survival curves of the high-exposure Rongelap group, the low-exposure Utirik population, and an unexposed group of Rongelap people matched by age and sex to the exposed Rongelap group in 1957 continue to be similar (Figure 1). This is not surprising because Japanese A-bomb survivors, which include a far greater number of radiation-exposed individuals, many of whom received a much higher radiation dose than the people of Rongelap, have also had no overall shortening of life-span, even when correlated with radiation dose (Kato et al. 1982). A separate study of Nagasaki A bomb survivors revealed their




Figure 1. Percent survivors of the different exposure groups since 1954. The curves are based on the total original populations, including those *in utero*.

1970-1984 age-specific death rates from all causes to be lower than controls, although it has been suggested that the programs providing health screening of these populations might have led to an underestimation of the effect of radiation on mortality (Okajima et al. 1985).

Clearly, therefore, concern over the consequences of the 1954 exposure transcends mortality statistics. The general health of the exposed population, morbidity directly or indirectly related to the exposure, and present and future risks continue to be monitored and evaluated by the Brookhaven National Laboratory Marshall Islands medical program. The program pursues two related objectives. One is the provision of a cancer-oriented annual examination that follows, as nearly as practicable, the recommendations of the American Cancer Society (1980). The other is a placing in perspective of the risks of radiation exposure as they relate to the overall health of the individual and the Marshallese community. Diabetes mellitus, for example, is a major health problem in the Republic of the Marshall Islands, affecting some 17% of the adults examined by the medical program. Attention to its attendant complications of renal failure, blindness, severe bacterial infection, peripheral neuropathy, impotence, and accelerated atherosclerotic disease should not be minimized because the focus of the program, as mandated by Public Law 95-134, is necessarily on radiation-related illness. The medical program has continued to address such problems by forwarding periodic

reports to the Health Services of the Government of the Republic of the Marshall Islands on public health matters identified by the Brookhaven medical teams. In 1983-1984 these public health reports included information concerning the prevalence of hepatitis B, the growth of Marshallese children, tuberculin skin-test positivity, a survey for syphilis in young adults, and the prevalence of anemia in Marshallese children. It was a related investigation, which identified high levels of fecal contamination of well water on Rongelap and Utirik, that led to the construction of a large concrete cistern on each of the two atolls. This was a joint effort of the Department of Energy Pacific Area Support Office and the Government of the Republic of the Marshall Islands. The contents of the public health reports are always presented to the Marshallese communities at the time of the "town meetings" which precede each medical examination session on the atolls visited by the medical team.

Exposure Groups

As in recent years, the medical program continues to examine and treat some 1200 to 1400 persons annually, half of whom are children. For purposes of comparison, however, the exposure groups defined in the last Brookhaven National Laboratory report are the same as those from which the statistics herein have been collected (Adams et al. 1984b). They are described below:

Rongelap

Now numbering 50, this group received an estimated 190 rads of absorbed external gamma radiation. Of the 67 persons originally exposed in 1954, 3 were *in utero*.

Ailingnae

Nineteen persons, including 1 *in utero*, received an estimated 110 rads of absorbed external gamma radiation. Twelve persons are now in this group.

Utirik

One hundred twelve persons are currently alive in this group. The original 167 individuals who were exposed, including 8 in utero, received an estimated absorbed external gamma radiation dose of 11 rads.

Comparison

In 1957, 86 unexposed Rongelap persons were individually matched by age and sex with the combined exposed Rongelap and Ailingnae groups (Conard et al. 1958). Sixty persons remain in this matched group, against which the overall survival of the exposed population is compared (Figure 1).

A second, larger unexposed group continues to be followed. Currently numbering 135, the age and sex distributions of its members are statistically similar to those of the combined Rongelap-Ailingnae groups and the Utirik group (Adams et al. 1984b). It is this larger unexposed population that is used for the statistical comparison of year-by-year medical events and that provides baseline prevalences from which unexpected consequences of the radiation exposure of persons from Rongelap and Utirik can be identified.

Unless otherwise specified, the term Rongelap, when referring to the high-exposure group, combines those who were on Rongelap and those who were on Ailingnae at the time of exposure.

The Brookhaven Medical Program

Under Public Law 95-134, the Department of Energy has a contract with the Brookhaven National Laboratory Medical Department to provide for diagnosis and treatment of radiation-related disease among the exposed populations of Rongelap and Utirik. Although considerable effort is spent on the care of acute and chronic illnesses of any etiology, a program is in place which is oriented toward the problems posed by their 1954 radiation exposure. The exposed population must be considered at increased risk for malignant disease (Wakabayashi et al. 1983), and chief among the responsibilities of an ongoing program is a cancer-related evaluation. There may be additional risks unrelated to malignancy. The current strategy of the medical program is outlined below.

1. A cancer-related examination is provided, using as a guide the current recommendations

of the American Cancer Society. The program now includes:

a. A review of systems and a complete medical examination.

b. Advice on decreasing risk factors and on self-detection of lesions.

c. Pelvic examinations with Papanicolaou smears.

d. Stool testing for occult blood.

e. A mammography unit and a flexible 65cm sigmoidoscope have been recently acquired.

2. Pursuant to the intent of PL 95-134, the examinations and procedures listed under (1) are performed more frequently than proposed by the American Cancer Society for populations not at increased risk for cancer. Therefore, the physical examinations are annual and include a pelvic examination and Pap smear for all exposed women. Annual mammograms, using a new low-dose mammography unit, will begin at age 35. Routine mammography was not begun earlier because older machines produced doses of x rays which were judged unacceptable for routine annual screening of a population already at increased risk for radiogenic breast cancer. Rectal examinations and stool testing for occult blood are done annually, starting at least by age 40. Routine flexible sigmoidoscopy will be offered before age 50 and will be repeated every other year, or more frequently if clinically indicated.

3. The delayed effects of radiation exposure are generally considered to be limited to malignant disease. The exposed Marshallese, however, receive additional attention for two reasons. First, their radiation exposure was of a unique type, and a tabulation of risks derived from the statistics of other irradiated populations may not cover the range of late consequences that could befall them. Second, data now collected by the Brookhaven medical program suggest previously undocumented late effects of radiation exposure in man. These include an increased incidence of pituitary neoplasms and a trend toward lower blood cell counts (Adams et al. 1984a, 1984b). Another late effect, hypothyroidism, was documented in some of the exposed Rongelap during earlier years of the program (Larson et al. 1982). Therefore, nonmalignant endocrine neoplasms, endocrine dysfunction, and hematologic abnor-

Ĩ

N-24

malities are actively sought. To this end, the medical program provides the following:

a. Annual thyroid examinations by an endocrinologist or surgeon.

b. Thyroid function testing for all exposed persons, annually for the people of Rongelap and biennially for those of Utirik.

c. Thyroid suppression (Synthroid) for all the Rongelap exposed. The intent is to decrease the likelihood of thyroid malignancy.

d. Serum prolactin levels on all exposed persons every three years. The most commonly encountered pituitary tumor in the United States is the prolactinoma.

e. Annual complete blood counts, including a platelet count.

f. Evaluation for "paraneoplastic" evidence of neoplasia, such as monoclonal spikes on serum protein electrophoresis (myeloma, lymphoma) and abnormal serum calcium levels (parathyroid adenoma, hypoparathyroidism, metastatic tumor).

4. There is ongoing evaluation for clinical evidence of depression in immunocompetence. The more recent medical surveys of serum immunoglobulins, toxoplasma antibodies, serologic markers of hepatitis B, and tuberculin sensitivity reveal no good evidence that the exposed Marshallese have a significant impairment of their immune mechanisms (Adams et al. 1984b). However, the matter should not be considered settled, and continued surveillance for evidence of increased risk for unusual manifestations of infectious disease is a part of the medical program.

5. The treatment of any neoplastic process which could conceivably be radiation related is done in referral facilities, generally in Honolulu, Hawaii. The exceptions are thyroid nodule surgery, which continues to be performed by Dr. Brown Dobyns, Professor of Surgery at Case Western Reserve University, and therapy for pituitary neoplasia, which has been done at the National Institutes of Health, Bethesda, Maryland. Few such lesions can be adequately treated in the health facilities of the Republic of the Marshall Islands. The medical program also refers almost all diagnostic workups for malignancy to Honolulu. For example, if the cause of persistent occult blood in the stool is not identified by the medical team, the patient receives x-ray studies, colonoscopy, etc. at one of the excellent medical facilities in Honolulu.

The Brookhaven Medical Team

Physicians, nurses, laboratory technicians, translators, and administrative personnel constitute a "Brookhaven medical team." This phrase does not adequately convey the variegated makeup of the medical missions that are mounted by the Medical Department of Brookhaven National Laboratory. For example, the following medical specialties were represented at least once during the four 1983-84 missions:

> Dentistry (pediatric and adult) Endocrinology Family Practice Gastroenterology Hematology Nephrology Obstetrics and Gynecology Ophthalmology Pediatric Cardiology Pediatrics Physical Medicine Rheumatology Surgery

The physicians and dentists represented in this listing are for the most part affiliated with excellent medical centers throughout the U.S., including Boston University, the National Institutes of Health, Western Reserve, Ohio State University, the University of Miami, the State University of New York (Stony Brook), the University of California (Irvine), Walter Reed Army Hospital, and Wills Eye Hospital (Jefferson Medical College). Other physicians were recruited from private practices in Honolulu, HI, and Portland, ME. The Brookhaven medical team, therefore, represents a broad cross section of medical practitioners in the U.S.; only two of the physicians are, in fact, from Brookhaven National Laboratory. Similarly, all the nurses and translators and half the laboratory personnel are Micronesian. It is clear, therefore, that the Brookhaven medical team is only slightly "Brookhaven" in professional composition.

The ability to recruit excellent doctors from around the U.S. has been one of the strengths of

the medical program. While the volunteer doctors provide the necessary medical examinations and care that are the core of each mission, they also provide consultations in their respective specialties that are often difficult to obtain in the remote atolls that are visited. They also are available for consultations at the Marshall Islands district hospitals on Ebeye and Majuro. Their participation in the medical missions entails in every instance some degree of personal sacrifice. The medical program cannot satisfactorily repay them for their personal and professional efforts in assisting the biennial missions.

In recent years the Straub Hospital and Clinic in Honolulu has been selected as the diagnostic and therapeutic center for Marshallese requiring Brookhaven National Laboratory-sponsored medical referrals. The Brookhaven program is most fortunate in having Dr. Henry Preston of the Department of Internal Medicine at the Straub Clinic volunteer his service as the coordinator and overseer of their care while in Honolulu. The Marshall Islands medical program is very grateful for his fine work.

Laboratory Support

Most medical activities and all laboratory services of the Brookhaven National Laboratory medical surveys are conducted aboard a chartered U.S. Oceanography vessel, Liktanur II. Exceptions include the examinations performed in Brookhaven National Laboratory facilities on Ebeye and pediatric examinations at Rongelap and Utirik which, for reasons of the children's safety, are carried out in dispensaries on shore.

Laboratory support during the medical trips is provided by three to four technicians. Routine five-parameter blood counts are performed on a J.T. Baker 500A electronic particle counter and sizer. Leukocyte differentials and phase contrast platelet counts are done concurrently. A battery of clinical tests (including serum creatinine, glucose, amylase, uric acid, and liver function tests) are carried out on a Beckman spectrophotometer with commercially available reagent kits. Serum and urine sodium and potassium measurements are made on a Beckman Instruments Electrolyte 2 system. Urinalysis (dipstick and microscopic), stool examinations (for occult blood and parasites), and bacteriologic cultures (aerobic and anaerobic) with antibiotic sensitivity testing are available. Hemoglobin A_{1c} determinations, syphilis testing, and erythrocyte sedimentation rates are also provided. Serum is routinely separated and frozen for thyroid function tests and other studies which must be sent to commercial or university laboratories. Fingerstick techniques are used on young children whenever possible. An x-ray machine is available for most commonly required roentgenograms. Electrocardiograms are also available.

Referral laboratories for studies mentioned in this report include: BioScience Laboratories in Honolulu (special chemistries, serologic tests); Pathologists' Laboratories, Inc., in Honolulu (Papanicolaou smear readings); the Endocrinology Laboratory at Brigham and Women's Hospital, Boston (thyroid function tests); Hazleton Laboratories American, Inc., Immunoassay Department, Vienna, VA (prolactin levels); Hepatitis Branch, Division of Viral Diseases, Centers for Disease Control, Atlanta, GA (hepatitis B serology); Brookhaven National Laboratory, Clinical Chemistry Laboratory (serum cholesterol, high-density lipoproteins, triglycerides); and Hematopathology Laboratory, University of California, Irvine Medical Center (free erythrocyte protoporphyrin assays).

Medical Findings

Recent Mortality

The following seven deaths occurred during 1983-84:

Rongelap

Subject No. 80. At the time of his last medical examination in 1982, this 72-year-old man gave clinical evidence of chronic obstructive pulmonary disease. His cigarette smoking history exceeded 60 pack-years. Congestive heart failure was not considered to be the cause of chronic dyspnea. His electrocardiogram showed atrial fibrillation in 1981. It had been present since at least 1965, but his pulse rate was not rapid in 1982. He died in January 1983.

Ailingnae

None

Utirik

Subject No. 2194. When examined in March 1983 this 64-year-old woman had proteinuria, a serum creatinine of 2.3 mg/dl, a hemoglobin of 10.8 g/dl, and diabetic retinopathy. Proteinuria, anemia, and hyperglycemia had been noted as early as 1979, and diabetic retinopathy and a serum creatinine of 2.2 mg/dl were present in 1976. A papillary carcinoma of the thyroid was removed in 1976. A thyroid scan in January 1983 showed minimal residual thyroid in the region of the isthmus; no evidence of metastatic disease was present, although the thyroglobulin level was elevated at 64 ng/ml. The patient was advised to take thyroid hormone replacement, but compliance was poor. In January 1984 she died of a "massive cerebro-vascular accident" in the Majuro hospital following outpatient care of cellulitis.

Subject No. 2157. Diabetes mellitus, mild urinary retention compatible with benign prostatic hypertrophy, and dyspnea on exertion associated with normal lung markings on chest x-ray were noted on this man's 1983 examination when he was 55 years old. He died in January 1984 while residing on Utirik. The cause of death, as diagnosed by the local health aid, was diabetic ketoacidosis.

Subject No. 2168. This patient, a 47-yearold man, had chronic low back pain, a 1-cm left axillary lymph node, and possible hepatomegaly noted in March 1983. His hemoglobin was 15.5 g/dl, and liver function tests were normal except for a slightly elevated serum aspartate aminotransferase level. He had no history of excessive ethanol intake. He died in March 1984 after being admitted to the Majuro Hospital for massive gastrointestinal bleeding. The death certificate identified bleeding from esophageal varices secondary to liver cirrhosis as the cause of death. Serologic tests for hepatitis B, performed on stored serum from his 1983 examination, revealed a positive test for hepatitis B surface antigen.

Subject No. 2185. In March 1983, at age 61, this man had a chronic cough associated with a positive tuberculin skin test and a chest x ray showing no pulmonary disease. He was a cigarette smoker, and cardiology consultation indicated no evidence of cor pulmonale. His weight had remained stable. In January 1984, while returning to Utirik atoll from a fishing trip, the vessel carrying him capsized and he was drowned.

Comparison

Subject No. 1575. This lady died in 1984 at age 78. Her last examination was in March 1981 at which time two thyroid nodules were observed. These were first noted in 1978, but surgery was not performed because of "her age and general senile state." Nevertheless, no serious health problems had been identified and the cause of death is unknown.

Subject No. 1005. In 1982, at age 49, this man's examination revealed no serious medical problems. He had a chronic complaint of shortness of breath. There was a 60-pack-year history of cigarette smoking, but a chest x ray in 1981 had been normal. In 1983 the diagnosis of lung cancer with metastases was made at the Majuro hospital. He died in January 1984.

Hematology

No malignant hematologic disease was diagnosed in 1983-84 in either the exposed or the unexposed populations. Mean values for neutrophils, lymphocytes, and platelets continue to follow the trends of earlier years (Figure 2). Mean hemoglobin levels and monocyte and basophil counts of the Rongelap, Ailingnae, and Utirik groups remain within a few percent of control values (Table 1). Occasionally macrocytosis is seen. It occurs in all groups and is generally borderline in degree. The only person with a clear-cut elevation (MCV of 109 fl) in 1983 was an exposed 72-year-old Rongelap woman. There was concern when a similar value was obtained on her in 1984. It was then learned that prescribed vitamin B12 had not been started. A follow-up MCV was found to be 98 fl. Despite the diagnosis of possible or probable vitamin B_{12} deficiency among Marshallese, intrinsic factor antibodies have yet to be detected. Facilities are not satisfactory for performing Schilling tests, and thus the diagnosis of pernicious anemia remains to be established.

Hepatitis B Serological Survey

The prevalence of hepatitis B is known to be high in Asia and the Western Pacific. For

	Ta	able 1		
Hemoglob	in Concentration, Mor	ocyte Counts, and	l Basophil Counts	
	Rongelap	Ailingnae	Utirik	Comparison
	1	983		
Hemoglobin (M) (g/dl) (F)	$15.2 \pm 1.5^{*}$ 13.6 ± 1.4	$\begin{array}{rrr} 14.9 \ \pm \ 0.9 \\ 13.7 \ \pm \ 0.4 \end{array}$	$\begin{array}{c} 15.7 \ \pm \ 1.2 \\ 13.3 \ \pm \ 1.5 \end{array}$	$\begin{array}{rrr} 15.3 \ \pm \ 1.3 \\ 13.5 \ \pm \ 1.1 \end{array}$
Monocytes/µl	322 ± 148	377 ± 255	316 ± 163	340 ± 179
Basophils/µl	19 ± 37	7 ± 20	19 ± 41	$27\ \pm 49$
	1	.984		
Hemoglobin (M) (g/dl) (F)	$\begin{array}{c} 14.6 \ \pm \ 1.5 \\ 13.5 \ \pm \ 0.7 \end{array}$	$\begin{array}{rrr} 14.0 \ \pm \ 1.0 \\ 12.9 \ \pm \ 0.7 \end{array}$	$\begin{array}{c} 15.7 \ \pm \ 1.1 \\ 13.4 \ \pm \ 1.1 \end{array}$	$15.0 \pm 1.3 \\ 13.5 \pm 1.2$
Monocytes/µl	290 ± 143	234 ± 149	$315\ \pm 157$	285 ± 151
Basophils/µl	20 ± 43	20 ± 34	16 ± 38	18 ± 39
* One standard deviation.				




Figure 2. Mean blood cell counts of the different exposure groups (age 5 years or more) expressed as percent of control, beginning two years after exposure. Values for both sexes are grouped for neutrophils and lymphocytes. Detailed annual observations on Utirik blood cell counts were not begun until 1973. Leukocyte differentials or platelet counts were not obtained for six and five annual examinations, respectively, although for graphing purposes the 100% line has not been broken at those years.

ŧ

example, approximately 60% of inhabitants of American Samoa and 40% of the population of Ponape are reported to have serologic evidence of past infection with this virus (Wong, Purcell, and Rosen 1979). The clinical significance of the cellular immune response in hepatitis B infection is unclear (Hanson et al. 1984; Rustgi et al. 1984). In contrast to hepatitis A, serious late manifestations of disease (chronic active hepatitis, cirrhosis, and hepatocellular carcinoma) are not rare with hepatitis B. It has been suggested that Japanese atomic bombing survivors in the United States do not have a deficit in natural cell-mediated cytotoxicity (Bloom et al. 1983), but studies of the Radiation Effects Research Foundation have revealed an impaired response of lymphocytes to phytohemagglutinin in Japanese receiving >100 rads (Akiyama et al. 1983). If the radiation-exposed Marshallese have an impaired immune mechanism, it is possible that they will be atincreased risk for serious hepatic sequelae if they acquire the infection. For this reason, a serological evaluation of radiation-exposed and unexposed Marshallese was performed in conjunction with the Hepatitis Branch, Division of Viral Diseases, Centers for Disease Control, Atlanta, GA (Dr. Howard Fields and Dr. Stephen Hadler).

Analysis of the results of serologic testing of 314 Marshallese tested revealed that 91.8% gave serologic evidence of past hepatitis B infection. The surveyed population included 98% of the Rongelap group, 82% of the Utirik group, 70% of the comparison population, and 46 younger persons. The last group, ranging in age from 10 to 28 years, was included to evaluate the agespecific prevalence of infection. A tabulation of the hepatitis experience of the different subgroups is presented in Table 2.

There was no difference in the prevalence of serologic evidence of hepatitis B infection among the three exposure groups. However, a significant group difference in the prevalence of hepatitis B surface antigen was detected, with the high-exposure Rongelap group having the lowest prevalence (X^2 =8.17, df=2, p<0.02). This finding contrasts with that of the Radiation Effects Research Foundation, which indicated that the Japanese atomic bombing survivors who received > 100 rads had a significantly higher prevalence of hepatitis B surface antigen than the low-dose groups (3.4% vs 2.0%) (Kato et al. 1983). The reason for the relative infrequency of hepatitis B surface antigenemia among the exposed Rongelap group (2 of 61 persons tested) is not known. However, it is more likely related to local factors rather than to radiation dose because the prevalence of this hepatitis B marker among the unexposed comparison population was not significantly different from that of the Rongelap exposed (X²=1.93, df=1, p>0.10).

Serological evidence of delta agent was not found in any of the persons tested. Delta agent is a co-infecting virus which can affect the host response to hepatitis B. Since the frequency of serious chronic liver disease can be much greater in delta antigen-positive individuals, its absence in the Marshallese is reassuring from the public health perspective.

Tuberculin and Candida Sensitivity

Impaired cellular immunity increases the risk of many types of infection. A survey of skin test responsiveness to mycobacteria and *Candida* was therefore undertaken to determine whether the exposed Marshallese reacted appropriately to these antigens. Another reason for the choice of *M. tuberculosis* testing is the increasing prevalence of tuberculosis in many parts of the world, including Micronesia.

Most persons were evaluated in March 1983. Screening was performed with the Mantoux tuberculin test, where 0.1 ml of PPD containing 5 TU was injected intracutaneously into the forearm in a manner recommended by the American Thoracic Society. A dosage of 0.1 ml of Candida antigen was injected into the opposite arm to test for anergy. After 48 to 72 hours, the amount of induration was measured, with 10 mm or more of induration being considered a positive test. Most individuals with a positive test had a chest x ray taken. Exceptions included those persons who were known, either by personal history or from the medical program records, to have had a positive PPD in earlier years.

A total of 323 PPD tests were applied and read in adults (those ≥ 15 years of age). Of those tested, 147 had a positive test, for a prevalence of 45.5%. One hundred and ten persons received a chest x ray; none revealed evidence of tuber-

	Number Tested		or More ive Tests		BsAg sitive
By sex					
Male	134	123	(91.8)*	20	(14.9)
Female	180	165	(91.7)	16	(8.9)
Combined	314	288	(91.7)	36	(11.5)
By age (yr)					
< 29	46	43	(93.5)	3	(6.5)
29-49	175	158	(90.3)	20	(11.4)
> 49	93	87	(93.3)	13	(14.0)
By atoll of residence **					
Kwajalein	100	89	(89.0)	10	(10.0)
Majuro	74	68	(91.9)	4	(5.4)
Rongelap	61	58	(95.1)	3	(8.5)
Utirik	76	70	(92.1)	19	(25.0)
By radiation exposure group					
Rongelap exposed	61	50	(82.0)	2	(3.3)
Utirik exposed	112	103	(92.0)	21	(18.8)
Rongelap comparison	9 5	86	(90.5)	10	(10.5)
By atoll of residence, excluding Rongelap exp	osed				
Ebeye	69	63	(91.3)	6	(8.7)
Majuro	61	58	(95.1)	4	(6.6)
Rongelap	44	42	(95.5)	3	(6.8)
Utirik	76	70	(92.1)	19	(25.0)

Summary of Positive Serologic Tests for Hepatitis B Surface Antigen (HBsAg), Antibody to Surface Antigen, and Antibody to Core Antigen Among 314 Marshallese

Table 2

* Percent of the total population tested is shown in parentheses.

** Three persons resided outside the atolls listed.

Table 3

Radiation Category	No. in Each Category	No. Tested	Tuberculin Negative	<i>Candida</i> Negative
Rongelap	62	38	16 (42.1%)	2 (5.3%)**
Utirik	137	72	39 (54.2%)	0 (0.0%)
Comparison	135	68	35 (51.5%)	2(2.9%)

Skin Test Responsiveness by Radiation Exposure Group*

* See text for definition of positive and negative tests.

** Two persons, an 83-year-old Rongelap exposed man and a 43-year-old unexposed woman, had positive tuberculin tests despite negative reactions to *Candida* antigen.

culosis. A tabulation of the prevalence of positive and negative tuberculin and *Candida* tests according to radiation group and island of residence at the time of testing is presented in Table 3. The results indicate that the prevalence of positive tuberculin tests and the prevalence of anergy, when analyzed by the chi-square test of independence between two or more samples, were similar among the radiation exposure groups.

The frequency of infection with atypical mycobacteria among Marshallese is unknown. An analysis of size distribution of positive tests indicated 2- to 5-mm induration responses from 14.4% of all persons tested, a finding compatible with past exposure to atypicals.

Hyperprolactinemia

Two exposed women have now been diagnosed as having pituitary tumors (Adams et al. 1984a). In the 1980-82 Brookhaven National Laboratory Marshall Islands report mention was made of another woman, 82 years of age, who had mild but persistent serum prolactin elevations (Adams et al. 1984b). In 1984 this Utirik patient, No. 2182, was brought to Cleveland Metropolitan Hospital for surgery for a suspected thyroid nodule. The presence of the nodule was not confirmed preoperatively, however, and surgery was not performed. Advantage was taken of the availability of CT scanning facilities at the hospital to evaluate her for a pituitary lesion. A CT scan of the skull. with and without contrast, was read as suggesting a lesion within the sella turcica. However, the interpretation of Dr. Azad Anand, neuroradiologist at University Hospital, SUNY, Stony Brook, indicated that there is no evidence for a pituitary tumor. Therefore, although it remains possible that such a tumor exists, no diagnosis can be confirmed at the present time.

Because the possibility of a third pituitary tumor in the small number of exposed persons still under observation would be a clinical finding without precedent, a survey of serum prolactin levels was undertaken in the unexposed comparison group. Of 110 persons tested, five were found to have mildly elevated levels. Four of these were found to be normal on repeat testing. One woman had a persistent mild elevation of serum prolactin (55 ng/ml). She was referred to the Republic of the Marshall Islands Health Service for further evaluation. The number of persons evaluated is too small to derive a prevalence of hyperprolactinemia among Marshallese. Therefore, this finding does not support or refute a conclusion that pathologic hyperprolactinemia and, by inference, prolactinomas are unusually common among the general Marshallese population.

Thyroid Hypofunction

Subclinical thyroid hypofunction, as assessed by thyroid-stimulating hormone (TSH) determinations and response to thyrotropin-releasing hormone (TRH), has been documented in 12 persons in the exposed Rongelap group (Larsen et al. 1982). Annual TSH testing has continued for this group, and biennial testing is provided for the Utirik group. Of 61 persons in the Rongelap group, 57 had TSH levels determined in either or both 1983 and 1984. No new cases of biochemical hypothyroidism were uncovered. However, since all members of this group are advised to take suppressive doses of thyroid hormone (Synthroid), it is possible that new cases are still emerging but are being masked by the administered thyroid hormone. Accurate diagnosis would require the discontinuation of thyroid hormone for several weeks, followed by TSH assays and perhaps TRH stimulation tests. Because little clinical benefit for the Rongelap group is likely, this approach has not been taken.

The Utirik group received much lower thyroid radiation doses in 1954 than did persons on Rongelap, and no thyroxin suppression has been prescribed for them. Thyroid hypofunction has yet to be diagnosed in this group, and, of 104 persons tested in 1983-84, the only elevated TSH levels found were in four individuals who had previously undergone thyroid surgery.

Hypothyroidism has numerous etiologies and occurs not uncommonly in all populations. Its spontaneous frequency is age related, and 4.4% of a Massachusetts population over 60 years of age have been found to have clearly elevated TSH levels (Sawin et al. 1985). The prevalence of biochemical hypothyroidism in unexposed Marshallese was evaluated in 1984. Of 90 persons tested, no TSH elevations were detected.

Hypothyroidism, which is sometimes associated with elevated serum cholesterol levels, may be a risk factor for coronary heart disease (Becker 1985). To determine whether an abnormality in serum lipids may have evolved in the exposed groups as an indirect consequence of radiation injury or thyroid surgery, serum levels of cholesterol, triglyceride, and highdensity lipoprotein were obtained in 1984. The results of an analysis by group are presented in Table 1. There was no significant difference between the mean serum cholesterol levels of the exposed Rongelap or Utirik groups and the unexposed. Since almost all the Rongelap exposed are receiving thyroid hormone in suppressive doses, it is unknown whether or not some of the cholesterol levels would be elevated if thyroxin were not being taken. At this point, then, questions concerning their risk of thyroidrelated hypercholesterolemia are moot. However, an analysis of Rongelap exposed and comparison group cholesterol levels in 1957 revealed the latter to be the higher by 17% (Conard et al. 1958). Analysis of serum cholesterol in persons with known thyroid hypofunction in 1984, as documented by an elevated TSH, and in persons who have had thyroid surgery revealed no values lying outside a normal range established by testing the comparison population (based on two standard deviations from the mean).

One finding that may be of clinical value is the relatively low level of high-density lipoprotein found in all three exposure groups. Since this lipid category, as an independent risk factor, shows an inverse association with coronary heart disease, the low levels found may indicate a propensity for the disorder among Marshallese. However, confirmation of the finding is required to rule out technical problems associated with transport and storage of serum specimens.

Thyroid Neoplasia

The Marshall Islands medical program is most fortunate to have the continued support of four eminent consultant pathologists who review the histologic sections of all thyroid nodules removed at surgery.* The same individuals were among the group of pathologists who, in 1981, reviewed all thyroid sections obtained throughout the history of the program. This has provided consistent year-to-year diagnostic categories of thyroid neoplasia.

In 1983-84, six persons underwent thyroid surgery at Cleveland Metropolitan Hospital

* Dr. L.V. Ackerman, Health Sciences Center, SUNY, Stony Brook, NY; Dr. W.A. Meissner, New England Deaconess Hospital Boston, MA; Dr. A.L. Vickery, Massachusetts General Hospital, Boston, MA; Dr. L.B. Woolner, Mayo Clinic, Rochester, MN.

Exposure Category	n	Cholesterol (mg/dl)	Triglycerides (mg/dl)	High-density Lipoprotein (mg/dl)
Rongelap	<u></u>			
(male)	21	$154 \pm 27^*$	$147 \hspace{.1in} \pm \hspace{.1in} 168$	36 ± 9
(female)	29	170 ± 32	121 ± 67	34 ± 11
Utirik				
(male)	42	177 ± 37	$222 \ \pm 139$	30 ± 5
(female)	49	$187\ \pm\ 35$	$153\ \pm\ 102$	33 ± 5
Comparison				
(male)	34	172 ± 27	173 ± 95	29 ± 6
(female)	60	179 ± 36	143 ± 143	35 ± 8

Table 4

11

(Table 5). Five were from the Utirik-exposed group and one was from the comparison group. The latter was judged to have an adenomatous nodule. Of the five Utirik patients, only four had significant thyroid pathology. Two of the four had occult papillary carcinomas. This is a neoplastic lesion of little clinical significance and is not considered the equivalent of papillary thyroid cancer. It is usually an incidental finding during thyroid surgery, and the prevalence of occult thyroid carcinomas has not been found to be increased in Japanese atomic bombing survivors (Wakabayashi et al. 1983). The other two patients did have papillary thyroid cancers, one of which was associated with lymph node metastases. All these new findings have been incorporated in the summary of thyroid lesions found throughout the history of the medical program (Table 6). An analysis of thyroid cancer risk as it relates to the exposed Marshallese was recently presented, and a summary is given in Appendix A.

INDIVIDUAL LABORATORY DATA

As in the last report, a computerized listing of laboratory test results obtained in 1983-84 and entered by identification number is presented in Appendix B.

Identification Number	Age at Diagnosis	Sex	Consensus Diagnosis
2248	44	F	Occult papillary carcinoma
944	58	Μ	Adenomatous nodule
2149	38	F	No tumor
2152	38	Μ	Papillary carcinoma
2167	44	Μ	Occult papillary carcinoma
2171	33	\mathbf{F}	Papillary carcinoma

Table 5

Table 6

Thyroid	Lesions	Diagnosed	at Surgery	Through 1984

	Adenomatous Nodules	Adenomas	Papillary Carcinomas	Follicular Carcinomas	Occult Papillary Carcinomas
Rongelap (67)*	17	2	4		
Ailingnae (19)*	4	—			1
Utirik (167)*	10	2	4	1†	3
Comparison (227)**	4	1	2	_	2††

NOT INCLUDED are the following unoperated (and therefore unconfirmed) nodules: Rongelap -1; Ailingnae - 1; Utirik - 1; Comparison - 5.

INCLUDED are all consensus diagnoses of a panel of consultant pathologists; two different lesions were detected in one person each from Rongelap, Ailingnae, and Utirik.

* Number of persons (including those in utero) who were originally exposed.

** This number includes all persons who have been in the comparison group since 1957. Some have not been seen for many years; others were added as recently as 1979.

+ Equally divided opinion in one case; follicular carcinoma vs atypical adenoma.

†† Majority opinion in one case; occult papillary carcinoma vs follicular carcinoma. The same patient had a lymphocytic thyroiditis.

Acknowledgments

The professional debts of the Brookhaven National Laboratory Marshall Islands medical program are many, but particularly to be acknowledged are the following:

Dr. Jacob Robbins, Chief of the Clinical Endocrinology Branch, the National Institutes of Health, continues to provide the medical program with exacting guidance in matters relating to clinical endocrinology.

Dr. Brown Dobyns, Professor of Surgery, Case Western Reserve University, Cleveland, OH, by his personal and professional commitment in assisting the exposed Marshallese with thyroid problems requiring surgery, remains an invaluable resource to the medical program.

Dr. P. Reed Larsen, Director of the Thyroid Diagnostic Laboratory, the Brigham and Women's Hospital, Boston, MA, makes available to the program endocrinologic testing of the highest quality.

Dr. Beverly Morgan, Professor and Chairperson, Department of Pediatrics, the University of California (Irvine), has personally and through her medical staff contributed enormously to an effective effort in pediatric care for the children of Rongelap and Utirik.

Acknowledged with particular pleasure is the Deputy for Pacific Operation, D.O.E. (Nevada),

Mr. Roger Ray. His long acquaintance with and sincere concern for the Marshallese people has helped foster a medical program which will benefit the exposed Marshallese for many more years.

The medical program continues to benefit from the highly competent efforts of the Department of Energy Pacific Area Support Office (William J. Stanley, Harry U. Brown, William D. Jackson). Their ability to manage logistic and political details so necessary for an effective and unencumbered medical program has been outstanding. Captain Keith Coberly and the crew of Liktanur II are entrusted with the safety and well being of the medical teams for much of each mission. That trust has been well earned; their efforts have been exemplary and extend far beyond those to be expected. The assistance of Ms. Geraldine Callister, whose enthusiasm and familiarity with the secretarial requirements of the medical program are so essential, is gratefully acknowledged. Dr. Robert Aronson kindly developed a computerized format for listing the references. Mrs. Mary Rustad has again competently guided a Marshall Islands report to completion. Dr. Victor Bond, in reviewing the manuscript, continues to add to an already long and notable record of support for the program.

13

Bibliography

Included in this listing are 1) references cited in the text, 2) all Brookhaven National Laboratory-sponsored publications which include Marshallese data, and 3) selected publications not from Brookhaven National Laboratory which bear on the radiation exposure of the people of Rongelap and Utirik.

- American Cancer Society. 1980. Guidelines for the cancer-related checkup: Recommendations and rationale. CA - A Cancer J. Clin. 30 (4).
- Adams, W.H., Harper, J.A., Rittmaster, R.S., and Grimson, R.C. 1984a. Pituitary tumors following fallout radiation and exposure. J. Am. Med. Assoc. 252:664-6.
- Adams, W.H., Harper, J.A., Rittmaster, R.S., Heotis, P.M., and Scott, W.A. 1984b. Medical Status of Marshallese Accidentally Exposed to 1954 BRAVO Fallout Radiation: January 1980 Through December 1982. BNL 51761.
- Adams, W.H., Harper, J.A., Heotis, P.M., and Jamner, A.H. 1984c. Hyperuricemia in the inhabitants of the Marshall Islands, *Arthritis Rheum*. 27:713-6.
- Akiyama, M., Yamakido, M., Kobuke, K., Dock, D.S., Hamilton, H.B., Awa, A.A., and Kato, H. 1983. Peripheral lymphocyte response to PHA and T-cell population among atomic bomb survivors. *Radiat. Res.* 93:572-80.
- Allison, A.C. and Blumberg, B.S. 1961. An isoprecipitation reaction distinguishing human serum-protein types. *Lancet* 1:634-7.
- Beasley, T.M., Held, E.E., and Conard, R.A. 1972. Iron-55 in Rongelap people, fish, and soils. *Health Phys.* **22**:245-50.
- Becker, C. 1985. Hypothyroidism and atherosclerotic heart disease: Pathogenesis, medical management, and the role of coronary artery bypass surgery. *Endocr. Rev.* 6:432-40.
- Bloom, E.T., Korn, E.L., Toji, D.S., and Makinodan, T. 1983. Natural cell-mediated cytotoxicity among A-bomb survivors residing in the United States. Int. J. Radiat. Biol. 44:213-8.
- Blumberg, B.S. and Gartler, S.M. 1959. High prevalence of high-level B-amino-isobutyric acid excretors in Micronesians. *Nature* 184: 1990-2.
- Bond, V.P., Conard, R.A., Robertson, J.S., Weden, E.A. Jr. 1955. Medical Examination of Rongelap People Six Months After Exposure to Fallout. Armed Forces Special Weapons Project, Sandia Base, Albuquerque, NM, Report No. 937.
- Cohn, S.H., Robertson, J.S., and Conard, R.A. 1960. Radioisotopes and environmental circumstances: The internal radioactive contamination of a Pacific

Island community exposed to local fallout. In: Radioisotopes in the Biosphere, Chapter 21, pp. 306-30, Caldecott, R.S. and Snyder, L.A. (Editors). Published by Center for Continuation Study, University of Minnesota.

- Cohn, S.H., Conard, R.A., Gusmano, E.A., and Robertson, J.S. 1963. Use of a portable whole-body counter to measure internal contamination in a fallout-exposed population. *Health Phys.* **9**:15-23.
- Conard, R.A. 1967. Further development of thyroid nodules in Marshallese population accidentally exposed to radioactive fallout in 1954. Suffolk County Med. Soc. Bull., p. 25.
- Conard, R.A., Shulman, N.R., Wood, D.A., Dunham, C.L., Alpen, E.L., and Browning, L.E. 1955. Skin Lesions, Epilation, and Nail Pigmentation in Marshallese and Americans Accidentally Contaminated With Radioactive Fallout. Naval Medical Research Institute and Naval Radiological Defense Laboratory, NMRI Report NM 006 012 04 82, pp. 423-500.
- Conard, R.A. 1956. The effects of fallout radiation on the skin: In: *The Shorter-Term Biological Hazards* of a Fallout Field, pp. 135-42, AEC-DOD, U.S. Government Printing Office, Washington, DC.
- Conard, R.A., Cannon, B., Huggins, C.E., Richards, J.B., and Lowrey, A. 1956a. Medical Survey of Marshallese Two Years After Exposure to Fallout Radiation. BNL 412 (T-80).
- Conard, R.A. et al. 1956b. Response of human beings accidentally exposed to significant fallout radiation from a thermonuclear explosion (summary).
 In: Progress in Radiobiology, pp. 491-3, Mitchell, J.S., Holmes, B.E., and Smith, G.G. (Editors), Oliver and Boyd, Edinburgh.
- Conard, R.A., Cannon, B., Huggins, C.E., Richards, J.B., and Lowrey, A. 1957. Medical survey of Marshallese two years after exposure to fallout radiation. J. Am. Med. Assoc. 164: 1192-7.
- Conard, R.A. 1957. Three year medical survey of the Marshallese people exposed to fallout in March 1954. *Radiat. Res.* 3:309.
- Conard, R.A., Meyer, L.M., Rall, J.E., Lowrey, A., Bach, S.A., Cannon, B., Carter, E.L., Eicher, M., and Hechter, H. 1958. March 1957 Medical Survey of Rongelap and Utirik People Three Years After Exposure to Radioactive Fallout. BNL 501 (T-119).
- Conard, R.A., Meyer, L.M., Robertson, J.S., Sutow, W.W., Wolins, W., and Hechter, H. 1959a. Effects of fallout radiation on a human population. *Radiat. Res. Suppl.* 1:280-95.
- Conard, R.A., Robertson, J.S., Meyer, L.M., Sutow, W.W., Wolins, W., Lowrey, A., Urschel, H.C., Barton, J.M., Goldman, M., Hechter, H., Eicher, M., Carver, R.K., and Potter, D.W. 1959b. Medical Sur-

paredness, pp. 55-8, Hubner, K.F. and Fry, S.A. (Editors), Elsevier North Holland, Inc.

- Conard, R.A. et al. 1980. Review of Medical Findings in a Marshallese Population Twenty-Six Years After Accidental Exposure to Radioactive Fallout. BNL 51261.
- Conard, R.A. 1984. Late radiation effects in Marshall Islanders exposed to fallout twenty-eight years ago. In: *Radiation Carcinogenesis, Epidemiology* and Biologic Significance, pp. 57-71, Boice, J.D. Jr. and Fraumeni, J.F. (Editors), Raven Press Inc., New York.
- Contreras, P., Generini, G., Michelsen, G., Pumarino, H., and Campino, C. 1981. Hyperprolactinemia and galactorrhea: Spontaneous versus iatrogenic hypothyroidism. J. Clin. Endocrinol. Metab. 53:1036-8.
- Cronkite, E.P., Bond, V.P., Browning, L.E., Chapman, W.H., Cohn, S.H., Conard, R.A., Dunham, C.L., Farr, R.S., Hall, W.S., Sharp, R., and Shulman, N.R. 1954. Study of Response of Human Beings Accidentally Exposed to Significant Fallout Radiation. Armed Forces Special Weapons Project, Sandia Base, Albuquerque, NM, Report No. WT-923.
- Cronkite, E.P., Dunham, C.L., Griffin, D., McPherson, S.D., and Woodward, K.T. 1955a. Twelve-Month Postexposure Survey on Marshallese Exposed to Fallout Radiation. BNL 384 (T-71).
- Cronkite, E.P., Bond, V.P., Conard, R.A., Shulman, N.R., Farr, R.S., Cohn, S.H., Dunham, C.L., and Browning, L.E. 1955b. Response of human beings accidentally exposed to significant fallout radiation. J. Am. Med. Assoc. 159:430-4.
- Cronkite, E.P., Bond, V.P., and Dunham, C.L. 1956. Some Effects of Ionizing Radiation on Human Beings. AEC-TID 5358.
- Dahl, L.K. 1961. Possible role of chronic excess salt consumption in the pathogenesis of essential hypertension. Am. J. Cardiol. VIII 4:571-5; 587-92.
- Demoise, C.F. and Conard, R.A. 1972. Effects of age and radiation exposure on chromosomes in a Marshall Island population. J. Gerontol. 27:197-201.
- Dungy, C.I., Morgan, B.C., and Adams, W.H. 1984. Pediatrics in the Marshall Islands. *Clin. Pediatr.* 23:29-32.
- Goldman, M. and Carver, R.K. 1959. An intestinal parasite survey on Rongelap atoll in the Marshall Islands. *Am. J. Trop. Med. Hyg.* 8:417-23.
- Hanson, R.G., Hoffnagle, J.H., Minuk, G.Y., Purcell, R.H., and Gerin, J.L. 1984. Cell-mediated immunity to hepatitis B surface antigen in man. *Clin. Exp. Immunol.* 57:257-64.
- Hardy, E.P., Rivera, J., and Conard, R.A. 1965. Cesium-137 and strontium-90 retention following an acute ingestion of Rongelap food. In: *Radioac*tive Fallout From Nuclear Weapons Tests, pp. 743-57, Klement, A.W. Jr. (Editor), AEC Conf. 765.
- James, R.A., 1964. Estimate of Radiation Dose to Thyroid of Rongelap Children Following the

BRAVO Event. Lawrence Radiation Laboratory Report, UCRL-12273.

- Kato, H., Brown, C.C., Howl, D.G., and Schull, W.J. 1982. Studies of the mortality of A-bomb survivors. Report of mortality, 1950-1978: Part II. Mortality from causes other than cancer and mortality in early entrants. *Radiat. Res.* **91**: 243-64.
- Kato, H., Mayumi, M., Nishioka, K., and Hamilton, H.B. 1983. The relationship of hepatitis B surface antigen and antibody to atomic bomb radiation in the adult health study sample, 1975-1977. Am. J. Epidemiol. 117: 610-20.
- Larsen, P.R., Conard, R.A., Knudsen, K., Robbins, J., Wolff, J., Rall, J.E., and Dobyns, B. 1978. Thyroid hypofunction appearing as a delayed manifestation of accidental exposure to radioactive fall-out in a Marshallese population. IAEA Meeting, *The Late Biological Effects of Ionizing Radiation, Proc.* Symp. Vol. 1, pp. 101-15.
- Larsen, P.R., Conard, R.A., Knudsen, K.D., Robbins, J., Wolff, J., Rall, J.E., Nicoloff, J.T., and Dobyns, B. M. 1982. Thyroid hypofunction after exposure to fallout from a hydrogen bomb explosion. JAMA, J. Am. Med. Assoc. 247: 1571-5.
- Lessard, E.T. and Conard, R.A. 1983. Exposure to fallout: The radiation dose experience at Rongelap and Utirik atolls. In: *Radiat. Res. Proc. 7th Int. Congr.* Vol. C., pp. C8-09, Broerse, J.J., Barendsen, G.W., Kal, H.B., and Van Der Kogel, A.J. (Editors), Martinus and Nijhoff, Amsterdam.
- Lessard, E.T. and Conard, R.A. 1984. Protracted exposure to fallout: The Rongelap and Utirik experience. *Health Phys.* **46**(3):511-27.
- Lessard, E., Miltenberger, R., Conard, R., Musolino, S., Naidu, J., Moorthy, A., and Schopfer, C. 1985. *Thyroid-Absorbed Dose for People at Rongelap*, *Utirik, and Sifo on March 1, 1954.* BNL 51882.
- Lessard, E.T., Brill, A.B., and Adams, W.H. 1986. Thyroid cancer in the Marshallese: relative risk of radioiodine and external radiation exposure. Radiopharmaceutical Dosimetry Symp. Proc. 4th Int. Conf., Oak Ridge, TN, Nov. 5-8, 1985 (in press).
- Lisco, H. and Conard, R.A. 1967. Chromosome studies on Marshallese people exposed to fallout radiation. *Science* **157**:445-7.
- Lyon, H.W., Conard, R.A., and Glassford, K.F. 1964. Long-term intraoral findings in humans after exposure to total body irradiation from sudden radioactive fallout. I. Five year post-detonation studies. J. Am. Dent. Assoc. 68: 49-56.
- Naidu, J.R., Greenhouse, N.A., Knight, G., and Craighead, E.C. 1980. Marshall Islands: A Study of Diet and Living Patterns. BNL 51313.
- Neel, J.V., Ferrell, R.E., and Conard, R.A. 1976. The frequency of "rare" protein variants in Marshall Islanders and other Micronesians. *Am. J. Human Genet.* 28:262-9.

vey of Rongelap People, March 1958, Four Years After Exposure to Fallout. BNL 534 (T-135).

- Conard, R.A. 1960a. Medical survey of Marshallese people five years after exposure to fallout radiation. *Int. J. Radiat. Biol. Suppl.* 1:269-81.
- Conard, R.A. 1960b. An attempt to quantify some clinical criteria of aging. J. Gerontol. 15:358-65.
- Conard, R.A., MacDonald, H.E., Lowrey, A., Meyer, L.M., Cohn, S., Sutow, W.W., Blumberg, B.S., Hollingsworth, J.W., Lyon, H.W., Lewis, W.H., Jaffe, A.A., Eicher, M., Potter, D., Lanwi, I., Riklon, E., Iaman, J., and Heikena, J. 1960a. Medical Survey of Rongelap People Five and Six Years After Exposure to Fallout (with an Addendum on Vegetation), BNL 609 (T-179).
- Conard, R.A., Meyer, L.M., Sutow, W.W., Blumberg, B.S., Lowrey, A., Cohn, S.H., Lewis, W.H. Jr., Hollingsworth, J.W., and Lyons, H.W. 1960b. Medical status of Marshall Islanders in 1959, five years after exposure to fallout radiation. *Nucl. Med.* 1:314-30.
- Conard, R.A. 1961. The biological hazards of a fallout field. In: *Radioactivity in Man*, pp. 249-65, Meneely, G.E. (Editor), C.C. Thomas, Springfield, IL.
- Conard, R.A., MacDonald, H.E., Meyer, L.M., Cohn, S., Sutow, W.W., Karnofsky, D., Jaffe, A.A., and Riklon, E. 1962. *Medical Survey of Rongelap People Seven Years After Exposure to Fallout.* BNL 727 (T-260).
- Conard, R.A. 1963. Effect of acute fallout radiation on a Marshall Island population. Presented at Second Annu. Conf. on Science and Human Survival, Congress of Scientists on Survival, Am. Mus. Nat. Hist., June 15, 1963. BNL 7145.
- Conard, R.A., Meyer, L.M., Sutow, W.W., Moloney, W.C., Lowrey, A., Hicking, A., and Riklon, E. 1963a. Medical Survey of Rongelap People Eight Years After Exposure to Fallout. BNL 780 (T-296).
- Conard, R.A., Meyer, L.M., Sutow, W.W., Moloney, W.C., Cannon, B., Hicking, A., and Riklon, E. 1963b. Medical Survey of the People of Rongelap and Utirik Islands, Nine Years After Exposure to Fallout Radiation (March 1963). Interim Report, BNL 7766.
- Conard, R.A. and Hicking, A. 1965. Medical findings in Marshallese people exposed to fallout radiation: Results from a ten-year study. J. Am. Med. Assoc. 192:457-9.
- Conard, R.A., Meyer, L.M., Sutow, W.W., Lowrey, A., Cannon, B., Moloney, W.C., Watne, A.C., Carter, R.E., Hicking, A., Hammerstrom, R., Bender, B., Lanwi, I., Riklon, E., and Anjain, J. 1965a. Medical Survey of the People of Rongelap and Utirik Islands Nine and Ten Years After Exposure to Fallout Radiation. BNL 908 (T-371).
- Conard, R.A., Meyer, L.M., Sutow, W.W., Robertson, J.S., Jesseph, J.E., Rall, J.E., Eicher, M., Gusmano, E.A., Hicking, A., and Lanwi, I. 1965b. Medical Survey of the People of Rongelap Island, Eleven

Years After Exposure to Fallout Radiation (March 1965). Interim Report, BNL 9698.

- Conard, R.A., Rall, J.E., and Sutow, W.W. 1966a. Thyroid nodules as a late sequel of radioactive fallout in a Marshall Island population exposed in 1954. *New Eng. J. Med.* 274:1392-9.
- Conard, R.A., Lowrey, A., Eicher, M., Thompson, K., and Scott, W.A. 1966b. Ageing studies in a Marshallese population exposed to radioactive fallout in 1954. In: *Radiation and Ageing*, pp. 345-60, Lindop, P.J., and Sacher, G.A. (Editors), Taylor and Francis, London.
- Conard, R.A., Meyer, L.M., Sutow, W.W., Robertson, J.S., Rall, J.E., Robbins, J., Jesseph, J.E., Deisher, J.B., Hicking, A., Lanwi, I., Gusmano, E.A., and Eicher, M. 1967. Medical Survey of the People of Rongelap and Utirik Islands Eleven and Twelve Years After Exposure to Fallout Radiation, March 1965 and March 1966. BNL 50029 (T446).
- Conard, R.A., Sutow, W.W., Colcock, B.P., Dobyns, B.M., and Paglia, D.E. 1969. Thyroid nodules as a late effect of exposure to fallout. In: *Radiation-Induced Cancer*, pp. 325-36, IAEA, Vienna.
- Conard, R.A. (abstract). 1970. Possible radiationinduced ageing as measured by immunohematological changes in a Marshallese population exposed to radioactive fallout. *Radiat. Res. Proc. 4th Int. Congr.*, 1970, p. 168.
- Conard, R.A. et al. 1970a. Medical Survey of the People of Rongelap and Utirik Islands Thirteen, Fourteen, and Fifteen Years After Exposure to Fallout Radiation: March 1967, March 1968, and March 1969. BNL 50220 (T-562).
- Conard, R.A., Dobyns, B.M., and Sutow, W.W. 1970b. Thyroid neoplasia as late effect of exposure to radioactive iodine in fallout. J. Am. Med. Assoc. 214: 316-24.
- Conard, R.A., Demoise, C.F., Scott, W.A., and Makar, M. 1971. Immunohematological studies of Marshall Islanders sixteen years after fallout radiation exposure. J. Gerontol. 26:28-36.
- Conard, R.A. 1972. Effects of ionizing radiations on aging and life shortening in human populations. *Front. Radiat. Ther. Oncol.* 6:486-98.
- Conard, R.A. 1975. Acute myelogenous leukemia following fallout radiation exposure. J. Am. Med. Assoc, 232:1356-7.
- Conard, R.A. et al. 1975. A Twenty-Year Review of Medical Findings in a Marshallese Population Accidentally Exposed to Radioactive Fallout. BNL 50424.
- Conard, R.A. 1977. Summary of thyroid findings in Marshallese 22 years after exposure to radioactive fallout. In: *Radiation-Associated Thyroid Carcinoma*, pp. 241-56, DeGroat, L.J. et al. (Editors), Grune and Stratton, New York.
- Conard, R.A. 1980. The 1954 Bikini Atoll incident. An update of the findings in the Marshallese people. In: *The Medical Basis for Radiation Accident Pre-*

- Okajima, S., Mine, M., and Nakamura, T. 1985. Mortality of registered A-bomb survivors in Nagasaki, Japan, 1970-1984. *Radiat. Res.* **103**: 419-43.
- Popp, R.A., Bailiff, E.G., Hirsch, G.P., and Conard, R.A. 1976. Errors in human hemoglobin as a function of age. *Interdiscip. Top. Gerontol.* 9:209-18.
- Rall, J.E. and Conard, R.A. 1966. Elevation of the serum protein-bound iodine level in inhabitants of the Marshall Islands. Am. J. Med. 40:883-6.
- Robbins, J., Rall, J.E., and Conard, R.A. 1967. Late effects of radioactive iodine in fallout. Ann. Intern. Med. **66**:1214-42.
- Robertson, J.S., Conard, R.A., Gusmano, E. 1966. Radionuclide body burdens in the Rongelap population. *Radiat. Res. Proc. 3rd Int. Congr.*, p. 189.
- Rustgi, V.K., Hoofnagle, J.H., Gerin, J.L., Gelmann, E.P., Reichert, C.M., Cooper, J.N., and Macher, A.M. 1984. Hepatitis B virus infection in the acquired immunodeficiency syndrome. Ann. Intern. Med. 101:795-7.
- Sawin, C.T., Castelli, W.P., Hershman, J.M., Mc-Namara, P., and Bacharach, P. 1985. The aging thyroid: thyroid deficiency in the Framingham Study. Arch. Intern. Med. 145:1386-8.
- Sharp, R. and Chapman, W.H. 1957. Exposure of Marshall Islanders and American Military Personnel to Fallout. Armed Forces Special Weapons

Project, Sandia Base, Albuquerque, NM, Report No. 938.

- Sussman, L., Meyer, L.M., and Conard, R.A. 1959. Blood groupings in Marshallese. *Science* **129**:644-5.
- Sutow, W.W., Conard, R.A., and Griffith, K.M. 1965. Growth status of children exposed to fallout radiation on Marshall Islands. *Pediatrics* **36**:721-31.
- Sutow, W.W. and Conard, R.A. 1969. Effects of ionizing radiation in children. J. Pediatr. 67:658-73.
- Sutow, W.W. and Conard, R.A. 1969. The effects of fallout radiation on Marshallese children. In: *Radiation Biology in the Fetal and Juvenile Mammal*, pp. 661-73, Sikov, M.R. and Mahlum, D.D. (Editors), USAEC.
- Sutow, W.W., Conard, R.A., and Thompson, K.H. 1982. Thyroid injury and effects on growth and development in the Marshallese children accidentally exposed to radioactive fallout. *Cancer Bull.* **34**:90-6.
- Wakabayashi, T., Kato, H., Ikeda, T., and Schull, W.J. 1983. Studies of the mortality of A-bomb survivors, Report 7: Part III. Incidence of cancer in 1959-1978. Based on the Tumor Registry, Nagasaki. *Radiat. Res.* **93**:112-46.
- Wong, D.C., Purcell, R.H., and Rosen, L. 1979. Prevalence of antibody to hepatitis A and hepatitis B viruses in selected populations of the South Pacific. *Am. J. Epidemiol.* 110:227-36.

Appendix A

Paper presented at the Fourth International Radiopharmaceutical

Dosimetry Symposium, Oak Ridge, Tennessee, November 1985.

THYROID CANCER IN THE MARSHALLESE: RELATIVE RISK OF SHORT-LIVED INTERNAL EMITTERS AND EXTERNAL RADIATION EXPOSURE

Lessard, E.T.,^a Brill, A.B.,^b and Adams, W.H.^b Brookhaven National Laboratory ^aSafety & Environmental Protection Division ^bMedical Department Upton, NY 11973

ABSTRACT

In a study of the comparative effects of internal versus external irradiation of the thyroid in young people, we determined that the dose from internal irradiation of the thyroid with short-lived internal emitters produced several times less thyroid cancer than did the same dose of radiation given externally. We determined this finding for a group of 85 Marshall Islands children, who were less than 10 years of age at the time of exposure and who were accidentally exposed to internal and external thyroid radiation at an average level of 1400 rad. The assumed risk coefficient for children, from external radiation alone, was derived from 1) values in The Effects on Populations of Exposure to Low Levels of Ionizing Radiation: 1980, National Academy Press, 2) values in Report of the Ad Hoc Working Group to Develop Radioepidemiological Tables, National Institutes of Health, and 3) values in Induction of Thyroid Cancer by Ionizing Radiation, National Council on Radi-ation Protection, Report 80. The risk from internal irradiation was computed from dose, health effect results which were reported in a recent BNL study, and an estimate of the external risk coefficient based on other studies. The external risk coefficient ranged between 2.5 and 4.9 cancers per million person-rad-years at risk, and thus, from our computations, the internal risk coefficient for the Marshallese children was estimated to range between 1.0 and 1.4 cancers per million person-rad-years at risk.

In contrast, for individuals more than 10 years of age at the time of exposure, the dose from internal irradiation of the thyroid with short-lived internal emitters produced several times more thyroid cancer than did the same dose of radiation given externally. The external risk coefficients for the older age groups were reported in the above literature to be in the range of 1.0 to 3.3 cancers per million person-rad-years-at risk. We computed internal risk coefficients of 3.3 to 8.1 cancers per million person-rad-years at risk for adolescent and adult groups. This higher sensitivity to cancer induction in the exposed adolescents and adults, is different from that seen in other exposed groups. The small number of cancers (9) in the exposed population and the influence of increased levels of TSH, nonuniform irradiation of the thyroid, and thyroid cell killing at high dose make it difficult to draw firm conclusions from these studies.

INTRODUCTION

The long-term health effects of external thyroid irradiation are known to include excess hypothyroidism, thyroid nodules, and thyroid cancer, and in this study we attempt to quantitate the relative risk of internal irradiation of the thyroid, for induction of thyroid cancer. The effects of external irradiation of child thyroids have been summarized in BEIR III (1) and by the NCRP (2). Internal irradiation of the thyroid from a mixture of radionuclides has occurred in children as a result of accidental exposure to fallout from nuclear weapons testing. Larger numbers of persons having received diagnostic and therapeutic doses from 131 I used in medical applications. Apart from the Marshallese, studies of internally irradiated human populations have not revealed an increased risk of thyroid malignancy (1,2). For example, studies of a group of children exposed to 90,000 person-rad in Utah have not revealed any excess thyroid cancer. The fallout in Utah contained 131 I and was reported to deliver up to several hundred rad of absorbed dose to thyroids of children who were less than 10 years of age (1,2). There are several studies which report no carcinogenic effect from large doses of 131 I (2). For example, Holm reported that persons irradiated with 131 I, with doses ranging between 6000 and 10,000 rad, exhibited no statistically significant increase in thyroid cancer (2). Studies of the children in the Marshall Islands conducted since 1954, on the other hand, do show a statistically significant increase in thyroid cancer in these irradiated subjects. Since the Marshall Islands' children were exposed simultaneously to external and internal irradiation, we have analyzed the data in an attempt to relate each type of exposure, internal versus external radiation, to the observed thyroid health effects. The mixture of radionuclides, contributing to internal dose in the Marshallese, included mostly short-lived ¹³³I and ¹³⁵I, and only 10-20% of the thyroid dose came from ¹³¹I, thus the radiobiological considerations differ greatly in these various exposure circumstances.

Estimates of thyroid-absorbed dose were recently reassessed for people exposed to fallout in the Marshall Islands (3). The accidental exposure of people on March 1, 1954, occurred as a result of nuclear weapons testing. Over the years, several estimates of thyroid-absorbed dose were made (4,5). The earliest estimate of thyroid dose was reported by Cronkite (4) who indicated a population-averaged thyroid dose. A 1962 study by James (5) listed the most probable thyroid dose to girls who were 3 to 4 years old at the time of exposure. However, the James dose estimate was flawed by the incorrect association of 133 I and 135 I dose relative to the dose from 131 I. The most recent assessment of dose provided detailed information on the type of nuclides in fallout, the mode of intake, and the contributions from internal and external sources. The study of Lessard et al. (3) established greater absorbed dose to people based upon greater intake of the shorter-lived radioiodines. The thyroid dose ranged from several hundred to five thousand rad, and the highest doses were assigned to young people. The revised dose estimates accounted for the radioactivity from all iodine isotopes.

Uncertainties with the dose estimates are associated with the amount of radioactivity measured in the urine of the exposed people, the intake of the short-lived radiotellurium and radioiodine isotopes and percent of thyroid uptake as as determined from a physiologic model, errors in estimating the exact amount of each radioiodine isotope, the dose rate and pattern of energy distribution from this radioiodine mixture, and the shape and thickness of the thyroid.

Adams et al. (6) reported the medical status of the Marshallese accidentally exposed to fallout. Through March 1985 there were 35 adenomatous nodules, 5 adenomas, 9 papillary carcinomas, 1 atypical adenoma or follicular carcinoma, and 2 occult papillary carcinomas. A comparison group of equal size exhibited 3 adenomatous nodules, 1 adenoma, 2 carcinomas, and 2 occult papillary carcinomas, one of which may have been a follicular carcinoma. Uncertainty was associated with diagnosis of follicular carcinoma, one in the exposed group and one in the comparison group, because of equally divided opinion among consulting pathologists. However, it was reasoned that both follicular carcinomas could be excluded from a risk coefficient estimate without seriously biasing the results. Diagnoses on five other individuals are pending. All five are from Utirik Atoll; three are in the <10-year old age group, and two are in the 10- to 18-year-old age group.

METHODS

Adams et al. (6) classified thyroid abnormalities following a scheme similar to that used by the World Health Organization and a committee of pathologists who had special expertise in diseases of the thyroid (7). The following nomenclature was used:

Adenomatous nodule: a focal proliferative lesion consisting of changes typical of adenomatous goiter; the lesions do not fulfill criteria of true neoplasms.

Adenoma: an encapsulated proliferative lesion with a uniform internal growth pattern and benign clinical course.

Occult papillary carcinoma: a small nonencapsulated sclerosing carcinoma, considered to be clinically benign even with positive regional lymph nodes.

Papillary carcinoma: larger, infiltrating carcinoma, usually containing both papillary and follicular components. The smallest lesion diagnosed as a papillary carcinoma, by the consultant pathologists, was 0.8 cm in diameter.

The recent computation of thyroid absorbed dose was performed for inhabitants of Rongelap, Utirik, and Ailingnae Atolls who were exposed to fallout on March 1, 1954. The amount of fallout activity taken into the body was estimated from the value of 131 excreted in urine obtained from 64 persons who were at Rongelap. The other components of fallout taken into the body, particularly 133 and 135 L, had to be inferred from studies on fallout composition. The authors of the reassessment study made dose estimates on the basis of actual BRAVO fallout composition. The intake pathway and the time post-detonation at which intake was likely to have occurred were obtained from interviews with the exposed people, and historical records and were factored into the new dose estimates. A detailed development of the dose reassessment was reported by Lessard et al. (3).

The radioepidemiological tables assembled by the Working Group (8) represented the best scientific judgment for the assignment of cancer risk from external radiation; thus we obtained one estimate of external exposure risk coefficient from this source. For persons less than 20 years of age, the Working Group adopted an average risk coefficient of 3.3 excess cancers per million person-rad-years at risk, and for persons 20 years or older they chose a value of 1.0 excess cancer per million person-rad-years at risk. A 10-year minimum latent period was chosen for thyroid cancer. The Working Group calculated thyroid cancer risk based on a linear dose-response function and maintained that the estimates of risk applied to external x and gamma irradiation, but not to the intake of radioisotopes of iodine.

The BEIR III (1) risk coefficients were based, in large part, on external

exposure of children less than 10 years of age, and upon data available through 1979. A central value of 4.0 cancers per million person-rad-years at risk was reported, but after review of their report, we modified the estimate to 4.9 cancers per million person-rad-years at risk. Our result, based on this modification, is discussed in the text and is noted in Table 7. The adjustment was based on weighting the risk coefficient from each study according to the number of excess cancers observed; that is, we gave more weight to cancer risk coefficients developed from studies reporting the greatest number of cancers. The BEIR risk coefficient was based on a minimum latent period of 10 years and on studies involving only external irradiation of the thyroid.

Risk coefficients for external and internal radiation were given in NCRP Report 80 (2), and these coefficients were estimated for a five-year latent period. Report 80 indicated the external risk coefficient applied to ¹³⁵I and ¹³³I intake, but not for ¹³¹I exposure. The two short-lived isotopes of iodine were assumed to have the same effectiveness as x rays, because of the fairly uniform distribution of dose, and because of the comparatively higher dose rates (2). In our analyses, we used risk coefficients for external exposure computed for 5- and 10-year latent periods derived from the following reports. We used external risk coefficients from NCRP Report 80 because they were based on a five-year latent period, and these appear in the results section along with the coefficients developed by the Working Group, which were based on a ten-year latent period.

Risk coefficient estimates, made here, were based on the total external and internal thyroid dose, the total number of cancers, the risk value published for external irradiation of the thyroid, and the partitioning of external and internal dose as follows

$$A B + C D = (A + C)E,$$
 (1)

where

- A = the person-rad to all thyroids from radioisotopes of iodine,
- B = the risk coefficient for internal exposure of the thyroid from radioisotopes of iodine, cancers per person-rad-years at risk,
- C = the person-rad to all thyroids from external gamma radiation,
- D = the risk coefficient from external exposure of the thyroid, for example, 1.0x10⁻⁶ cancers per person-rad-years at risk for adults, or in the case of children <10 years of age, 4.9x10⁻⁶ cancers per person-rad-years at risk, and
- E = the risk coefficient determined from the observed health effects, the total thyroid dose, and the spontaneous rates of thyroid disease in the Marshall Islands subjects. The value of E was computed from Eq. (2-1) given in NCRP Report 80 (2).

Computations of B and E were for latent periods of both 5 and 10 years, since the length of latent period affects the years at risk and the risk coefficient. Years at risk are the period from the end of the latent period to the time cancer is observed in a subject. The value for years at risk strongly affected the computation of risk coefficients.

RESULTS

The data in the Appendix are the result of 31 years of medical and

radiological follow-up and, in the case of cancer diagnosis, of consensus opinion of pathologists. The Appendix is provided to allow others to perform different analyses of the data, recognizing that the data base is incomplete. Verifying the data over the last seven years has resulted in changes in age, identification number, assigned dose, and diagnosis. Several independent groups reported age at exposure, and the Adams et al. (6) version was used here. Different ages at exposure influences the age distribution of cancers, which in turn impacts strongly on the risk coefficient for a given age group.

The external thyroid dose was due to gamma exposure from the fallout cloud and fallout on the ground, and was taken as equal to the external whole-body dose reported by Lessard et al. (3), i.e., 190 rad at Rongelap, 110 rad at Ailingnae, and 11 rad at Utirik.

These external doses were estimated for a point which was 1 meter above the ground, thus some variation in external thyroid dose with a person's height may have occurred. To a first approximation external thyroid dose is inversly proportional to height above the ground. We derived this proportionality by neglecting photon attenuation and buildup, and by limiting the height above ground to between 0.5 and 1.5 meters. The impact on the risk coefficient estimates, relative to assuming that external thyroid dose was height dependent, was minimal, since the person-rad from external exposure was much much less than the person-rad from internal exposure.

The data for the unexposed comparison groups are indicated in Table 1. In the age- and sex-matched comparison group used for this study, two papillary carcinomas have been observed. The summary is completed through 1983. To apply the data for risk coefficient determination, we modified the matched group results by the ratio of 31/29, which corrects for the difference in the number of reported observation years. The larger, less defined comparison population studied by Conard et al. (7) is shown in the first half of Table 1 to show that spontaneous cancer risk is not a strong function of group age for the Marshallese people. The comparison data indicated a spontaneous rate of $3x10^{-4}$ cancers per person-rad-years at risk. A lower spontaneous rate has been reported for the U.S. population, $1x10^{-4}$ per person per year (2). The Marshallese comparison data were used in the risk coefficient computations made here.

A summary of data in the Appendix appears in Tables 2 through 4. Note that out of 9 papillary cancers listed in the Appendix, only 2 were observed in males. This male to female ratio is similar to that reported in other studies (1,2,8). Tables 2 through 4 contain the input data which we used with Eq. (1). The data were grouped in the same manner as in other reports dealing with cancer and radiation exposure of the thyroid. The age groups were the same as that used by Conard et al. (7) and Adams et al. (6). To determine the average years post-exposure to onset of carcinoma, we set onset of carcinoma as the time of clinical observation of a thyroid nodule; thus, a latent period was assumed, but a period of several years could have elapsed before a nodule became large enough for detection by routine palpation by the physician. Therefore, the true latent period could be shorter than that assumed here. Tables 2 through 4 include the expected carcinomas, computed from the age- and sex-matched comparison group, and a summary of the total person-rad from manmade internal and external sources.

Ta	ble	1

Mars			malities in the son Groups 1954.	
Group Age 1954	Number	Total <u>Nodules</u>	Carcinoma	Hypofunction
<10	229	6	2	
10-18	79	6	· 1	1
>18	292	25	2	1
Total	600	37	5	2
Age- and Sem Matched Grow		5	2	
Followed Since 1954				

Summary of Thyroid Abnormalities in the

Table 2

Age Group <10 Data Summary

Number of Persons
Internal Exposure, Person-Rad 120,000
External Exposure, Person-Rad 5400
Number of Observed Carcinomas 3
Average Years Post-Exposure to Onset of Carcinoma
Assumed Latent Period, Years
Number of Expected Spontaneous Carcinomas

Table 3

Age Group 10 to 18 Data Summary

Number of Persons
Internal Exposure, Person-Rad 18,000
External Exposure, Person-Rad 2500
Number of Observed Carcinomas 3
Average Years Post-Exposure to Onset of Carcinoma
Assumed Latent Period, Years 5 and 10
Number of Expected Spontaneous Carcinomas

Table 4

Age Group >18 Data Summary

Number of Persons 120
Internal Exposure, Person-Rad 48,000
External Exposure, Person-Rad 8,000
Number of Observed Carcinomas 3
Average Years Post-Exposure to Onset of Carcinoma
Assumed Latent Period, Years 5 and 10
Number of Expected Spontaneous Carcinomas 1.1

Table 5

Risk Coefficients^a for Marshall Islanders, 10-Year Latent Period

		Excess	Total	Years at	Risk
Group		Thyroid			
<u>Age 1954</u>	Number	Cancers	Person-Rad	<u>Risk</u>	Coefficient
<10	85	2.2	120,000	12.2	1.5x10 ⁻⁶
10-18	32	2.7	21,000	17.7	7.4×10^{-6}
>18	120	1.9	56,000	6.2	5.4×10^{-6}
Total	237	6.8	200,000	11.3	3.0x10 ⁻⁶

^aThyroid cancers per person-rad-years at risk, based on thyroid dose from internal plus external sources.

		Excess		Years	
Group		Thyroid	Total	at	Risk
Age 1954	Number	Cancers	Person-Rad	<u>Risk</u>	Coefficient
<10	85	2.2	120,000	17.2	1.1x10 ⁻⁶
10-18	32	2.7	21,000	22.7	5.8x10 ⁻⁶
>18	120	1.9	56,000	11.2	3.0x10 ⁻⁶
Total	237	6.8	200,000	14.9	2.3x10 ⁻⁶

Risk Coefficients^a for Marshall Islanders, 5-Year Latent Period

^aThyroid cancers per person-rad-years at risk, based on thyroid dose from internal plus external sources.

Table 7

Estimated Risk Coefficient^a for Internal and External Exposure

		10-Year Latent Period		5-Year Latent Period	
		External	Internal	External	Internal
Group		Risk	Risk	Risk	Risk
<u>Age 1954</u>	Number	Coefficient	Coefficient	Coefficient	Coefficient
<10	85	3.3x10 ⁻⁶	1.4x10 ^{-6(b)}	2.5x10 ⁻⁶	1.0x10 ⁻⁶
10-18	32	3.3×10^{-6}	8.0x10 ⁻⁶	2.5x10 ⁻⁶	6.3×10^{-6}
>18	120	1.0×10^{-6}	6.1x10 ⁻⁶	1.3x10 ⁻⁶	3.3x10 ⁻⁶
Total	237	2.1x10 ⁻⁶	4.7x10 ⁻⁶	1.9x10 ⁻⁶	2.9x10 ⁻⁶

^aThyroid cancers per person-rad-years at risk.

^bA value of 1.3×10^{-6} results when 4.9×10^{-6} is used for the external risk coefficient.

The risk coefficient, E, for different age groups, computed from total dose resulting from internal plus external exposure for Marshall Islanders, ranged from 1.5×10^{-6} to 7.4×10^{-6} per person-rad-years at risk, assuming a 10-year latent period, and 1.1×10^{-6} to 5.8×10^{-6} , assuming a 5-year latent period. These data are indicated in Tables 5 and 6, respectively. The total risk coefficient, E, was used in Eq. (1) to determine the internal risk coefficient, B. For external risk coefficients and 10-year latent period, we chose 3.3×10^{-6} for age < 20 and 1.0×10^{-6} for age > 20 based on the Working Group study (8); for 5-year latent period we chose 2.5×10^{-6} for age < 18 and 1.3×10^{-6} for age >18, based on NCRP Report 80 (2). The results for internal risk coefficients are in Table 7. Finally, as we explained in the Methods, we chose a special value for the <10-year age group, since it was based on a large group of children exposed to x rays (1). This value was 4.9×10^{-6} cancers per person-rad-years at risk, and the estimate for the internal risk coefficient was 1.3×10^{-6} , virtually the same as the value given in Table 7 for the 10-year latent period.

A tabulation of risk coefficient versus internal thyroid dose is given in Table 8. These internal dose groupings resulted in little variation in external dose as a function of age. These groupings were made to examine the affect of dose on the value for internal risk coefficient.

Table 8

Average Dose Versus Internal and External Risk Coefficients, 10-Year Latent Period

	Average		Average			
	Internal	Internal	External	External	Total	
Group	Thyroid	Risk	Thyroid	Risk	Risk	
Age 1954	Dose, rad	Coefficient ^a	Dose, rad	<u>Coefficient^b</u>	<u>Coefficient^a</u>	
<10	1400	1.4×10^{-6}	63	3.3x10 ⁻⁶	1.5x10 ⁻⁶	
10-18	560	8.0x10 ⁻⁶	78	3.3×10^{-6}	7.4×10^{-6}	
>18	400	6.1x10 ⁻⁶	66	1.0x10 ⁻⁶	5.4x10 ⁻⁶	

^aThis study.

^bReference 8.

A sensitivity analysis, of the parameters in Eq. (1), shows that the value for the total risk coefficient, E, impacts greatly on the estimate of the internal risk coefficient, B, in this specific Marshall Islands study. This is because of the wide difference between internal thyroid dose, A, and external thyroid dose, C. Thus, our estimate of internal risk coefficient depends largely on the observed incidence of thyroid cancer because the total risk coefficient, E, is very sensitive to the small number of spontaneous and excess thyroid cancers observed.

DISCUSSION/CONCLUSION

Interest in the relative risk of ¹³¹I taken internally and external radiation dose to the thyroid relates to radiation protection and medical care issues. Unfortunately for those interested in obtaining information on this important issue, the complex mixture of radionuclides taken up by the Marshallese precludes such an analysis. The results obtained for these studies are specific to the case where the thyroid dose was due to a mixture of shortlived radioisotopes of iodine, some of which were produced by the decay of tellurium within the body. Current information on animal and human data was summarized recently in NCRP Report 80 (2). The Committee concluded that was less then one third as effective for thyroid cancer induction as external radiation. This can not be compared directly to the results of the present study because of the small amount of 131 I in the Marshallese exposures. In study because of the small amount of the time time time the small studies, which used rodents, high TSH levels were found to be 131 The solutions of the solution of the solut necessary co-factors for thyroid cancer induction. Thus, goitrogen plus exposures were needed to induce thyroid cancer, except in several studies using Long-Evans rats which behaved differently from all other strains studied. Results of ¹³¹I treatment of children for hyperthyroidism were reported in two large studies. In reviewing results of treatment of nine children, Sheline et al. (9) found that all of them subsequently developed thyroid nodules and one was diagnosed as having of thyroid cancer, about which there was disagreement regarding pathology. None of those children received thyroid replacement therapy after 131 I treatment, and all presumably developed high endogenous TSH levels. In Los Angeles, at a later date, 73 children were treated with approximately the same ¹³¹I dose, all were placed on thyroid replacement, and none developed thyroid nodules (10). Thus the relative risk of thyroid dose from internal emitters compared to external radiation for Marshall Islanders may be influenced by a high TSH co-factor, since thyroid replacement therapy began 11 years after exposure. Replacement therapy was recommended only for the high-dose group which, at that time, was thought to be the people at Rongelap.

Also no increased incidence of thyroid cancer was seen in large numbers of human subjects exposed to similar or higher doses of ¹³¹I in the treatment of thyrotoxicosis (11), or in children given ¹³¹I in lower diagnostic doses (12).

Hypothyroidism is a nonstochastic effect of ionizing radiation exposure, with estimated threshold for induction of 2000 rad to the thyroid (1). In the Marshallese children, whose thyroids were exposed to doses in the several thousand rad range, hypothyroidism and increased TSH levels certainly existed in the early years following exposure. In later years, uneven acceptance of thyroid supplementation by children may have led to persistent increased TSH levels. The combination of high TSH and high internal and external radiation doses may account for the unusually high incidence of nodules in this population, and in the unusual age distribution of sensitivity.

The numbers of individuals in the study are small, and statistical segregation of the interacting factors is not possible. Thus, it will be difficult to draw precise conclusions from this study with respect to apportionment of risk between internal and external doses. Further, the differences between the radiological characteristics of 131, 133, and 135 and the larger doses from 133 and 135 make it difficult to assess the relative risk of 131 and external radiation in this circumstance. A simple statistical model was used (3) to indicate the one sigma confidence interval. This confidence interval is indicated in the following paragraph in parentheses. The standard deviation of the risk estimate, E, was 1.5 times the average value for the risk estimate, and development of this standard deviation was given by Lessard et al. (3). The results support the notion that external risk coefficients are different from internal risk coefficients following exposure to a mixed radiation field. The total risk coefficients $[3.0x10^{-6} (\pm 4.5x10^{-6}) \text{ cancers}$ per person-rad-year at risk, 10-year latent period, and $2.3x10^{-6} (\pm 3.5x10^{-6})$ cancers per person-rad-year at risk, 5-year latent period] are similar to the literature values (1,2) for this age distribution and for external exposure. The literature values are $2.1x10^{-6}$ for a 10-year latent period and $1.9x10^{-6}$ for a 5-year latent period. However, if the risk is examined as a function of age or as a function of dose, differences are encountered. For example, the ratio of the risk coefficient for external exposure to the risk coefficient for internal exposure, in the <10 year age group, is 2.5 (0.38 to 4.6). In the 10- to 18-year age group, this risk coefficient ratio is 0.40 (0.22 to 2.6).

Small group size, in this study, and the uncertainties reported in studies on medical and fallout exposures make it difficult to establish relative risks of thyroid cancer from internal and external radiation doses to the thyroid. The possible synergistic effect of internal and external exposures and the modifying factors such as high TSH levels and nonuniform irradiation of thyroid cells complicate the biological interpretation of the risk. In this study, different age groups correspond to different dose levels, and very high dose to the thyroid may be a significant modifying factor. Because of the high interest in evaluating human sensitivity to ¹³¹I, continued efforts are needed to obtain data and to conduct analyses that will establish better estimates of risk coefficients than are now available. It is not likely that data for the Marshallese exposures will contribute to the answer to that important question.

ACKNOWLEDGMENT

The authors fully appreciate the efforts of John Baum, Associate Head for Research in the Safety and Environmental Protection Division, for his review of this manuscript.

This work was performed by members of Brookhaven National Laboratory, Associated Universities, Incorporated, under contract No. DE-ACO2-76CH00016 with the United States Department of Energy, and under contract No. Y01-CP-40503 with the National Cancer Institute.

REFERENCES

- Committee on the Biological Effects of Ionizing Radiation, The Effects on Populations of Exposure to Low Levels of Ionizing Radiation: 1980, National Research Council Report, National Academy Press, Washington, DC, 1980.
- Recommendations of the National Council on Radiation Protection and Measurements, Induction of Thyroid Cancer by Ionizing Radiation, NCRP Report No.80, Bethesda, MD, 1985.
- Lessard, E.T., R.P. Miltenberger, R.A. Conard, S.V. Musolino, J.R. Naidu, A. Moorthy, and C.J. Schopfer, Thyroid Absorbed Dose for People at Rongelap, Utirik and Sifo on March 1, 1954, Brookhaven National Laboratory Report, Upton, NY, BNL 51882, 1985.

- 4. Cronkite, E.P., V.P. Bond, and C.L. Dunham, A Report on the Marshallese and Americans Accidentally Exposed to Radiation from Fallout and a Discussion of Radiation Injury in the Human Being, in: Some Effects of Ionizing Radiation on Human Beings, U.S. Atomic Energy Commission, Washington, DC, USAEC-TID 5358, 1956.
- 5. James, R.A., Estimate of Radiation Dose to Thyroids of Rongelap Children Following the BRAVO Event, Lawrence Radiation Laboratory Report, Livermore, CA, UCRL-12273, 1964.
- Adams, W.H., J.A. Harper, R.S. Rittmaster, P.M. Heotis, and W.A. Scott, Medical Status of the Marshallese Accidentally Exposed to BRAVO Fallout Radiation: January 1980 Through December 1982, Brookhaven National Laboratory Report, Upton, NY, BNL 51761, 1982.
- Conard, R.A. et al. (9 co-authors), Review of Medical Findings in a Marshallese Population Twenty-Six Years After Accidental Exposure to Radioactive Fallout, Brookhaven National Laboratory Report, Upton, NY, BNL 51261, 1980.
- Ad Hoc Working Group to Develop Radioepidemiological Tables, Report of the National Institutes of Health, January 4, 1985. Office of the Director, NIH Publication No. 85-2748. U.S. Dept. HHS, PHS, VIH.
- 9. Sheline, G.E. et al., "Thyroid Nodules Occurring Late After Treatment of Thyrotoxicosis with Radioiodine," J. Clin Endoc. & Metab. 22, 8, 1962.
- Starr, P., "Later Results of I-131 Treatment of Hypothyroidism in 73 Children and Adolescents: 1967 Follow-up," J. Nucl. Med. 10, 586-590, 1969.
- 11. Dobyns, B.M. et al., "Malignant and Benign Neoplasms of the Thyroid in Patients Treated for Hypothyroidism: A Report of the Cooperative Thyrotoxicosis Follow-up Study," J. Clin, End. & Metab. 38, 976-998, 1974.
- 12. Holm, L.E. et al., "Incidence of Malignant Thyroid Tumors in Humans After Exposure to Diagnostic Doses of I-131," J. National Cancer Institute, 64, 1055-1059, 1980.
- 13. Prentice, R.L. et at., Radiation Exposure and Thyroid Cancer Incidence among Hiroshima and Nagasaki Residents, National Cancer Institute Monograph 62, 207-212, 1982.
- 14. Ishimary, T. et al., Incidence of Thyroid Cancer Among Atomic Bomb Survivors and Controls Hiroshima and Nagasaki, 1958-79: Relationship to Radiation Absorbed Dose in the Thyroid, RERF Technical Report 1984.

APPENDIX

Tabulation of Thyroid Dose and Thyroid Health Effects

ID Number	Sex	Age in 1954	Comment	Diagnosis	Internal Thyroid Dose, Rad	Years Post Exposure
*1	F	52	Died 1985		290	
2	М	1		Adenomatous Nodule	5000	11
3	м	1		Myxedema	5000	
4	м	36			1000	
5	М	1		Myxedema	5000	
*6	м	1		•	1300	
7	М	34			1000	
*8	F	5		Adenomatous Nodule	740	18.5
9	M	20			1000	1013
10	M	22			1000	
11	м	48			1000	
12	F	16			1200	
13	F	59	Died 1966		1100	
14	F	3	bicd 1900		3500	
15	F	5	Surgery(2x)	Adenomatous Nodule	2800	22;32
*16	M	37	Jurgery (2x)	Adenomatous	280	22,34
17	F	1		Adenomatous Nodule	5000	10.5
18	F	19		Papillary Carcinoma	1100	15.5
19	M	3		Adenomatous Nodule	3500	14.5
20	M	5		Adenomatous Nodule	2800	14.5
21	F	1		Adenomatous Nodule	5000	10.5
22	F	15		Adenomatous Nodule	1300	10.5
23	M	2		Adenomatous Nodule	4000	14.5
24	F	11		Adenomatous Nodule	1700	14.5
25	M	44	Died 1956		1000	
26	M M	13	Died 1962		1500	
20	M	33	Died 1962			
*28	r F	69	Died 1965		1000	
*29	M	65	Died 1965		290	
30	F	52	Died 1968		280 1100	
*31	г М	31	Died 1958		280	
32	M	2	Died 1990		4000	
33	F	1		Adenomatous Nodule	5000	12
34	F	43		Adenoidatous Module	1100	
35	r M	11			1700	
36	M	5		Adenomatous Nodule	2800	15.5
37	M	18		Adenomatous Nodule	1000	• •
37	л М	75	Diad 1957		1000	
38	л F	13	Died 1957		1500	
39 40	-				1000	
40 *41	M M	31 42			290	
42	r F			Adamamakan Madula	5000	
44	r	1		Adenomatous Nodule	290	

Rongelap and Ailingnae Population

31

ID Number	Sex	Age in 1954	Com	ment	Diagnosis	Internal Thyroid Dose, Rad	Years Post Exposure
*44	м	2					
*45	F	30			Adenomatous Nodule	290	19
46	М	76	Died	1962		1000	
47	м	6				2400	
*48	F	4				820	
49	F	13				1500	
*50	М	34	Died	1971		280	
*51	F	23	Died	1982	Follicular Adenoma	290	20
52	F	46	Died	1963		1100	
*53	F	5			Adenomatous Nodule with Occult Papillary Carcinoma	740	27
54	м	1	Died	1972	Adenomatous Nodule	5000	14.5
55	Я	76	Died			1000	
56	F	67	Died			1100	
57	F	98	Died			1100	
58	F	59	Died			1100	
*59	F	44	Died		Adenomatous Nodule	290	12
60	F	56	Died		Adenomatous Addute	1100	12
61	F	6	DIĘU	1972	Adenomatous Nodule	2400	12
62	F	55	Died	1959	Adenomatous Noulle	1100	12
63	F	34	preu	1937		1100	
64	F	28			Papillary Carcinoma	1100	11
65	F	1			Adenomatous Nodule	5000	12
66	F	29			Adenomatous Nodule	1100	25.5
67	F	12				1600	31
68	r M	44	Diad	1974	Papillary Carcinoma	1000	21
69	F	2	pred	19/4	Adenomatous Nodule	4000	10.5
*70	r F	5			Adenomatous Nodule	740	10.5
71	F	26		•		1100	
72	r M	5			Parillar Constant	2800	15.5
73	M	16			Papillary Carcinoma	1200	10.0
74	F	14			Partillary Constants		22
75	r F	10			Papillary Carcinoma Adenomatous Nodule	1400	18.5
61	r	10				1800	18.5
76	м	9			with Follicular Adenor	na. 2000	
70	M	24				1000	
78	л F	24 35				1100	
79	r M	35					
80	т М	37 44	NI - 2	1983		1000	
80 *81	л F	44 6	pred	1907		1000	
82	r M	49	D4 - 2	1000		640	
82	м м	49 In Utero		1980	Adapamakana Madula	1000	20
00	m M	In Utero			Adenomatous Nodule		20

Rongelap and Ailingnae Population

			Rongelap a	nd Ailingnae Population		
ID Number	Sex	Age in 1954	Comment	Diagnosis	Internal Thyroid Dose, Rad	Years Post Exposure
85 86	M F	In Utero In Utero		Adenomatous Nodule		25.5

*Ailingnae Exposed

			Ut	irik Population		
2101	м	48	Died 1968		150	
2102	M	3			480	
2103	М	43			150	
2104	F	22			160	
2105	м	45			150	
2106	M	4			430	
2107	F	25			160	
2108	М	11			250	
2109	F	45	Died 1978		160	
2110	М	47			150	
2111	F	6			340	
2112	м	53	Died 1968		150	
2113	F	3			480	
2114	м	40			150	
2115	м	1			670	
2116	F	21	Died 1960		160	
2117	F	24			160	
2119	F	18			160	
2120	М	4	Died 1982		430	
2121	м	57	Died 1965		150	
2122	м	82	Died 1959		150	
2123	м	15			200	
2124	м	2			550	
2125	M	37			150	
2126	F	5			390	
2127	М	68	Died 1959		150	
2128	F	8	Died 1985		310	
2129	F	17			160	
2130	F	3			480	
2131	F	29	Died		160	
2132	F	1		Adenomatous Nodule	670	27
2134	F	1			670	
2135	м	31	Died 1977		150	

Utirik Population

	Utirik Population												
ID Number	Sex	Age in 1954	Comment	Diagnosis	Internal Thyroid Dose, Rad	Years Post Exposure							
2136	м	3			480								
2137	M	14			220								
2138	F	4			430								
2139	F	44			160								
2140	F	45			160								
2141	F	53	Died 1968		160								
2142	M	5	Dica 1900		390								
2143	M	3			480								
2144	M	7			330								
2145	M	34			150								
2145	F	36	Died 1980		160								
2140	F	5	Died 1960	Adenomatous Nodule	390	25.5							
2147	r M	44		Adenouatous Noture		23.3							
2148	F	9		Disease Desides	150	20							
				Diagnosis Pending	300	30							
2150	M	10		Watte 1 41	270								
2150	М	12		Follicular Adenoma	240	22							
2151	F	4			430								
2152	M	17		Papillary Carcinoma	150	30							
2153	M	1			670								
2154	F	40	Died 1965		160								
2155	м	1			670								
2156	М	8			310								
2157	м	26	Died 1984		150								
2158	F	28			160								
2159	F	3			480								
2160	F	4		Papillary Carcinoma	430	21							
2161	F	29	Died 1981		160								
2162	F	32			160								
2163	М	65	Died 1964-65?		150								
2164	F	7	Died 1984		330								
2165	М	11			250								
2166	M	38			150								
2167	M	14			220								
2168	м	18	Died 1984	Diagnosis Pending	150	30							
2169	м	62	Died 1978		150								
2170	м	41	Died 1959		150								
2171	F	2		Papillary Carcinoma	550	30							
2172	F	12		Diagnosis Pending	240	30							
2174	М	1		- 5	670								
2175	M	57	Died 1970		150								
2176	М	10			270								
2177	м	5	Died 1961		390								
2178	М	19	Died 1972		150								
2179	м	2			550								
2180	M	70	Died 1960		150								

......

ID Number	Sex	Age in 1954	Com	ment	Diagnosis	Internal Thyroid Dose, Rad	Years Post Exposure
2181	м	65	Died	1967		150	
2182	F	52				160	
2183	м	56	Died	1965		150	
2184	М	60	Died	1961		150	
2185	м	32	Died	1984		150	
2187	F	56	Died	1959		160	
2188	м	3				480	
2189	F	26				160	
2190	F	75	Died	1964-65?		160	
2191	F	75	Died	1969		160	
2192	F	74	Died	1964-65?		160	
2193	F	31			Adenomatous Nodule	160	25
2194	F	35	Died	1984	Papillary Carcinoma	160	22
2195	F	24			Adenomatous Nodule	160	25
2196	F	38			Adenomatous Nodule	160	26.5
2197	F	3			Diagnosis Pending	480	31
2198	F	58	Died	1979		160	
2199	F	42	Died	1961		160	
2200	F	43				160	
2201	F	50	Died	1974		160	
2202	F	59	Died	1967		160	
2203	F	62	Died	1963		160	
2204	F	60	Died	1965		160	
2205	М	29				150	
2206	М	32				150	
2207	М	5				390	
2208	F	37			Adenomatous Nodule	160	19
2209	F	5				390	
2210	F	1				670	
2212	F	34			Adenomatous Nodules	160	19
2213	F	1				670	
2214	М	65	Died	1969		150	
2215	М	1			Adenomatous Nodule	670	25.5
					with Occult Papillary Carcinoma		
2216	F	33				160	
2217	F	22				160	
2218	F	1				670	
2219	F	54	Died	1957		160	
2220	F	25				160	
2221	F	52			Adenomatous Nodules	160	19
2222	F	60	Died	1957		160	
2223	F	66	Died	1967		160	
2224	F	31				160	
2225	F	6			Diagnosis Pending	340	30

Utirik Population

ID Number	Sex	Age in 1954	Com	ment	Diagnosis	Internal Thyroid Dose, Rad	Years Post Exposure
2226	F	1				670	
2227	F	4				430	
2228	F	8				310	
2229	F	18			Follicular Carcinoma	160	15.5
	-				Possible Atypical Ade		
2230	F	13				230	
2231	F	1				670	
2232	м	1				670	
2234	M	12				240	
2235	M	7				330	
2236	M	11			Follicular Adenoma	260	24
2237	M	7			POTTICUTAL MENOMA	330	24
2238	F	, 54	Died	1965		160	
2239	F	3	pred	1905	Adenomatous Nodule	480	27
2240	M	33	Died	1077	Adenomacous Noture	150	27
2240	F	28	Died			150	
2241	r M	1	pred	1901		670	
2242		46	D 4 - 4	1050			
	M		Died	1928		150	
2245	M	1	N / - 1	1071		670	
2246	F	8	Died	1971		160	
2247	F	8				310	
2248	F	15			Occult Papillary Carcinoma	200	29
2249	F	15				200	
2250	М	10				270	
2251	F	4				430	
2252	м	39	Died			150	
2253	М	45	Died	1965		150	
2254	F	5	•			390	
2255	F	1				670	
2256	F	5				390	
2257	М	7				330	
2258	М	47	Died	1971		150	
2259	F	21	Died	1968		160	
2260	F	1				670	
2261	M	26				150	
2268	М	In Utero	1				
2269	М	In Utero)				
2271	М	In Utero	1				
2273	м	In Utero	•				
2274	М	In Utero	•				
2276	м	In Utero					
2277	F	In Utero					
2548	М	In Utero)				

Utirik Population

Appendix B

Individual Marshallese laboratory data collected during the 1983 and 1984 medical surveys.

а

2.

Abbreviations:

IDN = Brookhaven National Labora	atory identification number
WBC = leukocyte count/ μ l	· · · · · · · · · · · · · · · · · · ·
PMN = neutrophil count/µl	TSH = thyroid stimulating hormone
BND = band forms/µl	level in µU/l
LYM = lymphocytes/µl	PRL = serum prolactin in ng/ml
MON = monocytes/µl	HBS = hepatitis B surface antigen
EOS = eosinophils/µl	AHBS = antibody to hepatitis B
BAS = basophils/µl	surface antigen
PLT = platelet count X $10^3/u1$	AHBC = antibody to hepatitis B core
HCT = percent	antigen
RBC = erythrocytes X $10^6/\mu l$	HDL = high-density lipoprotein in
MCV = mean corpuscular volume	mg/dl
in fl	CHO = cholesterol in mg/dl
HGB = hemoglobin level in g/dl	TRI = triglyceride in mg/dl

Comments:

 Identification numbers 1 to 86 belong to exposed persons of Rongelap and Ailingnae; numbers beginning at 2102 belong to the Utirik exposed; numbers from 805 through 1578 belong to the Comparison group.

С

- 2. Entries containing only 9s indicate no data were obtained.
- 3. Most normal ranges of the indicated tests are given in text. The value of 0.0 for TSH means the level was $< 2.5 \mu$ U/ml, (i.e., not elevated). Codes for HBS, AHBS, AHBC are 0, 1, 9, which indicate, respectively, not present, present, and not performed.

IDN	WBC	P M N	BND	LYM	MON	EOS	BAS	PLT	нст	RBC	MCV	HGB	TSH
1	6200	2852	62	2418	372	496	ø	198	43.4	4.37	99	14.2	ø.ø
2	67ØØ	3Ø15	134	268Ø	2Ø1	6Ø3	67	212	44.3	4.68	95	15.8	ø.ø
3	89ØØ	48Ø6	89Ø	24Ø3	356	45	ø	356	48.9	5.57	88	15.8	3.2
4	7400	3552	296	296Ø	222	3ØØ	ø	236	49.6	5.36	93	16.1	4.4
5	77.00	4466	154	1925	462	616	ø	249	44.4	4.39	1Ø1	14.Ø	152.Ø
6	48ØØ	1872	48	22Ø8	144	432	ø	237	43.5	4.39	99	14.1	ø.ø
7	6000	192Ø	Ø	342Ø	18Ø	48Ø	ø	252	43.Ø	4.34	99	14.Ø	5.6
8	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
9	6300	2961	ø	2898	315	126	ø	256	45.Ø	4.67	96	15.7	2.5
	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999		999.9
iĩ	5900	3422	118	1829	354	118	59	183	32.9	3.24	102	10.8	ø.ø
12	83ØC	415Ø	166	2739	415	93Ø	ø	400	40.8	4.18	98	13.9	3.Ø
14	58 <i>0</i> 0	2726	116	2494	29Ø	116	ø	337	40.8	4.Ø4	101	13.2	ø.ø
15	10500	4725	105	483Ø	63Ø	21Ø	õ	366	42.9	4.84	89	14.3	10.3
16	4300	2494	43	1462	129	172	õ	248	46.7	5.79	81	14.1	4.1
17	95ØØ	5985	855	1805	57Ø	190	95	251	41.4	4.5Ø	92	14.2	ø.ø
18	999999	99999	9999	9999	9999	9999	999	9 99	99.9	9.99	999	99.9	999.9
19	7000	455Ø	28Ø	14ØØ	35Ø	35Ø	7Ø	351	46.6	5.84	-8ø	15.5	80.0
19	7000 5300			2385	350	53	, io Ø	381	40.0	5.68	87	16.9	8.5
2Ø	42ØØ	2385 2184	159 Ø		318 252	53 84	Ø	200	49.2	4.94	89	14.Ø	8Ø.Ø
21	4200 5900	2065	236	1638		767	Ø	324	43.9 39.3	4.94 4.0Ø	98	13.4	31.Ø
22	10300			2065	177 412	515	ø	324	49.6	4.00	94	15.4	31.0
23	61ØC	4841 2745	3Ø9 61	4223		610	61	349	49.6	5.28 4.75	94 95	14.3	16.Ø 3.6
24				2257	427					4.75			3.0
27	79ØØ	3713	316	3239	474	79	79	186	5Ø.4	4.96	102	15.9 99.9	Ø.Ø 999.9
32	999999 9øøø	99999	9999	9999	9999	9999	999	999 438	99.9	9.99	999 84	13.4	333.3
33		504Ø	18Ø	1710	54Ø	630	ø		43.7	5.18			5.3
34	7300	2555	365	3942	219	365	ø	335	39.2	3.60	1Ø9	12.5	ø.ø
35	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
36	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
37	7200	36ØØ	142	2592	288	432	144	2Ø1	46.5	4.73	98	15.3	ø.ø
39	6500	3445	195	2080	39Ø	325	65	444	44.1	4.55	97	12.8	ø.ø
4Ø	6500	377Ø	195	1820	325	39Ø	ø	331	37.3	3.75	99	12.Ø	ø.ø
41	6100	2867	Ø	2257	366	549	61	221	45.5	4.44	100	14.8	3.9
42	8100	3969	324	2754	486	567	ø	263	43.3	4.20	1Ø3	14.Ø	10.9
44	84ØØ	4Ø32	336	3024	756	252	Ø	4Ø9	49.3	5.6Ø	88	15.5	ø.ø
45	7000	518Ø	21Ø	133Ø	21Ø	7Ø	ø	437	4Ø.5	4.3Ø	94	13.3	ø.ø
47	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
48	9999 9	99999	9999	9999	9999	9999	999	99 9	99.9	9.99	999	99.9	999.9
49	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
51		99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
53	10500	5565	315	3255	84Ø	525	ø	464	42.2	4.27	99	13.8	ø.ø
61	99ØØ	4653	ø	4752	198	297	ø	3Ø3	48.Ø	5.42	89	16.4	16.5
63	76ØØ	41Ø4	76	266Ø	45Ø	3Ø4	ø	3ØØ	43.8	4.55	96	14.Ø	ø.ø
64	99 999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
65	6300	3528	378	1323	567	5Ø4	ø	452	3Ø.5	3.35	91	9.4	55.8
66	11400	7638	798	285Ø	114	ø	ø	31Ø	4Ø.2	4.11	98	13.7	4.Ø
67	75ØØ	36ØØ	зøø	3000	225	375	ø	268	44.7	4.41	1Ø1	14.3	ø.ø
69	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
7Ø	4000	2Ø4Ø	ø	116Ø	12Ø	68Ø	ø	32Ø	4Ø.Ø	4.48	89	13.2	ø.ø
71	74ØØ	3774	37Ø	2516	296	444	ø	377	39.Ø	4.02	97	13.1	5.Ø

.

IDN	₩BC	PMN	BND	LYM	MON	ÉOS	BAS	PLT	нст	RBC	MCV	HGB	TSH
72	10200	5212	2Ø4	3264	4Ø8	51Ø	1Ø2	454	45.5	4.8Ø	95	14.8	48.2
73	71ØØ	497Ø	71	1775	142	142	Ø	244	5Ø.1	5.29	95	16.2	ø.ø
74	13900	82Ø1	139	4Ø31	417	1112	ø	324	48.4	5.22	93	15.8	ø.ø
75	8400	46Ø2	168	26Ø4	168	84Ø	ø	33Ø	39.5	4.31	92	13.6	15.1
76	7100	2414	71	4047	71	426	71	275	46.7	4.83	97	15.8	Ø.Ø
77	74ØØ 66ØØ	5254	74Ø 66	1184	74 33Ø	143 198	ø ø	3Ø7 325	46.9	5.1Ø 4.48	92 98	15.1 14.Ø	Ø.Ø 2.5
78 79	57ØØ	3762 342Ø	57	2244 1938	330	198 Ø	Ø	152	51.2	4.48	1ØØ	14.Ø 16.Ø	2.5 Ø.Ø
81	5700 6000	276Ø	18ø	216Ø	300	78ø	้อั	348	36.5	4.38	88	13.5	ø.ø
83	95ØØ	361Ø	285	418Ø	57Ø	76Ø	õ	359	49.4	5.Ø6	98	16.3	ø.ø
84	46 <i>0</i> 0	1932	46	2208	276	138	õ	375	49.6	4.98	100	16.1	999.9
85	9400	4324	376	376Ø	282	658	ø	3Ø1	53.3	5.66	94	16.4	ø.ø
86	88ØØ	6512	264	176Ø	88	176	ø	261	33.5	3.45	97	10.9	ø.ø
8Ø5	99999	99999	9999	9999	9999	9999	999	999.	99.9	9.99	999	99.9	999.9
811	9600	5184	576	3264	96	384	96	251	37.1	3.83	97	13.3	ø.ø
812	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
813	6600	2574	132	297Ø	33Ø	594	ø	324	47.6	4.68	1Ø2	16.1	999.9
814	81ØØ	2997	ø	3888	4Ø5	81Ø	Ø	262	5Ø.3	5.29	95	16.7	999.9
\$15	7100	355Ø	Ø	284Ø	284	355	Ø	347	49.6	5.2Ø	95	16.Ø	999.9
816	6800	3876	34Ø	1768	272	544	Ø	355	38.6	4.34	89	12.9	999.9
817	11100	5772	222	3885	888	333	Ø	274	52.Ø	5.33	98	17.2	999.9
818	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9 16.3	999.9
82Ø 821	85ØØ 99999	3025 99999	34Ø 9999	323Ø 9999	68Ø 9999	425 9999	ศ 1999	336 999	54.1 99.9	5.48 9.99	99 999	99.9	99.9 999.9
822	4900	1225	392	2842	294	147	555 Ø	2Ø5	48.5	5.28	92	15.Ø	999.9
823	4500	2385	9Ø	1665	100	165	45	254	44.4	4.79	93	15.6	999.9
825	6600	3234	Ĩø	2046	264	264	ี้ø	381	43.7	5.00	87	13.6	999.9
826	5300	2809	265	159Ø	212	371	õ	281	39.8	4.23	94	14.Ø	999.9
827	8400	4368	252	3Ø24	42Ø	252	84	285	45.6	4.66	98	14.6	999.9
829	6600	3Ø36	ø	3Ø36	396	66	66	999	42.4	4.52	94	14.Ø	ø.ø
83Ø	86ØØ	559Ø	172	2236	172	43Ø	ø	336	44.7	4.75	94	15.6	999.9
831	7400	259Ø	74	3848	444	296	148	298	46.3	4.81	96	15.3	999.9
832	7 2Ø Ø	238Ø	36Ø	3672	72	216	ø	329	39.8	4.62	86	13.3	999.9
833	46ØØ	1886	92	2162	23Ø	230	Ø	262	46.2	5.29	87	15.3	999.9
834	7600	418Ø	228	266Ø	456	76	ø	212	49.1	5.42	91	16.Ø	999.9
835	11800	6962	236	3422	354	826	Ø	2,77	42.6	4.35	98	14.8	999.9
836	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
838 839	99999 99999	99999 99999	9999 9999	9999 9999	9999 9999	9999 9999	999	999 999	99.9 99.9	9.99 9.99	999 999	99.9 99.9	999.9 999.9
84Ø	8100	3726	ووور لا	3Ø78	487	729	999 81	356	48.5	5.86	83	15.8	999.9
841	10500	7245	315	2205	630	105	ø	2Ø5	40.J 43.Ø	4.75	91	14.3	ø.ø
842	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
843	7500	้รวัดดี	255	225ø	375	45Ø	Ĩø	249	37.7	3.9ø	97	13.3	ø.ø
844	9000	406Ø	36Ø	3060	360	360	õ	275	44.5	4.56	98	14.0	999.9
845	7500	345Ø	225	3375	225	15Ø	75	299	46.4	5.00	93	14.4	999.9
846	10900	6758	874	25Ø7	436	327	ø	374	42.2	4.36	97	13.8	999.9
85Ø	99999	99999	9999	99 99	9999	9999	9 99	999	99.9	9.99	999	99.9	999.9
851	66ØØ	4Ø26	66	231Ø	66	132	ø	278	39.5	3.92	1Ø1	13.2	999.9
855	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
863	72ØØ	2808	144	3Ø24	432	288	ø	262	49.7	4.92	101	16.4	999.9

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	НСТ	RBC	MCV	HGB	TSH
864 865 867 868 869	66ØØ 63ØØ 99999 99999 99999	29Ø4 2835 99999 99999 99999	132 315 9999 9999 9999	27Ø6 2394 9999 9999 9999	198 189 9999 9999 9999	66Ø 567 9999 9999 9999	Ø 63 999 999 999	275 274 999 999 999	41.9 40.6 99.9 99.9 99.9	4.56 4.27 9.99 9.99 9.99	92 95 999 999 999	14.3 14.1 99.9 99.9 99.9	999.9 999.9 999.9 999.9 999.9
878 879 880 881	99999 99999 84ØØ 67ØØ	99999 99999 5376 2881	9999 9999 5Ø4 134	99999 9999 1848 335Ø	9999 9999 5Ø4 268	9999 9999 168 67	999 999 Ø	999 999 5Ø3 215	99.9 99.9 31.0 47.8	9.99 9.99 3.16 4.98	999 999 98 98	99.9 99.9 11.Ø 16.Ø	999.9 999.9 999.9 999.9
882 883 888 891	8500 8700 7600 99999	5525 2871 4636 99999	85 435 152 9999	2Ø4Ø 435Ø 22Ø4 9999	255 435 228 9999	51Ø 6Ø9 3Ø4 9999	85 Ø 76 999	315 27ø 288 999	41.7 44.4 41.3 99.9	4.75 4.24 4.43 9.99	88 1Ø5 93 999	14.8 14.6 13.6 99.9	Ø.Ø 999.9 999.9 999.9
892 896 9Ø9 911	17900	99999 4374 99999 10561	99999 162 9999 1253	99999 2511 9999 537Ø	9999 486 9999 537	9999 162 9999 Ø	999 Ø 999 179	999 322 999 433	99.9 41.2 99.9 36.6	9.99 4.47 9.99 4.Ø2	999 92 999 91	99.9 13.9 99.9 13.2	999.9 999.9 999.9 999.9
914 917 919 92Ø 922	87ØØ 99999 46ØØ 65ØØ 99999	522Ø 99999 2254 24Ø5 99999	174 9999 184 520 9999	2262 9999 1978 2665 9999	174 9999 138 455 9999	694 9999 46 455 9999	174 999 Ø Ø 999	298 999 247 313 999	41.2 99.9 44.Ø 45.3 99.9	4.64 9.99 5.Ø8 4.63 9.99	89 999 87 98 999	12.7 99.9 15.3 15.5 99.9	999.9 999.9 999.9 999.9 999.9
925 928 931 932	66ØØ 7ØØØ 75ØØ 75ØØ	363Ø 371Ø 39ØØ 39ØØ	33Ø 77Ø Ø 525	2310 2240 3000 2400	132 28Ø 45Ø 225	198 7Ø 15Ø 45Ø	0 0 Ø Ø	351 351 3Ø1 196	39.2 31.0 48.8 40.8	4.44 3.33 5.28 4.58	88 93 92 9Ø	13.2 10.5 16.5 13.4	999.9 999.9 999.9 999.9
934 938 939 942 943	8000 7600 9300 6400 8500	424Ø 4712 5673 32ØØ 3485	32Ø 38Ø 93 32Ø 11Ø5	28ØØ 1976 2697 23Ø4	32Ø 3Ø4 93 128	32Ø 228 279 448 85	Ø Ø Ø Ø	33Ø 263 248 294 355	42.4 37.2 44.6 34.Ø	4.83 4.26 4.78 3.37	88 87 93 1Ø1	14.4 12.3 15.4 11.4	999.9 Ø.Ø 999.9 37.1
943 944 95Ø 955 956	8700 8700 99999 9600 7000	5742 579999 4992 4410	435 9999 192 210	3315 1827 9999 2496 231ø	51Ø 435 9999 288 7Ø	261 9999 384 7Ø	0 8 999 Ø Ø	363 999 236 3Ø2	46.3 44.4 99.9 44.9 39.Ø	4.93 4.94 9.99 4.9Ø 3.98	94 9Ø 999 91 98	16.Ø 15.2 99.9 13.3 12.6	999.9 Ø.Ø 999.9 999.9 999.9
958 96Ø 962 963	8900 12300 99999 8200	4539 6765 99999 4264	178 492 9999 656	3649 369Ø 9999 2Ø5Ø	177 738 9999 82	267 615 9999 738	89 Ø 999 164	374 323 999 2 9 9	42.6 41.1 99.9 47.6	4.42 4.75 9.99 4.9Ø	96 86 999 97	13.3 13.Ø 99.9 15.9	999.9 999.9 999.9 999.9 999.9
965 966 969 97Ø 971	8900 5500 14900 12000 7400	5073 2850 8344 6840 3108	178 275 594 1Ø8Ø 296	2937 88Ø 5513 276Ø 34Ø4	356 11Ø 298 72Ø 518	356 33Ø 149 6ØØ 74	ช 55 ช ช ช	4Ø2 138 336 4Ø1 348	38.4 41.Ø 47.6 39.7 5Ø.9	4.33 4.22 4.64 4.32 5.55	89 97 1Ø3 92 92	13.3 13.8 15.1 12.6 15.8	999.9 999.9 999.9 999.9 999.9 999.9
975 977 978 98Ø	999999 999999 99 999 65ØØ	999999 99999 999999 351Ø	9999 9999 9999 13Ø	9999 9999 9999 2210	9999 9999 9999 26Ø	9999 9999 9999 26Ø	999 999 999 13Ø	999 999 999 274	99.9 99.9 99.9 44.5	9.99 9.99 9.99 4.89	999 999 999 999 91	99.9 99.9 99.9 14.5	999.9 999.9 999.9 Ø.Ø
981 991	74ØØ 999999	4292 99999	518 9999	1628 9999	444 9999	592 9999	0 999	212 999	49.1 99.9	4.97 9.99	99 999	16.8	999.9 999.9

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	нст	RBC	MCV	HGB	TSH
993	99999		9999	9999	9999	9999	999	999	99.9	9.99	999		999.9
998	96ØØ	7ØØ8	96	192Ø	192	384	Ø	223	46.1	5.15	9Ø	14.7	
1001	73ØØ	365Ø	365	2628	438	219	ø	287	40.5	4.66	87		
1005	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999		999.9
1007	6500	377Ø	130	221Ø	195	195	ø	315	4Ø.9	4.40	93	13.8	6.9
1Ø35	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	599	99.9	999.9
1036	8100	4050	162	3321	486	81	ø	222	51.4	5.88	87		999.9
1043	6600	3366	132	264Ø	198	264 44Ø	រ ស	386	44.6	4.99 4.33	89 95	14.2	999.9 999.9
1050	11000	6Ø5Ø	110	3740	660		Ø	424 19Ø	42.3				
1500	91ØØ 99999	5369	364 9999	3Ø94 9999	182 9999	91 9999	999	999	4Ø.7 99.9	4.55 9.99	89 999		999.9 999.9
15Ø5 1517	99999		9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
1517	6900	4140	207	1587	414	552	999 Ø	216	45.8	4.91	93		999.9
1520	8700	5481	174	2523	522	352 Ø	ø	336	45.Ø	5.16	90 90		999.9
1520	10100	4444	303	4646	505	202	ø	374	53.Ø	5.5Ø	96		999.9
1525	7600	418Ø	76	3116	76	228	õ	351	42.1	4.42	95		999.9
1526	999999	999999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
1533	99999		9999	9999	9999	9999	999	999	99.9	9.99	999		999.9
1541	7600	3952	228	2660	456	3Ø4	้ตี	381	42.1	4.54	93		999.9
1542	87 <i>@</i> Ø	3828	261	4002	522	87	õ	251	48.5	5.85	83	16.1	999.9
1546	99999		9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
1548	12300	6027	984	3Ø75	615	1722	ø	213	42.5	4.73	9Ø		999.9
1549	8700	522Ø	174	2262	174	694	174	298	41.2	4.64	89	12.7	
155Ø	9000	576Ø	18Ø	243Ø	36Ø	18Ø	9Ø	262	43.9	4.68	94	14.7	999.9
1552	58ØØ	174Ø	116	1972	348	464	ø	274	51.1	5.73	89	15.8	999.9
1553	99999		9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
1554	71ØØ	4544	284	17Ø4	142	284	142	248	43.8	4.9Ø	89	13.5	999.9
1555	99999		9999	9999	9999	9999	999	999	99.9	9.99	999		999.9
1556	75ØØ	45ØØ	375	195Ø	375	225	75	3Ø1	4Ø.2	3.92	1Ø3		
1558	6900	4002	2Ø7	1932	483	2Ø7	69	337	31.6	3.77	84	10.8	ø.ø
	15100		9Ø6	1963	Ø	6Ø4	Ø	325	47.Ø	5.47	86	14.6	999.9
156Ø	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999		999.9
1561	8700	6177	261	1827	261	87	Ø	312	42.5	4.38	98	13.5	999.9
1562	99999		9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
1563	6700	2948	268	2077	4Ø2	1005	ø	45Ø	43.8	4.64	94		999.9
1564	6800	2720	68	3332	272	340	68	351	41.6	4.47	93	13.5	2.5
1565	8600	3698	43Ø	3268	516	6Ø2	86	27Ø	51.7	4.93	105		999.9
1566	99999 99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
1567 1568			9999 9999	9999 9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
1569	999999	99999	9999	9999 9999	9999 9999	9999 9999	999 999	999 999	99.9 99.9	9.99 9.99	999 999		999.9 999.9
1570	999999		9999	9999	9999	9999	999	999	99.9 99.9	9.99	999		999.9
1570	99999		9999	9999	9999	9999	999	999 999	99.9 99.9	9.99	999	99.9 99.9	999.9
1572	7100	2527	355	284Ø	639	639	595 Ø	298	54.7	5.93	92		999.9
1577	8600	6364	344	1548	172	035 Ø	ø	275	34.7	5.53 4.ØØ	91 91		999.9
1578	9200	4784	276	3128	46Ø	46Ø	92 [°]	285	48.8	5.56	88	15.7	
2102	10100	5454	202	4141	3ø3	Ψ.	ี้ ดี	404	55.3	5.97	93	17.Ø	Ø.Ø
2103	96øø	7200	384	1536	96	192	õ	316	43.8	4.54	95	15.Ø	ø.ø
2104	5000	245Ø	25Ø	2000	2	5ø	5 ø	25Ø	40.9	4.38	93	13.2	2.9
	10200	6528	51Ø	2346	3Ø6	51Ø	ø	5ø3	4Ø.5	4.64	87	14.2	ø.ø

.

.

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	нст	RBC	MCV	HGB	ŤSH
	12400	558Ø	124	52Ø8	496	868	124	212	46.8	5.26	89	16.1	999.9
	13000	754Ø	1300	351Ø	65Ø	ø	ø	191	47.Ø	5.23	9Ø	14.8	Ø.Ø
21Ø8	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
211Ø	79ØØ	45Ø3	395	2212	237	395	Ø	385	4Ø.1	4.00	100	13.9	3.7
2111	76ØØ	342Ø	.76	3496	38Ø	328	ø	342	38.6	4.87	79	12.6	ø.ø
2113	9800	4410	392	2058	196	2744	ø	261	41.4	5.15	8Ø	14.3	Ø.Ø
2114	6900	3933	2Ø7	2139	276	345 9999	Ø 999	211 999	44.2 99.9	4.95 9.99	89 999	14.9 99.9	999.9 999.9
2115 2117	99999 11100	99999 6771	9999 666	9999 3441	9 999 111	111	999 Ø	363	46.4	5.Ø9	91	15.8	2.8
2119	8700	4002	348	3480	174	696	ø	325	44.2	4.73	92	14.2	999.9
2120	999999	999999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
2123	6400	4032	64	2112	ø	192	ø	151	42.6	4.51	94	14.7	Ø.Ø
2124			9999	9999	9999	9999	9 99	999	99.9	9.99	999	99.9	999.9
2125	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
2126	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
2128	10300	6 Ø77	515	2884	515	3Ø9	ø	234	33.6	4.11	82	11.3	2.6
2129	64ØØ	3136	ø	2432	384	128	128	363	39.Ø	5.Ø1	78	13.5	ø.ø
213Ø	75ØØ	4200	225	2175	45Ø	675	ø	271	36.9	4.18	88	12.8	ø.ø
2132	3500	1575	175	1 5 Ø5	175	7Ø	ø	155	22.1	2.33	95	7.9	ø.ø
2134	7400	3552	444	2516	444	444	ø	337	43.8	4.88	9Ø	14.7	Ø.Ø
2135	99999	99999	9999	9999	9999	9999	999	999 25 a	99.9	9.99	999		999.9 999.9
2136 2137	76ØØ 680C	3192 2584	152 2Ø4	3192 3128	456 4Ø8	6Ø8 476	Ø Ø	35Ø 352	47.9 45.3	5.Ø5 4.96	95 91	$15.5 \\ 14.8$	999.9 Ø.Ø
2137	7100	4118	284	1988	426	639	Ð	226	38.5	4.35	89	12.8	Ø.Ø
2139	12500	6625	25Ø	4625	420 5ØØ	375	125	301	4Ø.Ø	4.30	93	13.5	Ø.Ø
214Ø	5100	2958	102	1683	153	1.02	102	213	39.Ø	4.24	92	12.8	3.5
2142	3øøø	4500	45ø	351Ø	27Ø	27ø	ī	249	51.3	5.35	96	15.5	ø.ø
2143	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
2144	92ØØ	4416	552	3312	552	368	ø	249	51.3	5.21	98	17.6	ø.ø
2145	8500	3481	ø	4335	425	17Ø	85	331	42.4	4.41	96	13.7	ø.ø
2146	99999	99999	9999	99 99	9999	9999	999	9 99	99.9	9.99	999	99.9	999.9
2147	65ØØ	3Ø55	65	273Ø	39Ø	32Ø	ø	42Ø	45.7	4.99	92	15.Ø	ø.ø
2148	9200	5336	276	2852	552	184	ø	142	39.3	4.26	92	13.4	2.6
2149	68ØØ	3536	136	2788	272	68	ø	318	35.1	3.75	94	12.1	ø.ø
215Ø 2152	99ØØ 68ØØ	6237 36Ø4	297 68	297Ø 2924	198 68	198	ø	294	48.9	5.84	84	16.7	Ø.Ø
2152	6800	4488	2Ø4	1Ø88	136	136 984	ช ฮ	32Ø 336	45.Ø 46.Ø	4.93 5.53	91 83	14.Ø 15.Ø	Ø.Ø 4.7
2155	8200	4100	82	2132	574	123Ø	82	278	40.0	5.46	91	16.5	ø.ø
2155	6400	2752	192	2752	64	64	Ø	246	49.9	5.17	97	16.5	ø.ø
2157	10806	6004	Ĩø	4212	756	108	õ	229	44.4	4.83	92	15.7	ø.ø
2158	7100	3479	142	2769	284	426	õ	448	39.9	4.36	92	13.4	ø.ø
2159	7500	4125	300	2400	300	375	õ	449	46.1	5.07	91	15.2	ø.ø
216Ø	6200	2976	248	1984	372	62Ø	õ	385	41.8	4.62	9ø	14.1	9.9
2161	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
2162	13300	9177	133	2926	399	532	133	313	36.9	4.31	86	12.3	3.1
2164	89ØØ	445Ø	178	3471	267	534	ø	385	43.7	4.65	94	14.8	ø.ø
2165	13700	8494	137	4 11Ø	411	44.1	ø	363	5Ø.7	5.74	88	16.5	ø.ø
2166	9600	4512	96	3936	96	96Ø	ø	342	43.3	4.76	91	14.6	4.1
2167	97ØØ	6595	485	2522	97	Ø	ø	315	45.4	5.08	89	15.6	Ø.Ø
2168	67ØØ	3953	134	2144	335	134	ø	236	45.3	4.65	97	15.5	ø.ø

.

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	НСТ	RBC	MCV	HGB	тѕн
2171	85ØC	425Ø	425	3400	255	17Ø	ø	2Ø8	40.2	4.40	91	13.6	Ø.Ø
2172	77ØØ	4081	308	2772	385	154	Ø	335	42.3	3.82	88	13.9	Ø.Ø
2174	8600	5504	258	2Ø64	172	602	ø	260	46.7	5.19	9Ø	16.4	Ø.Ø
2176	9100	4277	91	4186	364	91	91	233	46.1	4.91	94	15.6	Ø.Ø
2179	127ØØ 58ØØ	6731 3Ø74	1Ø16 232	3683 1972	381 116	762 4Ø6	127 Ø	351 298	53.Ø 36.6	6.28 3.95	84 93	18.1 12.Ø	Ø.Ø 3.8
2182 2185	5000 9500	3074 4940	232 95	3895	475	19Ø	ø	219	43.3	4.21	1Ø3	12.0	3.0 Ø.Ø
2185	64ØØ	3328	25 Ø	2688	256	64	64	208	43.5	5.59	92	17.3	ø.ø
2189	11000	858Ø	77ø	660	22Ø	66Ø	Ø	524	38.2	4.31	89	13.5	ø.ø
2193	7400	4292	74	2516	37Ø	148	õ	276	39.1	4.20	93	14.Ø	2.8
2194	6200	3038	248	2666	186	62	õ	211	34.6	3.99	87	10.8	58.7
2195	7700	4Ø81	ø	3003	462	154	ø	423	39.7	4.64	86	14.3	ø.ø
2196	79ØØ	4740	474	2Ø54	79	553	ø	222	40.0	4.51	89	13.2	ø.ø
2197	7ØØØ	392Ø	ø	245Ø	14Ø	28Ø	ø	248	34.9	3.86	9Ø	12.2	4.4
22ØØ	6700	3752	67	2412	42Ø	67	ø	238	4 Ø.1	4.26	94	13.5	2.5
22Ø5	11000	7378	44Ø	253Ø	44Ø	220	ø	298	44.Ø	5.16	85	15.4	ø.ø
22Ø6	85ØØ	425Ø	34Ø	3315	51Ø	85	ø	298	45.8	4.97	92	16.Ø	ø.ø
2207	7400	296Ø	444	3478	222	296	Ø	221	46.7	5.54	86	15.4	2.7
2208	10700	5457	428	2675	642	1391	107	337	4Ø.8	4.33	94	13.7	3.2
2209	99999	99999 2646	9999 54	9999	9999 27Ø	9999 324	999 54	999 236	99.9	9.99	999	99.9	999.9 2.5
221Ø 2212	54ØØ 79ØØ	2646 316Ø	54 79	2Ø52 3792	270 79	324	54 79	236	4Ø.1 39.5	4.38 4.32	92 91	$13.9 \\ 13.4$	2.5 Ø.Ø
2212	7500 9100	5187	79 91	273Ø	455	637	Ø	286	40.2	4.32	92	13.4 13.5	Ø.Ø
2215	8500	357Ø	85	3825	425	595	ŝ	311	40.2	4.42	84	13.4	Ø.Ø
2216	11000	693Ø	ø	286Ø	660	55Ø	õ	423	40.8	4.64	88	14.3	ø.ø
2217	8800	5808	4 4 ø	2376	176	Ĩ	õ	237	46.9	4.89	96	14.2	ø.ø
2218	13600	748Ø	952	4488	4Ø8	272	ã	237	42.2	4.78	88	14.7	3.6
222Ø	77ØØ	4389	385	2233	3Ø8	385	ø	292	39.8	4.25	94	13.8	3.5
2221	6100	3294	488	1952	183	183	ø	242	39.5	4.22	94	13.4	7.5
2224	6000	336Ø	12Ø	198Ø	6Ø	48Ø	ø	323	37.6	3.97	95	12.8	ø.ø
2225	99ØØ	5742	198	2871	297	693	ø	3Ø1	36.3	4.21	86	12.1	3.8
2226	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999		999.9
2227	12700	9098	254	254Ø	ø	508	Ø	243	32.5	3.65	89	11.1	ø.ø
2228	10700	5136	321	4815	321	107	ø	416	36.4	4.06	9Ø	13.2	Ø.Ø
2229	77ØØ 77ØØ	5467 4ØØ4	231 231	1463 2849	231 231	308	Ø	375	43.6	4.74	92	14.1	6.Ø
223Ø 2231	8500	4675	231 17Ø	2049 3060	255	385 34ø	ø	437 999	48.9 4Ø.7	5.73 4.62	85 88	15.8 14.2	Ø.Ø Ø.Ø
2232	8300	3237	498	3984	332	249	ø	231	40.7	4.62 5.19	95	16.9	11.4
2233	8600	5762	344	2064	344	86	ø	286	49.3	5.35	92	17.1	Ø.Ø
2234	10700	6206	535	3317	642	ğ	õ	327	42.9	4.79	89	15.3	3.3
2235	7200	1872	216	4600	288	1 4 4	7 2 T	23Ø	46.6	4.98	94		999.9
2236	6800	3264	ā	3060	408	68	ġ	276	45.7	5.27	87	15.8	4.4
2237	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
2239	6800	4556	68	1428	2Ø4	544	ø	251	42.Ø	4.68	9Ø	13.5	ø.ø
224Ø	99999	99999	9999	99 99	9999	9999	999	999	99.9	9.99	999	99.9	999.9
2241	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	999.9
2242	57ØØ	3249	226	171Ø	171	342	Ø	276	47.2	5.00	94	15.8	ø.ø
2244	4600	1518	46	2438	276	276	46	249	43.8	4.56	96	14.1	ø.ø
. 2245	99999	99999	9999	9999	9999	9999	999	999	99.9	9.99	999	99.9	
2247	12600	7812	378	2898	63Ø	882	ø	363	32.9	3.71	89	11.7	ø.ø

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	нст	RBC	MCV	HGB	тѕн
2248 2249 2250 2251 2254 2255 2256 2260 2261 2268 2269 2269 2269 2267 2273 2273	6600 99999 7600 5200 9600 6700 5200 9700 8000 7800 10100 9700 99999 6500	4Ø92 99999 34380 3952 3432 5280 3618 2132 3880 3680 3744 8181 3880 99999 24Ø5	132 9999 152 228 288 268 312 291 24Ø 784 4Ø4 9999 Ø	1188 9999 2204 2052 1451 3072 2546 22546 2880 3120 1414 4850 999 3705	462 9999 152 312 268 312 268 312 268 312 268 312 488 468 101 485 9999 195	726 9999 684 1Ø64 768 2Ø8 2Ø8 291 72Ø 312 291 9999 195	Ø 9992 152 Ø Ø Ø Ø 999 Ø	284 9974 33283 283 2453 453 4554 2561 999 287	39.3 950.9 41.9 45.2 41.7 45.4 47.4 53.7 539.9 499.9 46.9	4.41 9.99 5.20 4.70 5.33 5.30 5.30 5.30 5.30 5.30 5.30 5.20 5.21	89 999 81 75 88 884 95 889 992 999 999 999	13.8 14.Ø 15.4 14.6 17.4 16.8 18.1 17.Ø 99.9 15.5	Ø.Ø 999.9 Ø.Ø 999.9 2.7 Ø.Ø Ø.Ø 2.8 Ø.Ø 2.8 Ø.Ø 3.4 999.9 Ø.Ø
2276 2277	84ØØ 8ØØØ	4368 536Ø	168 32Ø	3Ø24 16ØØ	336 16Ø	252 48Ø	0 8Ø	236 333	47.9 31.3	5.1Ø 4.72	94 66	16.4 8.9	Ø.Ø Ø.Ø

.

.

																			E A	
IDN	W8C	PMN	BND	LYM	MON	EOS	BAS	PLT	нст	RBC	MCV	HGB	TSH	PRL	HBS	AHBS	AHBC	HDL	сно	TRI
1	6600	2574	ø	3234	33Ø	396	66	22Ø	35.6	3.88	92	12.7	Ø.Ø	999.9	ø	1	1	46.0	134.0	66.Ø
2	9700	5044	194	3589	485	388	ğ	263	48.4	4.98	97	15.1	8.8	999.9	õ	ĝ	ġ		125.0	66.07
3	999999	99999	9999	99999	9999	9999	9999	999	99.9	9.99	999	99.9	999.9	999.9	ø	ø	1	99.9	999.9	999.9
- 4	6100	2562	ø	3172	122	244	ø	346	5ø.7	5.61	9Ø	15.8	Ø.Ø		ø	1	1			257.Ø
5	9800	6762	392	1862	294	392	98	25Ø	45.9	4.71	97	15.1	30.0		ø	ø	ø		164.Ø	73.07
6	4400	2200	88	1496	308	176	ø	161	41.2	4.58	90	14.3		999.9	Ø	1	1		142.0	
7	7200	4536	ø	2016	504	144	ø	191	40.0	4.11	97	13.5		999.9	Ø	1	ø		155.0	65.0
8 9	8600 8200	5848 4674	86 164	215Ø 246Ø	86 328	344 492	86 82	362 16Ø	41.4	4.69 4.6Ø	88 93	12.0	5.9 999.9	999. 9	ย ศ	ទ ទ	1		186.Ø 999.9	79.0
10	8700	5394	174	2697	174	174	87	174	50.6	5.71	89	16.2		999.9	Ø	1	1		183.Ø	
11	4600	2530	138	1564	138	230	Ŕ	231	28.2	2.83	100	10.0		999.9	ø	ģ	1		151.Ø	
12	67.00	3417	201	2680	268	134	õ	387	49.3	5.16	96	14.3		999.9	õ	õ	1		198.0	
14	6300	3465	63	22.05	315	252	ø	178	38.2	3.78	101	13.2		999.9	ø	ø	ī		167.0	63.0
15	10000	6300	ø	3100	580	100	ø	355	42.5	4.59	93	13.4	4.0	999.9	ø	1	1	99.9	999.9	99.9
16	13200	10296	264	2244	132	264	ø	363	45.Ø	5.82	77	13.9		999.9	ø	ø	1		135.Ø	74.Ø
17	9700	5432	ø	3686	291	291	ø	375	43.7	5.Ø4	87	13.4		999.9	ø	ø	1		124.0	44.0
18	6900	4347	276	1863	276	138	Ø	275	39.0	4.21	90	13.2		999.9	ø	ø	1		161.0	
19	5400	3456	216	1404	108	216	ø	374	45.Ø	5.72	79	14.6	Ø.Ø		ø	ø	<i>8</i> A		156.0	510.0
20	10400	7384	104	1768	2Ø8 54	936	<i>ឲ</i> ស្រែន	263 185	51.5 4Ø.3	5.55 4.31	93 91	16.Ø 13.4	0.0	999.9 999.9	97 67	Ø	9		136.Ø 141.Ø	71.Ø 36.Ø
21	54ØØ 54ØØ	378Ø 2592	54 Ø	1296 2592	216	1Ø8 Ø	1.08 Ø	389	40.3	4.31	100	13.4	Ø.Ø		ø	1	1		194.0	36.0 111.0
22 23	999999	99999	9999		9999	9999	9999	999	99.9	9.99	999	99.9	999.9	999.9	ø	à	1		999.9	
24	5800	2900	232	2030	348	290	Ø	291	41.Ø	4.21	97	13.2		999.9	ã	ĩ	i		170.0	93.Ø
27	11100	6438	Ĩø	4218	333	111	õ	237	48.7	4.76	102	16.Ø	0.0		õ	ī	ī			102.0
32	999999	99999	9999		9999	9999	9999	999	99.9	9.99	999	99.9	999.9	999.9	ø	1	1	99.9	999.9	999.9
33	8300	4399	83	2656	332	664	166	3Ø2	41.4	4.56	91	12.8	40.0	999.9	ø	1	1	34.Ø	176.Ø	131.0
34	6700	2211	134	3886	134	335	ø	281	39.3	3.64	1Ø8	12.8	Ø.Ø		ø	1	1		232.Ø	350.0
36	999999	99999	9999		9999	9999	9999	999	99.9	9.99	999	99.9	999.9		ø	1	1		999.9	
37	5900	3068	ø	1829	118	826	ø	225	42.0	4.31	97	13.0		999.9	1	Ø	1		110.0	42.0
39	6700	3417	ø	2680	335	268	ø	574	42.8	4.33	99	13.5	3.6	999.9	ø	1	1		183.Ø 999.9	110.0
40	6200	3224	124	2604	124	124	េ ស	395 166	46.3 42.9	4.79 4.42	97 97	14.3 14.Ø		999.9 999.9	រា ស	Ø	1		143.0	999.9 65.0
41 42	65ØØ 73ØØ	3835 4Ø15	13Ø 73	2275 2263	13Ø 219	13Ø 73Ø	ø	229	42.3	4.22	1.03	13.8		999.9	ต	ģ	Å		103.0	87.0
42	5100	3060	1.072	1734	1.02	102	õ	203	48.2	5.70	85	15.0		999.9	ตี	ĩ	ĩ		135.0	52.0
45	5200	2808	ΞŰΖ	1872	208	260	5 2	298	38.7	3.93	98	12.5	ต.ต		ã	i	i		207.0	153.0
48	5800	3074	58	2262	174	232	ø	182	39.2	4.Ø1	98	13.2	2.7		ø	ø	ø	30.0	138.Ø	62.0
49	8900	3916	267	3827	534	356	ø	224	48.9	5.4Ø	91	13.7	Ø.Ø	999. 9	ø	1	1	30.0	213.Ø	269.Ø
53	7400	4144	ø	2442	592	222	ø	326	43.2	4.65	93	13.9	Ø.Ø		ø	I	1		17Ø.Ø	95.Ø
61	8800	3784	ø	2816	352	88	ø	229	46.6	5.1Ø	91	14.9	7.3		ø	1	ø		2Ø7.Ø	
63	7400	444Ø	296	222Ø	37Ø	296	ø	298	45.7	4.73	97	14.2		999.9	Ø	1	Ø		191.Ø	71.0
64	999999	99999	9999		9999	9999	9999	999	99.9	9.99	999	99.9	999.9	999.9	ø	1	1		999.9	
65	6100	2562	61	1403	183	1769	122	214 229	39.0	3.84	1Ø2 95	11.7 13.Ø		999.9	9 0	1	1		202.0	
66	93ØØ 78ØØ	4185 3822	0 234	4185 312Ø	372 468	465 156	ø ø	229	38.7 42.3	4.Ø7 4.32	98	13.0		999.9	D D	1	1		999.9	
67 69	9999999	3822	234 9999		9999	9999	9999	200 999	99.9	9.99	999	99.9	999.9		Ø	1	à		999.9	
7Ø	47.00	3243	47	1128	188	94	Ø	164	39.Ø	4.28	91	12.6		999.9	õ	ø	ĩ		137.0	74.0
71	14600	7446	584	5986	438	146	õ	266	44.3	4.71	94	13.8	999.9		Ø	ĩ	ø	9.9	99.9	99.9
72	8800	5984	ø	2112	352	352	ø	331	41.2	4.42	93	13.Ø	Ø.Ø	999.9	1	ø	1	3Ø.Ø	153.Ø	17 4 .Ø
73	6700	3953	268	2077	268	134	ø	275	47.5	5.Ø4	94	14.2	ø.ø		ø	1	1			159.0
74	10200	54£76	3ø6	3468	612	306	102	274	46.4	5.Ø8	9 i	15.2	Ø.Ø	999.9	ø	ø	ø	30.0	144.Ø	93.Ø

.

46

PAGE 1

.

PAGE 2

		D MAN		1																
IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	HCT	RBC	MCV	HGB	TSH	PRL	HBS	AHBS	AHBC	HDL	СНО	TRI
75	9000	531Ø	ø	2430	18Ø	1Ø8Ø	ø	239	47.1	4.97	95	14.1	10 0	999.9	ø	1	1	75 Ø	176.Ø	95.Ø
76	6100	2989	183	2501	183	244	õ	237	44.4	4.46	100	14.3	3.3		้ตั	â	ġ		169.0	
77	12000	9480	ø	1920	480	120	ø	233	41.5	4.36	95	13.6		993.9	õ	ĩ	ĩ		157.0	62.0
78	6300	3024	63	2898	126	189	ø	453	39.3	4.00	98	13.1	0.0	999.9	ø	ø	ø	24.0	196.0	100.0
79	7300	3723	219	2774	219	365	ø	162	48.4	4.97	97	15.8	Ø.Ø		ø	1	1		162.0	77.Ø
8Ø	999999	99999	9999		9999	9999	9999	999	99.9	9.99	999	99.9	999.9		ø	1	1		999.9	
81	7200	5184	144	1440	216	216	ø	208	44.1	4.84	91	13.0		999.9	ø	ø	1		159.0	
83 84	52ØØ 999999	27Ø4 99999	ø 9999	2184 99999	2Ø8 9999	1Ø4 9999	ø 9999	3Ø1 999	48.2 99.9	5.ØØ 9.99	97 999	16.Ø 99.9	Ø.Ø 999.9	11.1	រា ស	1	1		19Ø.Ø 999.9	
85	97.00	4074	291	3977	582	679	97	324	47.8	5.09	94	15.2	999.9 Ø.Ø		ต	Ø 1	1		199.0	
86	6300	4158	, Ø	1701	126	315	ø	328	40.3	4.44	91	12.9	3.2		ø	1	i		140.0	87.0
805	6000	2640	120	2460	420	360	ø	349	41.3	4.71	88	12.3	0.0	5.0	ĩ	ġ	1		155.0	
811	75ØØ	3600	300	2925	375	225	75	276	41.4	4.22	98	13.7	0.0	4.8	ø	ø	ø	36.0	164.0	93.Ø
813	8900	4005	89	4894	356	356	ø	248	47.2	4.81	98	16.1	0.0	8.0	ø	1	1			232.0
815	6600	3630	ø	264Ø	132	198	Ø	239	46.6	5.08	92	15.3	ø.ø	7.3	ø	ø	ø		174.0	
816	8000	3760	160	2880	160	CØØ	16Ø	263	40.4	4.42	91	12.8	Ø.Ø		ø	ø	1		167.0	44.0
818	7600	2964	304	3724	228	380	Ø	464 248	45.8	5.07	9Ø	14.8	999.9	999.9	9	9	9		999.9	
821 822	64ØØ 6ØØØ	3648 348Ø	51Z Ø	1792 228Ø	384 180	64 6Ø	ø	3.05	38.8 46.4	4.16	93 94	12.7	999.9 Ø.Ø	999.9 4.6	ø ø	Ø	ø		151.Ø 169.Ø	37.0
823	8400	3948	ø	2436	84	1848	84	249	47.9	4.95	99	15.3	Ø.Ø	4.5	ต	1	1		133.0	
825	8400	4956	252	2940	168	84	ø	374	40.4	4.91	82	14.0	ø.ø	15.8	ต์	1	i		139.0	87.0
826	5100	2703	102	1683	357	255	õ	245	41.0	4.39	90	12.8	3.Ø	0.1	õ	1	î		139.0	87.0
827	10300	5562	2Ø6	2987	309	1236	ø	284	46.3	4.71	98	14.3	Ø.Ø	5.8	ø	1	1		166.0	355.0
829	5900	3186	118	2419	118	59	ø	261	41.9	4.38	96	12.6	Ø.Ø	7.6	ø	1	1	32.0	151.Ø	166.0
830	5400	3078	7.02	1350	54	108	108	2Ø1	43.6	4.48	97	14.5	ø.ø	3.1	ø	1	1		166.0	
831	8500	3400	17Ø	3655	34Ø	85Ø	85	306	56.9	5.90	96	16.7	Ø.Ø	11.1	1	ø	1		190.0	
832	7400	4144	296	2442	222	296	ø	279	38.1	4.91	78	13.6	0.0	55.6	ø	1	1		203.0	95.0
833	5100	27Ø3	1.02	2091	1.02	102	ø	287	48.6	5.65	86	15.2	Ø.Ø	7.2	ø	ø	1		173.0	
834 835	83ØØ 95ØØ	3735 57ØØ	Ø 95	3818 2945	332	415 285	Ø	29Ø 289	41.9	5.ØØ 4.8Ø	84 99	15.3	Ø.Ø Ø.Ø	12.6	ย ย	1	-1		184.Ø 136.Ø	66.0
838	9500	5320	38Ø	3040	190	570	õ	286	57.2	5.83	98	18.1	Ø.Ø	2.4	1	ø	1		128.0	
841	7900	4740	ø	2212	316	553	7 9	275	39.1	4.21	93	12.7	Ø.Ø	23.2	1	ã	i		217.0	
842	6700	3752	õ	2278	335	335	ø	158	45.3	4.64	98	14.3	Ø.Ø	4.4	ø	ĩ	ī		124.0	57.0
843	9200	5612	276	2576	368	368	ø	273	39.1	3.94	99	12.7	Ø.Ø	4.4	ø	1	1	30.0	134.0	137.0
844	4600	2070	138	1978	368	46	ø	295	35.5	4.10	87	12.4	Ø.Ø	11.3	ø	1	1		193.Ø	
845	7900	4108	ø	3239	316	237	ø	211	42.6	4,46	96	13.7	Ø.Ø	13.0	1	ø	1		207.0	
846	5800	3190	290	1798	29Ø	232	ø	300	41.1	4.38	94	12.8	2.5	7.6	ø	1	1		203.0	
851	6100	3233	183	2074	183	427	ø	239	37.6	3.77	100	12.5	Ø.Ø 999.9	9.8	Ø	1	1 9		231.Ø 999.9	96.Ø 999.9
863	84ØØ 75ØØ	4116 2625	252 3ØØ	3696 375ø	336 3øø	ø 525	Ø	257 227	47.5	5.15 4.84	92 87	13.5	999.9	999.9 999.9	9	9	9		999.9	
864 865	5900	2478	3.00 Ø	295Ø	59	413	ø	249	42.5	4.53	96	14.4	Ø.Ø	9.2	ĝ	ø	Ø		174.8	99.Ø
867	9100	4732	364	3458	182	364	ø	334	51.9	5.47	95	17.1	ø.ø	5.3	ต	1	1		212.0	
879	7700	4004	ø	2926	539	231	õ	413	42.7	4.77	9ø	13.3	ø.ø	15.0	õ	i	ī		149.0	86.0
881	6300	3276	ø	252Ø	378	126	ø	184	45.5	4.83	94	14.8	0.0	5.4	ĩ	ø	ĩ		203.0	
882	4900	2450	98	1911	147	196	98	224	52.1	5.83	89	14.7	Ø.Ø	7.4	ø	ø	1		174.0	
883	9400	2444	94	4888	470	15Ø4	ø	348	44.9	4.40	1Ø2	14.4	3.6	6.8	ø	1	1		167.Ø	59.Ø
888	7000	3570	140	2660	210	35Ø	ø	264	43.Ø	4.90	88	14.3	999.9	999.9	9	9	. 9		999.9	
891	6400	4096	256	1536	256	256	Ø	192	41.8	4.19	100	13.6	Ø.Ø	999.9	1	ø	1			140.0
896	8200	4182	82	2788	492	574 348	82	201	38.3	4.54	84	14.1	Ø.Ø Ø.Ø	7.0	Ø	1	Ø		219.Ø 148.Ø	148.0
9ø9	87 <i>00</i>	4372	ø	3219	261	340	ø	228	38.4	4.63	83	12.0	0.0	7.1	2	1	1	410.10	140.0	02.0

PAGE 3

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	НСТ	RBC	MCV	HGB	TSH	PRL	HBS	AHBS	AHBC	HDL	СНО	TRI
911	89ØØ	5785	89	178Ø	445	8Ø1	ø	4Ø4	36.Ø	4.12	87	12.8	Ø.Ø	999 .9	ø	1	1	44.Ø	153.0	77.Ø
917	6200	3844	124	186Ø	124	248	ø	231	41.8	4.95	84	12.7	Ø.Ø	13.4	ø	1	1	28.Ø	193.Ø	184.Ø
919	14500	9135	435	435Ø	58Ø	ø	ø	25Ø	49.7	5.59	89	15.2	Ø.Ø	10.7	ø	1	ø		151.Ø	87.Ø
92Ø	6100	2867	122	2806	<i>9</i>	305	ø	181	46.4	5.02	92	14.6	999.9	999.9	9	9	9		999.9	
922 925	6500 7000	299Ø 364Ø	65 Ø	3Ø55 273Ø	325 14Ø	65 49Ø	Ø Ø	28Ø 381	46.9 39.2	5.18 4.61	91 85	14.0	999.9 Ø.Ø	8.1 16.6	ต ต	1	1		2Ø8.Ø 145.Ø	43.0
926	5400	2592	រេទទ	2484	108	108	ø	355	40.2	4.71	85	13.6		999.9	a a	i	1		158.0	43.0 81.0
928	6900	3933	345	2001	552	138	õ	288	37.6	4.09	92	11.5	ø.ø	19.9	้ต์	1	1		173.0	90.0
931	85ØØ	4420	ø	374Ø	340	ø	ø	375	49.7	5.29	94	16.2	Ø.Ø	10.4	Ĩ	ø	ø		139.Ø	
932	7200	468Ø	72	1872	216	36Ø	ø	244	36.6	3.71	99	11.8	Ø.Ø	999.9	ø	1	ø	40.0	214.0	240.0
934	6200	2666	31Ø	2294	372	3107	ø	335	45.2	5.26	86	14.7	ø.ø	8.2	ø	ø	ø		295.Ø	
938	7300	4891	146	1533	219	438	73	222	4ø.ø	4.56	88	13.Ø	ø.ø	11.0	Ø	1	1		155.0	60.0
939	10300	4635	ø	4944	824	103	ø	218	45.7	4.84	94	14.8	Ø.Ø	6.7	ø	1	1		203.0	
941	95 <i>00</i> 68 <i>00</i>	5415 3264	19Ø 136	3325 2312	475 272	95 68Ø	0 0	241 342	4Ø.7 38.1	4.3Ø 3.8Ø	95 1øø	13.4	999.9 999.9	16.2 2Ø.5	0 8	1	1		197.Ø 198.Ø	
942 943	11000	5264 671Ø	136 Ø	2312 385Ø	220	220	ø	238	53.Ø	5.54	96	16.6	999.9 Ø.Ø	20.5	ы 1	ġ	1		179.0	
944	9900	4752	396	4158	198	396	ã	258	46.1	5.17	89	14.8	ø.ø	8.3	ģ	ã	ġ		195.Ø	
95Ø	9400	4888	188	3760	94	47Ø	ø	448	44.2	4.82	92	14.3	ø.ø	7.1	ĝ	õ	õ		201.0	
955	6800	4Ø12	136	2108	34Ø	2Ø4	ø	249	47.7	5.01	95	13.1	Ø.Ø	5.4	ø	1	ø	40.0	212.0	79.Ø
956	7300	4453	ø	2555	219	73	ø	379	4Ø.7	4.27	95	12.3	ø.ø	12.9	ø	1	1		207.0	
958	5300	2067	ø	2756	212	265	ø	243	38.1	4.35	88	12.0	999.9	999.9	9	9	9		999.9	
959	15000	10000	ø	3150	600	45Ø	Ø	251	45.5	4.84	94	14.8	Ø.Ø	11.4	ø	ø	1		212.0	79.Ø
960	7000	371Ø 4644	28Ø 1Ø8	2100 5076	21Ø 756	56Ø 216	14Ø Ø	336 146	33.8 4Ø.1	4.30	79 94	12.1	Ø.Ø 999.9	999.9 999.9	ø	Ø	1 9		187.Ø 999.9	84.Ø 999.9
963 965	10800 12200	9028	1586	854	366	366	Ø	651	38.9	4.28	91	13.2		999.9	9	9	9		167.0	79.0
966	4700	2209	47	1974	141	282	47	205	45.4	4.59	99	13.9	Ø.Ø	8.9	ø	ĩ	1		191.Ø	95.Ø
969	8000	3680	320	3760	8Ø	80	8.0	240	44.5	4.60	97	14.8	999.9	999.9	ĩ	ġ	9		999.9	999.9
97Ø	11400	7182	114	3078	57Ø	342	114	239	33.7	3.57	94	10.7	Ø.Ø	6.4	ø	i	i		165.Ø	88.0
971	6800	3672	2Ø4	2312	4Ø8	204	ø	312	45.2	4.92	92	14.3	ø.ø	999.9	ø	1	1	24.Ø	131.Ø	23Ø.Ø
975	59ØØ	3835	Ø	1534	236	295	ø	133	46.1	5.18	89	15.1	ø.ø	3.5	ø	ø	1		151.Ø	
977	14900	8791	149	4917	447	596	ø	300	47.1	5.24	90	15.4	4.5	8.0	ø	1	1		149.0	56.Ø
98Ø	6000	2880	120	252Ø	240	240	ទ ស	192	42.2	4.60	92	13.4	Ø.Ø	999.9	ø	ø	1		155.0	44.0
981	89ØØ 63ØØ	64Ø8 3Ø2 4	ថ ខ	2225 2583	267 126	Ø 5Ø4	63	253 287	47.6 43.2	5.Ø8 4.85	94 89	16.1	Ø.Ø Ø.Ø	9.Ø 19.Ø	រា ស	1 1	л р		149.Ø 120.Ø	103.0 52.0
993 998	8500	4335	85	3145	255	340	85	240	41.9	4.59	91	14.0	Ø.Ø	999.9	Ø	1	1		201.0	52.0 75.0
1001	6000	4440	24Ø	1140	120	6.0	อัต	342	41.5	4.92	84	13.3	ğ.ğ	4.5	ตี	i	1		150.0	
1007	5900	3953	Ĩø	1711	177	59	õ	233	42.2	4.51	94	13.6	Ø.Ø	999.9	õ	i	ī		222.0	
1035	9200	5520	ø	2944	46Ø	276	ø	348	46.8	5.43	86	14.7	0.0	8.9	Ø	Ĩ	ĩ		173.Ø	
1043	9200	6716	ø	2024	276	184	ø	24Ø	43.1	5.Ø1	86	13.6	Ø.Ø	6.8	ø	ø	ø		174.Ø	5ø.ø
1050	9100	4459	182	2912	182	1365	ø	348	35.3	4.Ø2	88	12.8	Ø.Ø	11.Ø	ø	1	1		218.0	
1500	5800	3306	58	1914	29Ø	116	116	352	38.2	4.14	92	12.5	Ø.Ø	6.6	ø	1	1		187.0	75.0
1505	6000	2340	Ø	3Ø6Ø 255Ø	360	24Ø 75	Ø	298 28Ø	4Ø.8 47.4	4.29	95 95	13.8	Ø.Ø Ø.Ø	3.7 8.2	ø ø	1 1	1		179.Ø	
1519	75ØØ 64ØØ	465Ø 4224	Ø 64	2550	225 192	128	ы Ø	365	4/.4	4.97 5.21	89	15.2	999.9	9.9	ø	a na	1		217.Ø	
152Ø 1524	9200	5060	Ø	3956	92	92	ø	210	48.3	5.02	96	16.3	2.9	8.4	ดี	ĩ	i		182.0	
1524	6400	3840	64	1536	384	512	64	228	42.1	4.33	97	13.2	ø.ø	8.0	õ	ø	ø		123.0	84.0
1526	8300	4399	83	2988	166	498	166	255	42.6	4.9Ø	87	15.3	999.9	999.9	- 9	9	9		999.9	
1529	10800	5616	ø	4536	432	216	ø	248	53.Ø	5.99	88	16.9	Ø.Ø	11.3	ø	1	1		2Ø7.Ø	
1530	8800	6512	4 4 Ø	1056	4 4 Ø	352	ø	381	45.6	4.89	93	14.3	ø.ø	5.3	Ø	1	1		196.Ø	
1541	6900	4071	138	2277	138	345	ø	262	35.7	4.14	86	13.0	2.5	14.1	ø	1	1	28.0	190.0	247.0

 $\frac{43}{8}$

.

.

1

PAGE 4

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	НСТ	RBC	MCV	HGB	TSH	PRL	HBS	AHBS	AHBC	HDL	сно	TRI	
1542	7600	4Ø28	ø	3344	76	152	ø	324	44.8	5.33	84	15.6	ø.ø	5.1	ø	1	1	30.0	202.0	312.0	
1546	7800	3978	156	312Ø	312	234	ø	146	51.8	5.58	93		999.9		9	9	9	99.9	999.9		
1548	9200	5336	184	276Ø	552	368	° Ø	337	40.9	4.56	9Ø	13.8	Ø.Ø		1	ø	1		137.0	65.0	
1549	7400	4292	148	2516	222	222	ດ ø	175	48.5	5.10	94	14.6	Ø.Ø	7.5	ø	1	1		192.0	76.0	
155Ø 1552	73ØØ 7ØØØ	3431 392Ø	73 14Ø	3139 238Ø	292 42Ø	365 56Ø	ю Ø	411 357	45.4	4.83 5.Ø1	94 89	15.5	Ø.Ø 999.9	11.1	Ø	1 9	1 9	3Ø.Ø 99.9	198.Ø 999.9	225.0	
1552	6100	3538	244	1952	183	183	ø	271	43.2	4.27	101	13.9	995.9 Ø.Ø	11.4	ø	1	1		16Ø.Ø		
1555	10800	6588	g	3348	432	432	ã	236	51.1	6.35	80	15.4	ø.ø	8.8	õ	ġ	i	38.Ø	194.0		
1556	4400	176Ø	ø	2288	88	22Ø	44	281	43.Ø	4.33	99	13.3	999.9	16.7	Ø	ø	1	99.9	999.9	999.9	
1558	6200	2418	62	2666	372	682	ø	323	40.3	4.33	90	13.6	Ø.Ø	24.2	ø	1	1		161.Ø	7Ø.Ø	
1559	7000	4620	ø	1610	35Ø	420	Ø	236	42.8	4.85	88	12.1	Ø.Ø	993.9	ø	1	1	56.Ø	246.0		
1564	7200	3456	ø	2808	360	576	ø	290	43.1	4.89	88	13.3	13.0	22.8	ø	1	1		136.0	66.0	
1565 1567	82ØØ 56ØØ	5412 2128	. 82 . Ø	2Ø5Ø 2128	164 168	41Ø 1176	82 Ø	237 299	53.9 41.7	5.44 4.27	99 91	16.6	Ø.Ø Ø.Ø	8.2 34.1	Ø Ø	1	1	26.0	161.Ø 126.Ø	149.Ø 37.Ø	
1567 157Ø	10500	4200	ទ	5670	210	420	ฮ	299	41.7	4.27	91	14.8	Ø.Ø	Ø.Ø	Ø	1	1		260.0		
1572	7000	2870	õ	392Ø	149	7.0	õ	225	51.4	5.77	89	16.4	ø.ø	7.3	ã	ģ	i	26.0	130.0	75.Ø	
1573	6400	3200	25Ĝ	2496	192	192	64	202	51.3	5.41	95	16.8	Ø.Ø	6.7	ĩ	ø	ī	24.Ø		34Ø.Ø	
1577	12600	8316	756	2772	378	378	ø	351	43.Ø	4.57	94	14.1	Ø.Ø	34.7	ø	1	ø	44.Ø	157.Ø	44.Ø	
21Ø2	9200	6348	ø	2300	46Ø	92	ø	341	48.9	5.Ø4	97		999.9	999.9	ø	1	1	99.9	999.9		
2103	6400	3712	128	1984	32Ø	256	ø	222	43.6	4.30	101		999.9		1	ø	1		161.0		
2104	6700	4891	201	1139	201	268	Ø	330	37.5	3.90	96		999.9 999.9		1	ย ต	1	42.0			
21Ø5 21Ø6	10300 13500	5974 7155	ย ฮ	2575 58Ø5	618 4Ø5	1133 135	ø	425 232	44.8 47.4	4.72 5.20	95 91		999.9		1 Ø	1	1	28.Ø	160.0		
21.07	16300	9128	រ	5053	489	1467	163	252	46.0	4.84	95		999.9		1	ต่	ต่	40.0		229.0	
2108	5900	2183	118	3068	295	236	Ĩø	244	43.1	4.90	88		999.9		ġ	ê	- 9	99.9	999.9		
2110	8700	4872	261	2958	435	174	ø	285	40.9	4.04	1Ø1		999.9	999.9	ø	1	1	28.Ø	228.Ø	319.00	
2111	9000	4500	9Ø	324Ø	45Ø	72Ø	ø	316	42.8	4.95	86		999.9		1	ø	1	28.Ø	155.Ø		
2113	97ØØ	6208	ø	3007	388	97	ø	33Ø	44.5	5.67	78-				ø	1	ø		189.0		
2114	7200	5256	216	1152	144	36Ø	72	172	45.1	4.91	92		999.9		ø	1	1		210.0		
2117	10300 6500	6489 37ø5	2Ø6 65	3Ø9Ø 195Ø	2Ø6 325	3Ø9 455	Ø	312 298	46.3 46.8	4.96 5.Ø5	93 93		999.9 999.9		1 Ø	Ø 1	1	30.0 20 0	18Ø.Ø 167.Ø	420.0 90.0	
2119 2123	9000	5040	27Ø	2520	54Ø	455	9ø	186	40.0	4.86	97		999.9		ø	1	1		146.0		
2124	10000	5000	200	3400	300	300	Ĩø	271	53.5	5.93	9.0	16.5		999.9	ã	î	i	20.0	195.Ø		
2125	7200	4248	283	1656	36Ø	648	ø	374	48.4	5.02	96	15.8		999.9	ø	i	ĩ	26.Ø	232.0		
2126	7600	4028	ø	3040	228	3Ø4	ø	324	42.6	4.51	94		999.9		ø	1	1	46.Ø		9Ø.Ø	
2128	95ØØ	6080	57Ø	247Ø	95	285	ø	348	31.Ø	3.72	83		999.9		ø	1	1	24.0			
2129	8400	4536	84	2688	588	420	84	313	42.6	5.30	80		999.9		1	ø	ø		260.0		
2130	5400	3240	108	1620	1.08	324	Ø	253	42.5	4.47	95		999.9	999.9	ø	1	1	32.0	137.Ø 12Ø.Ø	53.Ø 52.Ø	
2132 2134	41 <i>00</i> 999999	2583 99999	41 9999	1189 99999	164 9999	123 9999	9999	2Ø1 999	41.7 99.9	4.8 <i>0</i> / 9.99	87 999	13.2	999.9		Ø	1	1		999.9		
2134	6500	377Ø	325	20/15	325	65	8	322	51.1	5.38	95		999.9		1	Ŕ	1	26.0	153.0	72.0	
2137	8900	3204	89	4984	356	267	õ	240	43.8	4.83	91				ī	õ	ø		999.9		
2138	9200	6Ø72	ø	2392	46Ø	276	ø	385	39.1	4.32	91	12.9	999.9	999.9	ø	1	ø		204.0	86.Ø	
2139	6900	3864	69	2346	414	2Ø7	ø	278	44.Ø	4.67	94		999.9		1	ø	1	38.Ø			
2140	7900	3713	79	2923	553	553	79	228	38.8	4.15	93		999.9		Ø	1	1		999.9		
2142	9800	5978	392	2842	490	98	ø	200	51.1	5.23	98		999.9		1	ø	1		198.0		
2143	9400	5922	94	2068	188	1128 34Ø	Ø	313	51.9	5.62	92 9Ø	15.3	999.9 Ø.Ø	999.9 Ø.Ø	រា ស	1	1		113.Ø 999.9		
2144	85ØØ 92ØØ	476Ø 4324	51Ø Ø	28Ø5 368Ø	85 368	34Ø 828	ю Ø	288 438	49.5 43.1	5.53 4.47	90 96		999.9	ط.ط 999.9	10 10	1	I Ø		190.0		
2145	9999999			999999	9999	9999	9999	430 999	43.1 99.9	9.99	999		999.9		ø	1	1		999.9		
2147				22225						2.22	222				~	•	-				

- 0	Δ		

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	нст	RBC	MCV	HGB	TSH	PRL	HBS	AHBS	AHBC	HDL	СНО	TRI
2148	6500	2795	13Ø	2795	455	13Ø	ø	200	44.3	4.77	93	13.9	999.9	999 9	ø	1	,	22 a	165.0	173 a
2149	7200	3816	216	2808	144	216	ø	274	41.6	4.53	92	12.3		999.9	ĝ	1	i		176.0	
2150	7500	4200	75	2400	525	300	ø	315	49.6	5.82	85	16.7		999.9	ĩ	ġ	i		188.0	
2152	7600	3300	ø	3268	38Ø	152	ø	318	48.Ø	4.84	99	15.9	0.0	999.9	ī	ø	1		19Ø.Ø	
2153	9999999	99999		99999	999 9	9999	9999	999	99.9	9.99	999			999.9	ø	1	1	99.9	999.9	999.9
2155	8200	5166	ø	27Ø6	246	82	ø	363	52.Ø	5.82	89		999. 9		ø	ø	1		154.Ø	
2156	6000	2520	60	2400	120	9ØØ	Ø	235	49.4	5.20	95		999.9		1	ø	1		192.Ø	
2157	9999999 7200	3744	9999 36Ø	99999 2ø16	9999 36Ø	9999 648	9999 72	999 249	99.9 4Ø.8	9.99 4.47	999 91		999.9		្រ ឆ	ย ย	Ø 1		999.9	
2158 2159	11000	8360	220	2010	110	220	ĝ	225	47.8	5.37	89		999.9		ø	1	1		174.Ø 166.Ø	
2160	8700	5307	2 L Ø	2610	348	435	õ	289	44.6	4.72	94	14.5		999.9	ø	à	1		133.Ø	
	9999999			999999	9999	9999	9999	999	99.9	9.99	999		999.9		ã	ĩ	ģ		999.9	
	9999999			999999	9999	9999	9999	999	99.9	9.99	999		999.9		ĩ	ø	ĩ		999.9	
2164	7900	3792	79	3239	553	237	ø	328	38.2	4.14	92	12.3	999.9	999.9	ø	ø	1	38.Ø	188.0	155.0
2165	16300	863 9	652	5868	652	489	ø	214	49.5	5.63	88		999.9		ø	ø	1		3Ø6.Ø	
2166	7100	2769	142	3337	568	284	ø	244	44.5	4.58	97		999.9		Ø	1	ø		164.0	
2167	12500	6250	ø 9999	4250	500	1500	Ø	226 999	51.6 99.9	5.47 9.99	94 999	17.0	<i>ط. ط</i>	999.9	ø	ø ø	1		128.0	
2168 2171	999999 9800	99999 588Ø	9999	99999 2744	9999 294	99 99 784	9999 Ø	234	45.1	4 .76	95	13.4		999.9	1 Ø	1 .	1		999.9 167.Ø	
2172	6100	3843	122	1891	183	61	. ø	326	40.0	4.53	88		999.9		ดี	1	1		167.0	
2174	11000	8140	Ĩø	2200	440	220	õ	325	44.6	5.05	88		999.9		ĩ	ø	i		200.0	
2176	76ØØ	3648	ø	3116	532	152	ø	298	49.Ø	5.15	95		999.9		ø	1	ī		178.0	
2179	85ØØ	493Ø	255	3Ø6Ø	17Ø	17Ø	ø	335	51.1	6.13	83		999.9		ø	1	1		110.0	
2182	7700	5159	154	2002	308	77	ø	285	36.4	3.94	92		999.9		1	ø	1		183.Ø	
2188	6400	3264	ø	2624	192	320	ø	227	51.9	5.45	95		999.9		ø	1	1		194.0	
2189	8800	5280	176	1936	352	968	88 Ø	4Ø1 325	35.2 43.4	3.80	93	10.8	Ø.Ø	Ø.Ø 999.9	Ø	ខ ព	រ ឆ		999.9	
2193 2194	8ØØØ 9999999	528Ø 99999	8Ø 9999	2 <i>000</i> 99999	48Ø 9999	16Ø 9999	9999	325 999	43.4	4.7Ø 9.99	92 999	13.2	999.9		ต ต	1	10 I		219.Ø 999.9	
2194	8100	3645	81	3483	243	648	Ø	348	42.2	4.95	85	13.4		999.9	ø	ġ	1		235.Ø	
2196	7200	3672	ě	3168	144	216	õ	363	40.3	4.48	9 <i>ø</i>	13.4		999.9	ĩ	õ	i			222.0
2197	6600	3234	66	27.06	198	396	ø	428	38.4	4,32	89		999.9		ø	1	1			98.0
2200	6500	351Ø	13Ø	24.05	26Ø	195	ø	215	41.1	4.49	92	13.2	999.9	999.9	ø	1	1	24.Ø	195.Ø	123.Ø
22Ø5	8700	4437	174	3915	87	87	ø	266	46.6	5.37	87		999.9		ø	1	1			283.0
2206	7800	46Ø2	78	2496	468	156	ø	212	45.4	5.04	90		999.9		1	Ø	1		218.0	96.0
2207	8500	4930	Ø	2890	510	170	រា ស	252	47.2 43.4	5.38	88 93		999.9	999.9	ø	1	1		160.0	
2208	95ØØ 10100	6555 5656	285 2Ø2	1995 3131	285 1 <i>0</i> 1	38Ø 9Ø9	101	285 285	43.4	4.65	91	14.1 13.Ø		999.9	Ø	1	1		211.Ø 143.Ø	68.0
22Ø9 221Ø	8500	4930	85	3145	170	170	I D I	341	44.7	4.9Ø	91		999.9		ø	ø	i		129.Ø	41.0
2212	76ØØ	5092	ğ	1596	3.04	6.08	ã	287	42.5	4.57	93	13.Ø		999.9	õ	ĩ	î		243.0	
2213	999999		9999		9999	9999	9999	999	99.9	9.99	999		999.9	999.9	Ø	ī	ĩ			999.9
2215	10000	5000	200	3600	7ØØ	400	100	391	44.8	5.16	87	14.7		999.9	1	ø	i		219.Ø	
2216	114 00	684Ø	ø	2964	456	1140	ø	374	42.6	4.96	86		999.9		ø	1	1		201.0	
2217	6400	3648	128	1856	128	640	ø	259	41.7	4.29	97	13.4	999.9		ø	1	1			215.Ø
2218	9800	4018	294	4410	392	588	98	319	48.9	5.77	85	14.4	0.0	0.0	1	Ø	1		999.9	
2220	82 <i>00</i> 7800	451Ø 4524	82 546	246Ø 1872	328 234	738 624	82 Ø	333 3Ø4	41.2 37.2	4.33 4.25	95 83	12.4	999.9 Ø.Ø	999.9 Ø.Ø	Ø Ø	1	1		212.Ø 999.9	
2221 2224	7800 6800	4524	136	2244	68	68	ø	336	36.3	4.25	94		999.9		ø	ឆ	Ø		244.0	
	999999	999999	9999	999999	9999	9999	9999	999	99.9	9.99	999			999.9	ø	1	1		999.9	
2226	5800	3422	116	2030	232	116	ø	28Ø	39.8	5.26	76	12.8	Ø.Ø	Ø.Ø	ĩ	ø	ī		999.9	
	999999			99999	9999	9999	9999	999	99.9	9.99	999	99.9	999.9	999.9	ø	1	ø		999.9	999.9

-

PAGE 6

.

IDN	WBC	PMN	BND	LYM	MON	EOS	BAS	PLT	нст	RBC	MCV	HGB	TSH	PRL	HBS	AHBS	AHBC	HDL	СНО	TRI
2228	12200	6954	366	3172	122	1586	ø	295	33.3	3.66	91	11.7	999.9	999.9	ø	1	1	38.0	191.Ø	147.0
2229	8200	5576	246	18Ø4	419	82	ø	255	44.9	4.62	97	13.8		999.9	ā	ī	ī		156.Ø	42.0
223Ø	11100	7104	ø	3219	333	444	ø	3Ø1	52.Ø	5.91	88		999.9		ø	i	ġ		259.Ø	
2231	6600	3498	66	2178	33Ø	528	ø	211	46.5	5.42	86	13.8	999.9	999.9	Ø	ø	- 1	30.0	215.Ø	620.0
2232	96ØØ	384Ø	96	4896	576	192	ø	256	56.5	5.72	98	18.1	3.3	999.9	ø	ĩ	ĩ	30.0	199.Ø	212.0
2233	9700	5529	ø	3783	388	ø	ø	249	52.2	5.6Ø	93	16.2	999.9	999.9	ø	1	ī	24.0	130.0	185.0
2234	6400	4352	ø	1536	256	256	ø	2ø5	45.7	5.16	89	15.1	999.9	999.9	ø	1	1	30.0	128.0	163.Ø
2235	8400	4032	84	252Ø	5.04	126Ø	ø	324	43.5	4.64	94	14.1	999.9	999.9	ø	1	1	26.Ø	184.0	
2236	9300	5952	ø	2697	372	279	ø	342	45.1	5.22	86	15.3	3.1	999.9	ø	1	1	3Ø.Ø	167.Ø	112.0
2239	8500	425Ø	ø	357Ø	255	425	ø	263	40.3	9.99	999	99.9	ø.ø	999.9	ø	1	1	38.Ø	135.Ø	47.Ø
2242	7100	355Ø	71	22Ø1	355	852	71	327	47.Ø	4.81	98	15.Ø	Ø.Ø	999.9	ø	1	1	32.00	149.0	52.Ø
2244	6800	2516	136	3876	136	136	ø	2Ø4	4Ø.8	4.2Ø	97	13.4	999.9	999. 9	Ø	ø	1	38.Ø	203.0	59.Ø
2245	8200	4756	246	27ø6	246	246	ø	275	5Ø.5	5.26	96	15.8	ø.ø	999.9	ø	1	1	36.Ø	155.Ø	176.0
2247	7200	3Ø24	ø	2952	432	72Ø	72	236	38.9	4.35	89	13.Ø	999.9	999.9	1	ø	1	32.Ø	160.0	124.00
2248	6400	3712	128	2176	64	32Ø	ø	400	42.6	4.95	86	14.4	ø.ø	999.9	ø	1	1	36.0	157.0	180.0
2249	999999	99999	9999	99999	9999	9999	9999	999	99.9	9.99	999	99.9	999.9	999.9	ø	ø	1	99.9	999.9	999.9
225Ø	8000	392Ø	ø	3040	48Ø	56Ø	ø	400	49.3	5.31	93	15.5	999.9	999.9	ø	1	1	32.Ø	151.Ø	215.Ø
2251	9100	5915	182	2184	364	273	182	413	40.4	4.75	85		999. 9		ø	1	1	26.Ø	155.Ø	149.Ø
2254	4800	2736	48	168Ø	ø	336	ø	425	43.8	5.12	84	12.8	999.9		ø	1	ø		2Ø5.Ø	59.Ø
2255	78ØØ	3666	ø	3354	234	546	ø	202	49.Ø	5.37	91	14.5		999.9	1	ø	1		192.Ø	
2256	7700	4389	77	3Ø8Ø	77	77	ø	445	42.7	4.82	89		999.9		ø	1	ø		181.Ø	84.Ø
2257	4600	2622	92	17Ø2	184	184	ø	244	45.5	5.17	88		999.9		1	ø	1		231.Ø	
2260	8600	3440/	86	3698	344	946	86	383	43.7	4,93	89	14.8	Ø.Ø		ø	1	i		176.Ø	
2261	55 <i>00</i>	2365	ø	264Ø	110	275	11Ø	287	5Ø.5	5.22	97	16.6	ø.ø		ø	1	1	30.0	189.Ø	
2268	77ØØ	4312	ø	2849	231	154	ø	222	5Ø.5	5.51	92				ø	1	1	3Ø.Ø	161.Ø	
2269	8000	432Ø	8Ø	312Ø	400	8Ø	ø	256	49.3	5.Ø8	97	16.Ø			ø	1	1	28.Ø		173.Ø
2271	7600	3496	ø	3952	76	76	Ø	341	48.5	5.34	91		999.9		ø	1	1		151.Ø	
2273	9999999	99999	9999	99999	9999	9999	9999	999	99.9	9.99	999				ø	ø	ø	99.9		
2274	5700	2109	171	3Ø21	171	228	ø	284	46.3	5.45	85		999.9		ø	1	1		156.Ø	
2276	93ØØ	3999	ø	4836	279	186	ø	287	55.Ø	5.86	94		999.9		1	ø	1		139.Ø	
2277	7600	4484	38Ø	2204	228	304	ø	33Ø	36.3	5.42	67	10.6	999.9	999.9	ø	ø	1	20.Ø	136.0	65.0
