

R



UNCLASSIFIED

UWFL-92

ATOLL SOIL TYPES IN RELATION TO THE DISTRIBUTION OF FALLOUT RADIONUCLIDES

ES DOC ARCHIVES	
RG	
Collection	1320
Box	9
Folder	33

August 1965

Laboratory of Radiation Biology  
University of Washington  
Seattle, Washington

HISTORIAN'S OFFICE INVENTORY CONTROL	
Job	1320
Box	9
Folder	33

UNITED STATES ATOMIC ENERGY COMMISSION • DIVISION OF TECHNICAL INFORMATION

UNCLASSIFIED

## LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

This report has been reproduced directly from the best available copy.

Printed in USA. Price \$3.00. Available from the Clearinghouse for Federal Scientific and Technical Information, National Bureau of Standards, U. S. Department of Commerce, Springfield, Virginia 22151.

DOE ARCHIVE

ATOLL SOIL TYPES IN RELATION TO THE  
DISTRIBUTION OF FALLOUT RADIONUCLIDES

E.E. Held, S.P. Gessel and R.B. Walker

Laboratory of Radiation Biology  
University of Washington  
Seattle, Washington

Lauren R. Donaldson  
Director

August 1965

Operated by the University of Washington under Contract No.  
AT(45-1)1385 with the United States Atomic Energy Commission.

## ABSTRACT

The redistribution of radionuclides in atoll soils following fallout from a nuclear device is described. The soils are calcareous, containing no inorganic colloids, and their exchange capacity is directly related to organic content. Comparison of gamma-ray spectra of depth increments from young and old soils shows that  $\text{Cs}^{137}$  and  $\text{Sb}^{125}$  move most readily in old soil, while the principal gamma-emitting radionuclide moving in young soil is  $\text{Sb}^{125}$ .  $\text{Sr}^{90}$  moves in both old and new soils, and quantitative differences in vertical movement between soil types is obscured by the highly variable surface distribution of the radionuclides. There is a vertical gradient in the distribution of radionuclides even within the surface inch. Litter redeposits  $\text{Cs}^{137}$  and  $\text{Sr}^{90}$  at the soil surface and bird droppings have added  $\text{Zn}^{65}$  and  $\text{Co}^{60}$ . In young soils the highest levels of radioactivity are associated with soil algae found as a surface crust in undisturbed areas and in coral fragments in eroded areas. Horizontal movement is localized and probably is of little overall importance. Buried organic horizons contain more  $\text{Cs}^{137}$  than adjacent soil layers, and roots are generally more radioactive than the surrounding soil except at the soil surface. Pumice particles in the soil adsorb radionuclides but pumice is found infrequently. Mechanical mixing by animals in old soils and by erosion in young soils

NOT RECORDED

is important in the redistribution of radionuclides near the soil surface.  $Cs^{137}$  and  $Sr^{90}$  are the principal radionuclides entering a cycle within the soil-plant system. Any loss from this system appears to be small, but a definite conclusion can not be drawn from the data.

E. L. HIND

## TABLE OF CONTENTS

	Page
INTRODUCTION	1
METHODS OF COLLECTION AND MEASUREMENT	4
RESULTS AND DISCUSSION	7
Depth Gradient of Gamma Spectra	12
Old Soil	12
Litter and Guano	14
Young Soil	18
Depth Gradient of Sr <sup>90</sup>	21
RADIOAUTOGRAPHS	25
Young Soil	25
Undisturbed	25
Eroded	25
Old Soil	28
Horizontal Movement	28
Buried Organic Layers	30
Roots	30
Pumice	31
CONCLUSIONS	33
SUMMARY	35
REFERENCES	37

DOE ARCHIVES

LIST OF TABLES

<u>Number</u>		<u>Page</u>
I.	Nutrient levels and soil reaction in the surface A <sub>1</sub> horizon of the five most extensive soil series.	5
II.	Sr <sup>90</sup> levels in soil samples from two sets of plots.	6
III.	Composition of representative soil types, from Rongelap Atoll (after Kenady 1962).	11
IV.	Relative Sr <sup>90</sup> content of depth increments collected in 1959 from two Rongelap Atoll soil types.	22
V.	Sr <sup>90</sup> and calcium in the surface 1-1/2 inches of Rongelap soil, Rongelap gravelly sand, collected in September 1959.	24

DOE ARCHIVES

## LIST OF FIGURES

<u>Number</u>		<u>Page</u>
1.	Beach Ridge Sand, Rongelap Island.	3
2.	Kabelle Sand.	8
3.	Gogan series, Kabelle Island.	9
4.	Lomuilal Sand, Kabelle Island.	10
5.	Gamma-ray spectra of equal amounts of old soil from different depths.	13
6.	Gamma-ray spectrum of a 9 to 10-inch depth increment from old soil compared with the gamma spectra of $Sb^{125}$ and $Cs^{137}$ .	15
7.	Gamma-ray spectra of depth increments of old soil taken from the surface inch.	16
8.	Gamma-ray spectra of litter and guano collected in an area of old soil.	17
9.	Gamma-ray spectra of equal amounts of young soil from different depths.	19
10.	Gamma-ray spectrum of young soil from a depth increment of 9 to 10 inches compared with the gamma spectrum of $Sb^{125}$ .	20
11.	Radioautograph obtained by exposing no-screen X-ray film approximately one cm below the soil surface <u>in situ</u> .	23
12.	Radioautographs and photographs of soil core sections.	26
13.	An area of young soil subject to erosion.	27
14.	Photograph and radioautograph of old soil showing correspondence of radioactive portions to organic matter.	29

DOE ARCHIVES



ATOLL SOIL TYPES IN RELATION TO THE  
DISTRIBUTION OF FALLOUT RADIONUCLIDES

INTRODUCTION

The redistribution of radionuclides in atoll soils following contamination with radioactive fallout is the subject of this paper. Rongelap Atoll, northern Marshall Islands, in the central Pacific Ocean, presents a unique opportunity for such studies since it was substantially contaminated with radioactive fallout only once. The fallout resulted from a thermonuclear device detonated at Bikini Atoll eighty miles to the west on March 1, 1954. Although there was some additional contamination from nuclear tests in 1956 and 1958 the total contribution of radionuclides from the fallout of these subsequent test series amounted to a fraction of one per cent of the amount from the 1954 fallout. Gamma radiation dose rates at Rongelap at detonation plus one day ranged from 3.5 r/hr at the southern islets of the atoll to 35 r/hr at the northern islets (Dunning 1957). These rates declined at approximately the rate predicted for mixed fission products by Miller and Loeb (1958).

Rongelap Atoll has a lagoon area of 388 square miles and an average depth of 168 feet (Nugent 1946, p. 748). The emergent land area is about three square miles, consisting of

DOF ARCHIVES

sixty-one small islets ranging in size from a fraction of an acre to the largest, Rongelap, which is four miles long and one-half mile across at its widest point. There is one small islet on the western reef and the other islets extend along the northern, eastern and southern reefs. The islets on the northern reef are not as well developed as those to the east and south. There are two seasons--a dry season from December to March and a wet season from April to November. Annual rainfall is less than fifty inches, and there is no well-developed fresh-water lens. Some important features of Rongelap Atoll including aerial photographs are given by Wiens (1962).

Classification and mapping of the soil types at Rongelap Atoll were reported by Kenady (1962). The parent material is primarily calcium carbonate, originating from corals, foraminifera, coralline algae and mollusk shells. There is a very small amount of pumice drift in the soils. Since these soils contain no inorganic colloids, exchange capacity and organic content are linearly related. In some areas, particularly along the seaward sides of the islets, buried  $A_1$  horizons are found as deep as eighty inches (Fig. 1). These highly organic horizons presumably result from storm debris covering previously established soil and vegetation.

The pH, determined in the field from a 1:1 soil-water ratio with a Beckman Model N-2 pH meter, is generally between




Fig. 1. Beach Ridge Sand, Rongelap Island. This soil type occurs on the island beach ridges. The pit shown above is seven feet deep and contains six buried horizons.

7 and 8 in the surface horizon but occasional values from 4.6 to 9.5 were found. The pH increases with depth and decreasing organic content.

The amounts of exchangeable cations in the different soil types are given in Table I . The calcium content is so high that more calcium is brought into solution with repeated extractions. Consequently, strontium units have little or no significance relative to atoll soils and, when given, are based on total calcium rather than on exchangeable calcium.

#### METHODS OF COLLECTION AND MEASUREMENT

The vertical distribution of radioactivity in soil and litter was studied by analyzing samples taken mostly by 1-inch soil increments, in a few cases by 1/4- and 1/8-inch increments and by radioautographs of sections of soil cores prepared by the method of Held et al. (1965). The increment collections were made during both the wet and dry seasons in 1958, 1959, 1961 and 1963. The cores were collected only in 1963. Because there is considerable horizontal variation in the levels of radioactivity (Table II) each set of increments was collected to insure the sampling of a single vertical column.

The large amount of horizontal variability also made it more profitable to compare the relative amounts of radionuclides at different depths from many profiles than to make precise

Table I. Nutrient levels and soil reaction in the surface A<sub>1</sub> horizon of the five most extensive soil series (from Kenady 1962).

Soil Series	Rongelap Gravelly Sand	Gogan Gravelly Sandy Loam	Lomuialal Sand	Beach Ridge Sand	Kabelle Sand
Per Cent Nitrogen	0.57	1.71	0.26	0.09	0.14
Per Cent Organic Matter	16.7	35.6	6.4	4.5	7.7
*Exchange Capacity	22.2	37.7	12.6	3.7	5.7
*Sodium	3.36	4.01	2.68	1.16	1.52
*Magnesium	4.19	11.06	3.21	2.55	1.92
*Potassium	1.95	1.80	0.79	0.37	0.46
Phosphorus (ppm)	81.7	985.2	54.2	32.8	32.1
pH	8.1	7.8	8.4	8.6	8.6

\* meq per 100 grams of oven dry soil

Table II. Sr<sup>90</sup> levels in soil samples from two sets of plots. Samples collected in September 1961. Values in pCi/gm dry. Each plot is 16-1/2 by 33 feet.

Kabelle Islet, Pit 38 Area																
Depth in inches	Plot Sample	Subsample	A		B		C		D		E		F		G	
			1	2	1	2	1	2	1	2	1	2	1	2	1	2
0-1	"	1	100	110	94	81	220	280	530	82	40	69	71	100	76	62
	"	2	78	100	95	94	240	200	410	130	49	81	49	100	91	73
1-2	"	1	33	62	180	60	150	41	45	61	22	54	30	26	13	29
	"	2	25	51	180	91	160	43	49	78	24	52	29	22	21	22
2-3	"	1	44	50	33	50	65	29	41	49	8	36	19	14	8	17
	"	2	36	52	25	69	79	23	41	51	11	32	25	17	7	21

Rongelap Islet, Pit 22 Area													
Depth in inches	Plot Sample	Subsample	A		B		C		D		Pit 22		
			1	2	1	2	1	2	1	2	1	2	
0-1	"	1	230	230	130	110	180	170	160	180	130	140	
	"	2	230	150	150	170	190	180	150	350	150	200	
1-2	"	1	150	190	84	47	100	53	54	140	77	100	
	"	2	160	180	89	47	95	61	63	140	79	97	
2-3	"	1	63	49	19	6	28	13	25	65	25	43	
	"	2	55	50	18	4	30	12	21	65	19	36	

FOIA ARCHIVES

quantitative determinations from a few profiles. Gamma-ray spectra of equal amounts of soil from different depths were compared directly to arrive at (1) a qualitative evaluation of the vertical distribution of the radionuclides and (2) a semiquantitative estimate of relative amounts of radionuclides at different depths. Gamma-ray spectra were made with a system which included a 3-inch by 3-inch solid, thallium-activated, sodium iodide crystal and a 256-channel analyzer. In addition, selected samples were taken for analyses of the pure beta-emitter,  $\text{Sr}^{90}$ .

#### RESULTS AND DISCUSSION

Vertical distribution in relation to soil type is made by comparing young and old soils. Examples are Kabelle Sand, a young soil, (Fig. 2) and Gogan Gravelly Sandy Loam (Fig. 3) and Lomuial Sand (Fig. 4), old soils. Their characteristics are given in Tables I and III.

The young soil has little organic material except that in an algal surface crust about 1 cm thick. Hermit crabs, Coenobita perlatus, at the base of a few shrubs, the borrowing ghost crab, Ocypode ceratophthalmia, and ants are the main animals present. Litter accumulation is found only at the bases of the scattered shrubs and is a minor part of this soil system.





Fig. 2. Kabelle Sand. A very young soil composed of fine lagoon sands deposited on the lagoon side of Kabelle Island.





Pit 4, Gogan Series, Kabelle Island



1 inch

3

16


26

Micromonolith of Pit 38



Stand of *Pisonia grandis*, Kabelle Island, pit 38 area.


Fig. 3. Gogan series, Kabelle Island. This is a well-developed, productive soil, usually associated with *Pisonia grandis* and *Cordia subcordata*.



Soil pit 12, Kabelle Island. Dark strata are buried horizons



Messerschmidia argenticum at soil pit 12.



Micromonolith  
Pit 21

Fig. 4. Lomuilal Sand, Kabelle Island.

Table III. Composition of representative soil types from Rongelap Atoll (after Kenady 1962).

Kabelle Sand, a young soil			
Soil Depth - inches	0-1	1-11	11-38
Per Cent Material > 2 mm	94	8	2
Per Cent Nitrogen	0.22	0.02	0.01
Per Cent Organic Matter	8.1	--	2.6
* Exchange Capacity	6.3	0.3	0.1
* Sodium	1.57	3.01	1.97
* Magnesium	1.37	1.04	1.16
* Calcium	3.01	2.88	2.65
* Potassium	0.57	0.15	0.20
Phosphorus (ppm)	30.0	12.0	12.0
pH	8.9	9.1	9.2

Lomuilal Sand, an old soil					
Soil Depth - inches	0-3	3-10	10-12	12-21	21-48
Per Cent Material > 2 mm	0	5	18	12	2
Per Cent Nitrogen	0.29	0.07	0.08	0.04	0.02
Per Cent Organic Matter	2.8	2.3	2.2	1.9	1.7
* Exchange Capacity	14.2	2.3	2.6	1.1	0.6
* Sodium	2.73	0.73	0.82	0.82	0.84
* Magnesium	4.66	8.36	1.05	0.85	0.83
* Calcium	5.02	1.87	2.50	2.31	2.35
* Potassium	1.09	0.18	0.19	0.16	0.16
Phosphorus (ppm)	105.9	15.1	14.1	5.0	5.0
pH	8.4	8.6	8.3	8.8	9.1

Gogan Gravelly Sandy Loam, an old soil						
Soil Depth - inches	1-0	0-1	1-5	5-12	12-20	20-26
Per Cent Material > 2 mm	10	20	20	27	39	56
Per Cent Nitrogen	1.54	1.96	0.42	0.18	0.07	0.05
Per Cent Organic Matter	21.4	--	5.9	6.8	2.6	2.6
* Exchange Capacity	20.5	43.6	17.9	7.2	2.6	1.7
* Sodium	2.0	3.0	0.8	0.4	0.4	0.4
* Magnesium	7.0	7.4	4.0	2.2	1.2	1.1
* Calcium	10.3	24.2	14.1	6.2	7.7	7.8
* Potassium	--	1.80	--	--	--	--
Phosphorus (ppm)	1330.4	892.8	415.5	216.3	150.6	25.1
pH	7.4	7.1	7.9	8.2	8.6	8.8

\* meq per 100 gm dry soil

DOE ARCHIVES

In contrast, the old soil has well-developed  $A_0$  and  $A_1$  horizons and supports dense vegetation, which produces a heavy litter fall during the dry season and which contributes to the redistribution of radionuclides in the system.

Sea birds nest in the vegetation and land crabs, predominantly Birgus latro and Coenobita perlatus, burrow in the soil. Earthworms are seldom found and terrestrial isopods, although found in old soils on some islets, are few. There are also few soil insects, which are mainly tenebrionid and carabid beetles as well as ants.

#### Depth Gradient of Gamma Spectra

##### Old Soil

The gamma spectra, with background subtracted, of the 0 to 1-inch, 1 to 2-inch, and 9 to 10-inch depth increments from old soil collected in March 1959 are given in Fig. 5. The spectra of the increments between 2 inches and 9 inches have been omitted from the figure for clarity but show a gradual change from the condition at 1 to 2 inches to that at 9 to 10 inches. The radionuclides corresponding to the photopeaks are indicated in the figure. The photopeaks of  $Co^{60}$ ,  $Zn^{65}$ ,  $Mn^{54}$ ,  $Ce^{144}$ - $Pr^{144}$  and  $Eu^{155}$  show significant counts in the surface increments but are not detectable in samples from greater depths. The amounts of  $Cs^{137}$  and  $Sb^{125}$  decrease and the proportions of  $Cs^{137}$  and




Fig. 5. Gamma-ray spectra of equal amounts of old soil from different depths.

$\text{Sb}^{125}$  change with increasing depth. In the 0.60 to 0.66-mev photopeak region of the 0 to 1-inch increment, the relatively broad peak is a combination of the 0.60-mev photopeak of  $\text{Sb}^{125}$  and the 0.66-mev peak of  $\text{Cs}^{137}$ - $\text{Ba}^{137\text{m}}$ . In the 1 to 2-inch increment the peak becomes sharper and is oriented toward the 0.66-mev photopeak of  $\text{Cs}^{137}$ - $\text{Ba}^{137\text{m}}$ . At the 9 to 10-inch increment there is almost complete orientation toward the  $\text{Cs}^{137}$ - $\text{Ba}^{137\text{m}}$  peak, with little  $\text{Sb}^{125}$  remaining. In Fig. 6 a comparison is given of the spectrum of the 9 to 10-inch increment (Fig. 5) and the gamma spectra of  $\text{Cs}^{137}$  and  $\text{Sb}^{125}$  spikes.

Fig. 7 shows the gamma spectra of increments taken from an undisturbed area in 1958. The first spectrum is from the 0 to 1/4-inch depth and the subsequent spectra are from 1/8-inch depth increments to a depth of 1 inch. The highest levels of  $\text{Sb}^{125}$ , the rare earths, and  $\text{Co}^{60}$ , which move more slowly than  $\text{Cs}^{137}$  or  $\text{Sr}^{90}$ , are in the 1/2 to 5/8-inch increments, whereas the  $\text{Cs}^{137}$ , which moves most rapidly in this soil type, is in the 3/4 to 7/8-inch increment.

#### Litter and Guano

The gamma-ray spectrum of litter, consisting of leaves, twigs and floral parts splattered with tern droppings, collected from old soil in 1961 (Fig. 8) shows the  $\text{Cs}^{137}$  photopeak to be much higher than the  $\text{Sb}^{125}$  peak. The 1.17 peak of  $\text{Co}^{60}$  is skewed to the left, indicating the presence of the 1.12-mev




Fig. 6. Gamma-ray spectrum of a 9 to 10-inch-depth increment from old soil compared with the gamma spectra of Sb<sup>125</sup> and Cs<sup>137</sup>.




Fig. 7. Gamma-ray spectra of depth increments of old soil taken from the surface inch. The origin of each spectrum has been arbitrarily separated from the preceding one by a factor of ten to facilitate inspection of the figure.






Fig. 7. (continued)




Fig. 8. Gamma-ray spectra of litter and guano collected in an area of old soil.

Zn<sup>65</sup> peak. The presence of Zn<sup>65</sup> is corroborated by the 0.5-mev peak. Fig. 8 also shows the gamma-ray spectrum of noddy tern guano collected in this area. The 1.12-mev peak predominates over the 1.17 peak of Co<sup>60</sup> and the 0.51 peak of Zn<sup>65</sup> is evident. The foliar contribution to the litter contains only Cs<sup>137</sup> from among the gamma-emitters.

In undisturbed areas Cs<sup>137</sup> and Sr<sup>90</sup> are being deposited with the litter and are thus replacing at the surface some of the Cs<sup>137</sup> and Sr<sup>90</sup> lost by leaching. There is not sufficient data from the field work to determine whether there eventually will be a loss of these radionuclides from the soil-plant system, or a steady state (excluding physical decay of the radionuclides). Long-term experiments, under simulated field conditions, with monolith lysimeters and controlled and uniform addition of the radionuclides would define this point.

#### Young Soil

Fig. 9 gives the spectra of the 0 to 1-inch, 1 to 2-inch, and 9 to 10-inch increments of a young soil. Co<sup>60</sup>, Zn<sup>65</sup>, Ce<sup>144</sup>, Pr<sup>144</sup>, and Eu<sup>155</sup> were detected only in the surface layers, and with increasing depth the 0.60 to 0.66-mev photopeak region of the spectra shifts toward the 0.60-mev peak of Sb<sup>125</sup>. The spectrum of the 9 to 10-inch increment is compared with that of an Sb<sup>125</sup> spike in Fig. 10, showing that the photopeaks of the soil and spike gamma spectra are identical.




Fig. 9. Gamma-ray spectra of equal amounts of young soil from different depths.




Fig. 10. Gamma-ray spectrum of young soil from a depth increment of 9 to 10 inches compared with the gamma spectrum of  $Sb^{125}$ .

The radionuclide content of leachates collected in the field agrees with these results (Cole et al. 1961). Leachates from young soil contained only  $\text{Sb}^{125}$  and  $\text{Sr}^{90}$ , while leachates from better developed soils contained mainly  $\text{Sr}^{90}$  and  $\text{Cs}^{137}$ , with traces of other gamma-emitters.

#### Depth Gradient of $\text{Sr}^{90}$

The relative  $\text{Sr}^{90}$  content of depth increments of the two soil types is given in Table IV. There is a rapid decrease in the amount of  $\text{Sr}^{90}$  with depth, and the differences in  $\text{Sr}^{90}$  content between soils probably are not significant. The extreme values from the results of  $\text{Sr}^{90}$  analyses of subsamples of replicate samples taken from small areas differ by a factor of more than ten (Table II). It is likely that the variability is due largely to the spotty nature of the distribution of the fallout radionuclides, which is evident in the radioautographs discussed below and from X-ray films that were exposed at the soil surface and just below the surface (Fig. 11), and to small differences in the characteristics of the soil within a single soil type.

Sampling by 1/8-inch increments in 1959 of an undisturbed old soil, (Fig. 4 and Table I) on Rongelap Island indicates a gradient of  $\text{Sr}^{90}$  levels in the top inch of soil (Table V). The levels in the second inch are about one tenth those in the

Table IV. Relative Sr<sup>90</sup> content of depth increments collected in 1959 from two Rongelap Atoll soil types.

Depth in inches	Percent of total in surface ten inches	
	<u>Old Soil</u> (Pit 4)	<u>Young Soil</u> (Pit 6)
0 - 1	81.0	52.9
1 - 2	4.9	16.6
2 - 3	2.8	16.2
3 - 4	2.0	5.8
4 - 5	1.4	4.0
5 - 6	2.2	2.2
6 - 7	2.0	1.2
7 - 8	0.8	0.7
8 - 9	2.5	0.2
9 - 10	0.6	0.1
Total	100.2	99.9



Fig. 11. Radioautograph obtained by exposing no-screen X-ray film approximately one cm below the soil surface in situ.

not available



Table V.  $\text{Sr}^{90}$  and Ca in the surface 1-1/2 inches of Rongelap soil, Rongelap Gravelly Sand, collected in September 1959.

Depth in inches	$\text{Sr}^{90}$ Percent of total in top 1-1/2 inches	Ca mg/gm
0 - 1/8	27	292
1/8 - 1/4	26	291
1/4 - 3/8	16	315
3/8 - 1/2	10	337
3/4 - 7/8	3.2	346
7/8 - 1	2.7	348
1 - 1-1/8	2.0	306
1-1/8 - 1-1/4	2.6	338
1-1/4 - 1-5/8	1.5	329
1-5/8 - 1-3/4	2.2	328
1-3/4 - 1-7/8	3.6	351
1-7/8 - 1-1/2	2.2	369
Total	99.0	

first inch and remain at approximately the same level from 7/8 inch to 1-1/2 inches in depth.

## RADIOAUTOGRAPHS

### Young Soil

#### Undisturbed

Radioautographs of cores taken in 1963 show marked differences in the distribution of radioactivity. Fig. 12 shows photographs and the corresponding radioautographs of sections of cores of young soils from an undisturbed area (Fig. 2) and from an area subject to erosion (Fig. 13) and an old soil (Fig. 3). The radioactivity corresponds closely to the dark area in the photograph of the core from the undisturbed young soil. This dark area is composed almost entirely of a mixture of soil algae, forming a crust which has retained most of the fallout radionuclides.

#### Eroded

In an eroded area of young soil the radioactivity is associated with large coral fragments which are infiltrated with algae. This area is subject to erosion by both wind and water, which accounts for the coral fragments containing algae and radionuclides occurring below the soil surface. Radioactivity is not associated primarily with smaller particles as




Fig. 12. Radioautographs and photographs of soil core sections. From left to right: undisturbed young soil, eroded young soil, old soil.




Fig. 13. An area of young soil subject to erosion.

in other soils, presumably because smaller particles have been eroded away since fallout occurred.

### Old Soil

A radioautograph of a core from the old soil (Fig. 12) shows that the radioactivity extends to a greater depth and is more diffusely distributed than in the young soils. The surface litter contains little radioactivity compared with the soil surface.

### Horizontal Movement

There is evidence of localized horizontal movement of radionuclides in the soil, at least near the surface, which in turn may affect vertical distribution. Fig. 14 is a photograph and radioautograph of a section of a soil core collected in old soil on Kabelle Islet. There is a darkened funnel-shaped area at the top center of the radioautograph and a corresponding darker area of high organic content in the core section. The amount of activity in the stem of the "funnel" appears to be too great to be accounted for by leaching from the surface directly above. Either a depression was filled with radioactive organic matter after fallout, or the depression was filled before fallout and the radionuclides were adsorbed when surface runoff water filtered through the funnel of organic matter. The latter interpretation seems more probable since gamma-ray spectra of

DOF ARCHIVE




Fig. 14. Photograph and radioautograph of old soil showing correspondence of radioactive portions to organic matter.

layers of the funnel show only  $\text{Cs}^{137}$  and  $\text{Sb}^{125}$  in the deepest layer.

### Buried Organic Layers

The organic matter in buried horizons may become important in the vertical distribution of radionuclides in the soils. No detectable gamma-emitters were found in buried horizons during the period 1958-1961. In September 1963 a core was taken through a buried horizon at 14 inches in old soil. Radioautographs of sections of the core show that a small amount of radioactivity has accumulated in the buried horizon. The diffuse distribution of the radioactivity in this case indicates that there is adsorption of the radionuclides from solution as water percolates through the soils, although the possibility of translocation by plants cannot be ruled out.  $\text{Cs}^{137}$  and naturally occurring  $\text{K}^{40}$  were the only radionuclides present in the buried horizon.

The accumulation of  $\text{Cs}^{137}$  in the buried horizons is of particular significance in terms of plant-soil relationships, because a proliferation of roots is often found in the buried horizons and  $\text{Cs}^{137}$  is the principal fallout radionuclide taken up by atoll plants.

### Roots

Roots, which contain  $\text{Cs}^{137}$  and  $\text{Sr}^{90}$ , in most instances have higher levels of radioactivity than the surrounding soil except near the soil surface (Fig. 12). The influence of poplar roots on redistribution in a continental soil has been well demonstrated

by Witkamp and Frank (1963). It may be possible to distinguish between translocation within the roots and downward movement along root channels at Rongelap by comparing the ratios of radio-nuclides within the roots and in the immediately surrounding soil in core sections. For example, if there is greater movement along root channels, we would expect  $Sb^{125}$ , which is not absorbed by the plants, to be most abundant in the soil adjacent to the roots at depth, if there is appreciable channelization.

#### Pumice

Radionuclides are also adsorbed by pumice particles. No detailed morphological examinations of the soil sections have been made, but it is obvious in some core sections that a few of the larger "hot spots" several inches below the surface are associated with pumice fragments. The retention of radionuclides by pumice fragments may be of importance in considering soil-plant relationships in a few highly localized areas since proliferation of roots around pumice fragments has been observed (Sachet 1955; Kenady 1962). However, as pumice is rarely found beneath the surface, the effects of this material would not be generally important.

\* \* \*

We can not explain the differences in distribution of radio-nuclides between soil types but assume that the greater retention

1967 APR 11 11:17 AM



of all radionuclides, except  $\text{Sb}^{125}$  and  $\text{Sr}^{90}$ , at the surface of the young soil is associated with the algal crust. The observation that mosses and lichens collected from trees at Rongelap Atoll in 1961 show essentially the same gamma-ray spectra as the algal crust lends supporting evidence to this assumption. The retention of radionuclides by algal crust, mosses and lichens must be related to adsorptive surfaces or to the metabolism of the organisms, although it is impossible to determine from the field data the mechanism or combination of mechanisms involved. Similar observations have been made with arctic lichens (Palmer et al. 1964).

Since the algal crust at Rongelap Atoll has retained the radionuclides for nine years and from all indications will continue to do so for years to come, it is possible that it also retains a variety of mineral nutrients, thus providing a reservoir of nutrients in otherwise barren areas. This reservoir might be tapped upon invasion of the areas by higher plants and the concomitant activity of animals in mixing the upper layers of the soil.

Distribution of the radionuclides in the old soils arises from a combination of processes which are difficult to delineate. No doubt much of the movement is due to leaching and readsorption

of radionuclides moving through the soil. This is certainly true for  $\text{Cs}^{137}$  and  $\text{Sr}^{90}$ , which were abundant compared to other radionuclides in the leachates. It also was demonstrated by comparing the gamma-ray spectra of depth increments that there is a more rapid movement of  $\text{Cs}^{137}$  than of other gamma-emitters present. It should be recalled that the exchange capacity of these soils originates from the organic content, which, as is obvious from the photographs of the core sections and Table III, is far higher in the old than in the young soil. There is thus more opportunity for exchange and retention of radionuclides in old soil. Mechanical mixing, due mainly to the activity of land crabs, plays an important role in redistribution in the surface layers. This effect is obvious in areas where there has been active burrowing, and is probably occurring to a small extent throughout densely vegetated areas as is indicated by the presence of  $\text{Ce}^{144}$ - $\text{Pr}^{144}$  and  $\text{Eu}^{155}$  in the litter. These radionuclides could only have come directly from the soil by upward mixing since they were not found in the vegetation which contributed significantly to the litter.

#### CONCLUSIONS

Different plant and soil environments on single islets have a different vertical distribution pattern of radionuclides from the same fallout material. The vertical distribution of

radionuclides in old soils is as follows, in order of greatest penetration:  $\text{Cs}^{137}$ ,  $\text{Sr}^{90}$ ,  $\text{Sb}^{125}$ ,  $\text{Co}^{60}$ ,  $\text{Zn}^{65}$ ,  $\text{Cr}^{144}$ - $\text{Pr}^{144}$ ,  $\text{Eu}^{155}$  and probably other rare earths. In the young soils, consisting almost entirely of parent material, the positions of  $\text{Cs}^{137}$  and  $\text{Sb}^{125}$  are reversed and the other radionuclides appear to be more completely retained in the surface algal crust. The maximum concentration of fallout radionuclides remains at the soil surface, a few inches or less in depth, except in areas where there has been erosion. In the eroded areas large particles containing both soil algae and radionuclides are randomly distributed to a depth of a few inches.

There is some horizontal movement of radionuclides but such movement appears to be very localized and thus is of little consequence in the overall picture of distribution.

$\text{Cs}^{137}$  and  $\text{Sr}^{90}$  are the principal radionuclides entering a cycle within the soil-plant system. Any loss from this system appears to be small (a fraction of one per cent per year), but a definite conclusion can not be drawn from the data.

## SUMMARY

1. Rongelap Atoll received a single heavy dose of radioactive fallout in 1954.
2. The atoll soils are calcareous and contain no inorganic colloids; the exchange capacity is related to organic content.
3. Comparison of gamma-ray spectra of depth increments from old and new soils shows that  $\text{Cs}^{137}$  and  $\text{Sb}^{125}$  move most readily in the old soil; the principal gamma-emitting radionuclide moving in new soil is  $\text{Sb}^{125}$ .
4.  $\text{Sr}^{90}$  moves in both old and young soils.
5. The distribution of radionuclides at the surface is very spotty.
6. There is a vertical gradient in the distribution of radionuclides within the surface one-inch layer.
7. Litter redeposits  $\text{Cs}^{137}$  and  $\text{Sr}^{90}$  over the soil surface and bird droppings have added  $\text{Zn}^{65}$  and  $\text{Co}^{60}$ .
8. The principal reservoir of radionuclides in young soils is the surface algal crust.
9. In eroded areas radioactivity is associated with large coral fragments, which are infiltrated with algae.
10. There is some localized horizontal movement of radionuclides in old soils.

DO NOT REPRODUCE

11. Buried organic horizons contain more  $\text{Cs}^{137}$  than adjacent soil layers.
12. Roots are generally more radioactive than the surrounding soil except at the soil surface.
13. Pumice particles in the soil adsorb radionuclides.
14. Mechanical mixing by animals in old soils and by erosion in young soils results in a redistribution of radionuclides in the surface layers.
15.  $\text{Cs}^{137}$  and  $\text{Sr}^{90}$  are the principal radionuclides entering a cycle within the soil-plant system.

## LITERATURE CITED

- Cole, D. W., S. P. Gessel and E. E. Held. 1961. Tension lysimeter studies of ion and moisture movement in glacial till and coral atoll soils. *Soil Sci. Soc. of Am.*, Proc. 25, 4.
- Dunning, G. M. (Ed.). 1957. Radioactive contamination of certain areas in the Pacific Ocean from nuclear tests. U.S. Atomic Energy Commission, Washington, D. C.
- Held, E. E., S. P. Gessel, L. J. Matson and R. F. Billings. Autoradiography of sectioned soil cores. In Proceedings of [IAEA] Symposium on Radioisotope Sample Measurement Techniques in Medicine and Biology, Vienna, May 24-28, 1965. (In press)
- Kenady, R. M. 1962. The soils of Rongelap Atoll, Marshall Islands. M.S. Thesis, University of Washington Library.
- Miller, C. F. and P. Loeb. 1958. Ionization rate and photon pulse rate of fission products from slow neutron fission of  $U^{235}$ . U. S. Naval Radiological Defense Laboratory report USNRDL-TR-247.
- Nugent, L. E., Jr. 1946. Coral reefs in the Gilbert, Marshall, and Caroline Islands. *Geol. Soc. Am.*, Bull. No. 57, 8.
- Palmer, H. E., W. C. Hanson, B. I. Griffin, and D. M. Fleming. 1964. Radioactivity measurements in Alaskan Eskimos in 1963. *Science* 144:1620.
- Sachet, M. H. 1955. Pumice and other extraneous volcanic materials on coral atolls. *Atoll Res. Bull.*, No. 37.
- Wiens, H. J. 1962. Atoll environment and ecology. New Haven, Conn., Yale University Press. 532 p.
- Witkamp, M. and M. L. Frank. 1964. First year of movement, distribution and availability of  $Cs^{137}$  in the forest floor under tagged tulip poplars. *Rad. Botany* 4(4):485-495.